



United States
Department of
Agriculture

In cooperation with
Illinois Agricultural
Experiment Station

Soil Survey of Sangamon County, Illinois



Natural
Resources
Conservation
Service




How To Use This Soil Survey

The **detailed soil maps** can be useful in planning the use and management of small areas.

To find information about your area of interest, locate that area on the **Index to Map Sheets**. Note the number of the map sheet and turn to that sheet.

Locate your area of interest on the map sheet. Note the map unit symbols that are in that area. Turn to the **Contents**, which lists the map units by symbol and name and shows the page where each map unit is described.

The **Contents** shows which table has data on a specific land use for each detailed soil map unit. Also see the **Contents** for sections of this publication that may address your specific needs.



This soil survey is a publication of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (formerly the Soil Conservation Service) has leadership for the Federal part of the National Cooperative Soil Survey.

Major fieldwork for this soil survey was completed in 1999. Soil names and descriptions were approved in 1999. Unless otherwise indicated, statements in this publication refer to conditions in the survey area in 1999. This survey was made cooperatively by the Natural Resources Conservation Service and the Illinois Agricultural Experiment Station. It was partially funded by the Illinois Department of Agriculture and the Sangamon County Soil and Water Conservation District. The survey is part of the technical assistance furnished to the Sangamon County Soil and Water Conservation District.

Soil maps in this survey may be copied without permission. Enlargement of these maps, however, could cause misunderstanding of the detail of mapping. If enlarged, maps do not show the small areas of contrasting soils that could have been shown at a larger scale.

The United States Department of Agriculture (USDA) prohibits discrimination in all of its programs on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice or TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue SW, Washington, DC 20250-9410, or call 202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Cover: A farm pond adjacent to strongly sloping, well drained Hickory and moderately sloping, well drained Navlys soils.

Additional information about the Nation's natural resources is available on the Natural Resources Conservation Service homepage on the World Wide Web. The address is <http://www.nrcs.usda.gov>.

Contents

How To Use This Soil Survey	3	<i>Camden Series</i>	35
Numerical Index to Map Units	9	134A—Camden silt loam, 0 to 2 percent slopes	36
Foreword	11	134B—Camden silt loam, 2 to 5 percent slopes	36
General Nature of the County	13	134C2—Camden silt loam, 5 to 10 percent slopes, eroded	37
History	13	<i>Clarksdale Series</i>	38
Farming and Industry	13	257A—Clarksdale silt loam, 0 to 2 percent slopes	39
Transportation Facilities	15	<i>Cowden Series</i>	40
Physiography and Drainage	15	112A—Cowden silt loam, 0 to 2 percent slopes	42
Climate	15	<i>Denny Series</i>	42
How This Survey Was Made	15	45A—Denny silt loam, 0 to 2 percent slopes	43
Formation and Classification of the Soils	17	<i>Drury Series</i>	44
Formation of the Soils	17	7075B—Drury silt loam, 2 to 5 percent slopes, rarely flooded	45
Factors of Soil Formation	17	536—Dumps, mine	46
Parent Material	17	<i>Edinburg Series</i>	46
Climate	18	249A—Edinburg silty clay loam, 0 to 2 percent slopes	47
Plants and Animals	18	<i>Elburn Series</i>	47
Topography	19	198A—Elburn silt loam, 0 to 2 percent slopes	49
Time	19	<i>Elco Series</i>	49
Processes of Soil Formation	20	119C2—Elco silt loam, 5 to 10 percent slopes, eroded	50
Soils and Soil-Landscape Units	20	119D—Elco silt loam, 10 to 18 percent slopes	51
Classification of the Soils	22	119D2—Elco silt loam, 10 to 18 percent slopes, eroded	51
Soil Series and Detailed Soil Map Units	25	119D3—Elco silty clay loam, 10 to 18 percent slopes, severely eroded	52
<i>Alvin Series</i>	26	<i>Elkhart Series</i>	52
131C2—Alvin fine sandy loam, 5 to 10 percent slopes, eroded	27	567C2—Elkhart silt loam, 5 to 10 percent slopes, eroded	53
131D2—Alvin fine sandy loam, 10 to 18 percent slopes, eroded	27	567D2—Elkhart silt loam, 10 to 18 percent slopes, eroded	54
<i>Assumption Series</i>	27	<i>Fayette Series</i>	54
259C2—Assumption silt loam, 5 to 10 percent slopes, eroded	28	280C2—Fayette silt loam, 5 to 10 percent slopes, eroded	55
259D2—Assumption silt loam, 10 to 18 percent slopes, eroded	29	280D2—Fayette silt loam, 10 to 18 percent slopes, eroded	56
<i>Blackberry Series</i>	29	280D3—Fayette silty clay loam, 10 to 18 percent slopes, severely eroded	56
679B—Blackberry silt loam, 2 to 5 percent slopes	30		
<i>Broadwell Series</i>	31		
684B—Broadwell silt loam, 2 to 5 percent slopes	32		
684C2—Broadwell silt loam, 5 to 10 percent slopes, eroded	32		
<i>Buckhart Series</i>	33		
705A—Buckhart silt loam, 0 to 2 percent slopes	34		
705B—Buckhart silt loam, 2 to 5 percent slopes	34		

<i>Harrison Series</i>	57	630C2—Navlys silt loam, 5 to 10 percent	
127B—Harrison silt loam, 2 to 5 percent		slopes, eroded	77
slopes	58	630D2—Navlys silt loam, 10 to 18 percent	
127C2—Harrison silt loam, 5 to 10 percent		slopes, eroded	78
slopes, eroded	58	630D3—Navlys silty clay loam, 10 to 18	
<i>Hartsburg Series</i>	59	percent slopes, severely eroded	78
244A—Hartsburg silty clay loam, 0 to 2		801C—Orthents, silty, rolling	79
percent slopes	60	<i>Oscos Series</i>	79
<i>Herrick Series</i>	60	86B—Oscos silt loam, 2 to 5 percent slopes	80
46A—Herrick silt loam, 0 to 2 percent slopes	62	86C2—Oscos silt loam, 5 to 10 percent	
<i>Hickory Series</i>	62	slopes, eroded	81
8D—Hickory silt loam, 10 to 18 percent		864—Pits, quarries	81
slopes	63	862—Pits, sand	82
8D2—Hickory loam, 10 to 18 percent slopes,		<i>Plano Series</i>	82
eroded	64	199A—Plano silt loam, 0 to 2 percent slopes	83
8D3—Hickory clay loam, 10 to 18 percent		199B—Plano silt loam, 2 to 5 percent slopes	83
slopes, severely eroded	64	<i>Proctor Series</i>	84
8F—Hickory silt loam, 18 to 35 percent		7148A—Proctor silt loam, 0 to 2 percent	
slopes	65	slopes, rarely flooded	85
<i>Huntsville Series</i>	65	<i>Radford Series</i>	85
3077A—Huntsville silt loam, 0 to 2 percent		3074A—Radford silt loam, 0 to 2 percent	
slopes, frequently flooded	66	slopes, frequently flooded	86
<i>Ipava Series</i>	66	<i>Ross Series</i>	87
43A—Ipava silt loam, 0 to 2 percent slopes	67	3073A—Ross silt loam, 0 to 2 percent	
<i>Kendall Series</i>	68	slopes, frequently flooded	87
242A—Kendall silt loam, 0 to 2 percent		<i>Rozetta Series</i>	88
slopes	69	279B—Rozetta silt loam, 2 to 5 percent	
7242A—Kendall silt loam, 0 to 2 percent		slopes	89
slopes, rarely flooded	69	<i>Sable Series</i>	89
<i>Keomah Series</i>	70	68A—Sable silty clay loam, 0 to 2 percent	
17A—Keomah silt loam, 0 to 2 percent		slopes	90
slopes	71	<i>Sawmill Series</i>	91
830—Landfills	71	3107A—Sawmill silty clay loam, 0 to 2	
<i>Lawson Series</i>	72	percent slopes, frequently flooded	92
3451A—Lawson silt loam, 0 to 2 percent		<i>Shiloh Series</i>	92
slopes, frequently flooded	72	138A—Shiloh silty clay loam, 0 to 2 percent	
<i>Marseilles Series</i>	73	slopes	93
549G—Marseilles silt loam, 35 to 60 percent		<i>Spaulding Series</i>	94
slopes	74	712A—Spaulding silty clay loam, 0 to 2	
<i>Middletown Series</i>	75	percent slopes	95
685B—Middletown silt loam, 2 to 5 percent		<i>Tama Series</i>	95
slopes	76	36B—Tama silt loam, 2 to 5 percent slopes	96
685C2—Middletown silt loam, 5 to 10 percent		36C2—Tama silt loam, 5 to 10 percent	
slopes, eroded	76	slopes, eroded	97
<i>Navlys Series</i>	77	<i>Thebes Series</i>	97

212C2—Thebes silt loam, 5 to 10 percent slopes, eroded	98	Soil Properties	127
<i>Tice Series</i>	98	Engineering Index Properties	127
3284A—Tice silty clay loam, 0 to 2 percent slopes, frequently flooded	99	Physical Properties	128
533—Urban land	100	Chemical Properties	129
<i>Vesser Series</i>	100	Water Features	130
8396A—Vesser silt loam, 0 to 2 percent slopes, occasionally flooded	101	Soil Features	131
<i>Virden Series</i>	101	References	133
50A—Virden silty clay loam, 0 to 2 percent slopes	102	Glossary	135
<i>Worthen Series</i>	103	Tables	147
7037A—Worthen silt loam, 0 to 2 percent slopes, rarely flooded	103	Table 1.—Temperature and Precipitation	148
<i>Zook Series</i>	104	Table 2.—Freeze Dates in Spring and Fall	149
3405A—Zook silty clay loam, 0 to 2 percent slopes, frequently flooded	104	Table 3.—Growing Season	149
Use and Management of the Soils	107	Table 4.—Classification of the Soils	150
Interpretive Ratings	107	Table 5.—Acreage and Proportionate Extent of the Soils	151
Rating Class Terms	107	Table 6.—Limitations and Hazards Affecting Cropland and Pasture	153
Numerical Ratings	107	Table 7.—Land Capability and Yields per Acre of Crops and Pasture	157
Crops and Pasture	107	Table 8.—Prime Farmland	161
Management Considerations on Cropland ...	108	Table 9.—Hydric Soils	162
Explanation of Criteria	109	Table 10.—Windbreaks and Environmental Plantings	167
Management Considerations on Pasture	109	Table 11.—Forestland Productivity	182
Explanation of Criteria	110	Table 12.—Forestland Management (Part 1) ...	190
Yields per Acre	110	Table 13.—Forestland Management (Part 2) ...	197
Pasture and Hay Yields	110	Table 14.—Forestland Management (Part 3) ...	204
Land Capability Classification	111	Table 15.—Forestland Management (Part 4) ...	210
Prime Farmland	111	Table 16.—Forestland Management (Part 5) ...	214
Hydric Soils	112	Table 17.—Recreation (Part 1)	220
Windbreaks and Environmental Plantings	113	Table 18.—Recreation (Part 2)	226
Forestland	113	Table 19.—Wildlife Habitat	231
Forest Productivity	114	Table 20.—Building Site Development (Part 1)	236
Forest Management	115	Table 21.—Building Site Development (Part 2)	243
Recreation	116	Table 22.—Sanitary Facilities (Part 1)	252
Wildlife Habitat	118	Table 23.—Sanitary Facilities (Part 2)	261
Engineering	120	Table 24.—Construction Materials (Part 1)	268
Building Site Development	120	Table 25.—Construction Materials (Part 2)	274
Sanitary Facilities	121	Table 26.—Water Management (Part 1)	281
Construction Materials	123	Table 27.—Water Management (Part 2)	288
Water Management	124	Table 28.—Water Management (Part 3)	297
Waste Management	125	Table 29.—Engineering Index Properties	303

Table 30.—Physical Properties of the Soils	316	Table 32.—Water Features	332
Table 31.—Chemical Properties of the Soils	325	Table 33.—Soil Features	338

Issued 2004

Numerical Index to Map Units

8D—Hickory silt loam, 10 to 18 percent slopes	63	212C2—Thebes silt loam, 5 to 10 percent slopes, eroded	98
8D2—Hickory loam, 10 to 18 percent slopes, eroded	64	242A—Kendall silt loam, 0 to 2 percent slopes	69
8D3—Hickory clay loam, 10 to 18 percent slopes, severely eroded	64	244A—Hartsburg silty clay loam, 0 to 2 percent slopes	60
8F—Hickory silt loam, 18 to 35 percent slopes	65	249A—Edinburg silty clay loam, 0 to 2 percent slopes	47
17A—Keomah silt loam, 0 to 2 percent slopes	71	257A—Clarksdale silt loam, 0 to 2 percent slopes	39
36B—Tama silt loam, 2 to 5 percent slopes	96	259C2—Assumption silt loam, 5 to 10 percent slopes, eroded	28
36C2—Tama silt loam, 5 to 10 percent slopes, eroded	97	259D2—Assumption silt loam, 10 to 18 percent slopes, eroded	29
43A—Ipava silt loam, 0 to 2 percent slopes	67	279B—Rozetta silt loam, 2 to 5 percent slopes	89
45A—Denny silt loam, 0 to 2 percent slopes	43	280C2—Fayette silt loam, 5 to 10 percent slopes, eroded	55
46A—Herrick silt loam, 0 to 2 percent slopes	62	280D2—Fayette silt loam, 10 to 18 percent slopes, eroded	56
50A—Virden silty clay loam, 0 to 2 percent slopes	102	280D3—Fayette silty clay loam, 10 to 18 percent slopes, severely eroded	56
68A—Sable silty clay loam, 0 to 2 percent slopes	90	533—Urban land	100
86B—Osco silt loam, 2 to 5 percent slopes	80	536—Dumps, mine	46
86C2—Osco silt loam, 5 to 10 percent slopes, eroded	81	549G—Marseilles silt loam, 35 to 60 percent slopes	74
112A—Cowden silt loam, 0 to 2 percent slopes	42	567C2—Elkhart silt loam, 5 to 10 percent slopes, eroded	53
119C2—Elco silt loam, 5 to 10 percent slopes, eroded	50	567D2—Elkhart silt loam, 10 to 18 percent slopes, eroded	54
119D—Elco silt loam, 10 to 18 percent slopes	51	630C2—Navlys silt loam, 5 to 10 percent slopes, eroded	77
119D2—Elco silt loam, 10 to 18 percent slopes, eroded	51	630D2—Navlys silt loam, 10 to 18 percent slopes, eroded	78
119D3—Elco silty clay loam, 10 to 18 percent slopes, severely eroded	52	630D3—Navlys silty clay loam, 10 to 18 percent slopes, severely eroded	78
127B—Harrison silt loam, 2 to 5 percent slopes	58	679B—Blackberry silt loam, 2 to 5 percent slopes	30
127C2—Harrison silt loam, 5 to 10 percent slopes, eroded	58	684B—Broadwell silt loam, 2 to 5 percent slopes	32
131C2—Alvin fine sandy loam, 5 to 10 percent slopes, eroded	27	684C2—Broadwell silt loam, 5 to 10 percent slopes, eroded	32
131D2—Alvin fine sandy loam, 10 to 18 percent slopes, eroded	27	685B—Middletown silt loam, 2 to 5 percent slopes	76
134A—Camden silt loam, 0 to 2 percent slopes	36	685C2—Middletown silt loam, 5 to 10 percent slopes, eroded	76
134B—Camden silt loam, 2 to 5 percent slopes	36		
134C2—Camden silt loam, 5 to 10 percent slopes, eroded	37		
138A—Shiloh silty clay loam, 0 to 2 percent slopes	93		
198A—Elburn silt loam, 0 to 2 percent slopes	49		
199A—Plano silt loam, 0 to 2 percent slopes	83		
199B—Plano silt loam, 2 to 5 percent slopes	83		

705A—Buckhart silt loam, 0 to 2 percent slopes	34	3284A—Tice silty clay loam, 0 to 2 percent slopes, frequently flooded	99
705B—Buckhart silt loam, 2 to 5 percent slopes	34	3405A—Zook silty clay loam, 0 to 2 percent slopes, frequently flooded	104
712A—Spaulding silty clay loam, 0 to 2 percent slopes	95	3451A—Lawson silt loam, 0 to 2 percent slopes, frequently flooded	72
801C—Orthents, silty, rolling	79	7037A—Worthen silt loam, 0 to 2 percent slopes, rarely flooded	103
830—Landfills	71	7075B—Drury silt loam, 2 to 5 percent slopes, rarely flooded	45
862—Pits, sand	82	7148A—Proctor silt loam, 0 to 2 percent slopes, rarely flooded	85
864—Pits, quarries	81	7242A—Kendall silt loam, 0 to 2 percent slopes, rarely flooded	69
3073A—Ross silt loam, 0 to 2 percent slopes, frequently flooded	87	8396A—Vesser silt loam, 0 to 2 percent slopes, occasionally flooded	101
3074A—Radford silt loam, 0 to 2 percent slopes, frequently flooded	86		
3077A—Huntsville silt loam, 0 to 2 percent slopes, frequently flooded	66		
3107A—Sawmill silty clay loam, 0 to 2 percent slopes, frequently flooded	92		

Foreword

This soil survey contains information that affects land use planning in this survey area. It contains predictions of soil behavior for selected land uses. The survey also highlights soil limitations, improvements needed to overcome the limitations, and the impact of selected land uses on the environment.

This soil survey is designed for many different users. Farmers, foresters, and agronomists can use it to evaluate the potential of the soil and the management needed for maximum food and fiber production. Planners, community officials, engineers, developers, builders, and home buyers can use the survey to plan land use, select sites for construction, and identify special practices needed to ensure proper performance. Conservationists, teachers, students, and specialists in recreation, wildlife management, waste disposal, and pollution control can use the survey to help them understand, protect, and enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. The information in this report is intended to identify soil properties that are used in making various land use or land treatment decisions. Statements made in this report are intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are shallow to bedrock. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

These and many other soil properties that affect land use are described in this soil survey. The location of each soil is shown on the detailed soil maps. Each soil in the survey area is described. Information on specific uses is given for each soil. Help in using this publication and additional information are available at the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

William J. Gradle
State Conservationist
Natural Resources Conservation Service

Soil Survey of Sangamon County, Illinois

By James K. Hornickel, Natural Resources Conservation Service

Original fieldwork by J.F. Steinkamp (survey leader), G.V. Berning, R.E. Bourland, T.M. Goddard, K.A. Gotsch, D.C. Hallbick, G.W. Hudelson, L.M. Kiefer, D.R. Mapes, J.W. Scott, and J.A. Thompson, Soil Conservation Service

Updated fieldwork by John W. Ford, Paula K. French, James K. Hornickel, William M. Teater, and Robert A. Tegeler, Natural Resources Conservation Service

Map compilation by Dale A. Baumgartner, Gerald V. Berning, John W. Ford, Kenneth A. Gotsch, James K. Hornickel, William M. Teater, and Robert A. Tegeler, Natural Resources Conservation Service

United States Department of Agriculture, Natural Resources Conservation Service, in cooperation with the Illinois Agricultural Experiment Station

General Nature of the County

SANGAMON COUNTY is in the central part of Illinois (fig. 1). It has a total area of 560,755 acres, or about 876 square miles. According to the 2000 census, the population of the county is 188,951 and the population of Springfield, the largest city in the county, is 111,545 (United States Census Bureau, 2002). Springfield is the county seat and the State capital.

This soil survey updates the survey of Sangamon County published in 1980 (Steinkamp, 1980). It provides additional information and has larger maps, which show the soils in greater detail.

History

Settlement of the survey area began more than 175 years ago. In 1818, Springfield was settled by people from South Carolina, Kentucky, and Virginia. It became the county seat in 1821, when Sangamon County was established. In 1831, Springfield was incorporated as a town. Partly through the efforts of Abraham Lincoln, it was made the State capital in 1837.

The area now known as Sangamon County at one time was part of the territory claimed by the Kickapoo

and Pottawatomie Indians. At that time, prairie grasses covered about 70 percent of the survey area, mainly the nearly level to strongly sloping uplands. The rest of the survey area, mostly the more sloping land near streams, was timbered. The early settlers cleared and farmed the timbered areas. By 1830, metal plows were available, and the settlers started plowing the prairie. About 1875, underground tile drains became available. They provided the artificial drainage needed for cultivation of the wet, nearly level soils.

Farming and Industry

Hal Pyle, District Conservationist, Natural Resources Conservation Service, helped prepare this section

Soil is the most important natural resource in the county. Crops produced on the soil and livestock that feed on some of these crops contribute to the wealth of the county. According to the Census of Agriculture, 466,956 acres, or 82 percent of the county, was farmland in 1997 (USDA, 1997). The county had 993 farms, which averaged 470 acres in size. Since 1980, the average farm size has increased 61 percent and the number of farms has decreased. In 1999,




Figure 1.—Location of Sangamon County and the major land resource areas (MLRAs) in Illinois (USDA, 1981).

Sangamon County ranked 7th in the State in soybean production with 42 bushels per acre. It ranked 12th in the State in corn production with 145 bushels per acre (Illinois Agricultural Statistics Service, 2002; <http://www.agstats.state.il.us/website/reports.htm>).

The water supply in the county is adequate for domestic use and for watering livestock (Illinois Department of Natural Resources, 2000b). The county is well supplied with surface water, but the upper layers of Pennsylvanian bedrock are unreliable sources of good-quality ground water. Lake Springfield supplies water for most of the county. Good sites for reservoirs are available along the Sangamon River and along many small streams.

Small oilfields are in scattered areas throughout the east-central part of the county. Oil-producing strata consist of Devonian and Silurian rocks, mainly

limestone or dolomite limestone. In most places they are within a depth of 2,000 feet. The Carbonale geologic formation is a limited source of coal. Most of the coal has been mined out by underground operations.

Sand and gravel are extracted mainly from the low terraces and flood plain along the Sangamon River east of Springfield. Deposits of resource quality are on hills and ridges and in other areas in the valley of the Sangamon River west of Springfield. In the past, limestone, clay, and shale were mined from the Pennsylvanian rocks. The limestone was used in roadbeds and in the construction of buildings and the production of agricultural lime. The shale was used in the manufacture of tile drains, sewer pipes, pottery, stoneware, and other clay products.

Transportation Facilities

A system of railroads and highways facilitated the rapid settlement and development of the county. The main highways crossing the county are Interstate 55, from north to south, and Interstate 72, from east to west. Other highways are U.S. Highway 36 and State Routes 4, 29, 54, 97, 104, and 125. All farms have access to surfaced roads. Capitol Airport, northwest of Springfield, provides freight and passenger service. Several railroads also provide freight and passenger service.

Physiography and Drainage

Sangamon County is in the Springfield Plain region (Bergstrom et al., 1976). Shallow river valleys dissect the Springfield Plain. Moderately thick glacial till was deposited during Illinoian and earlier stages of glaciation. The Buffalo Hart moraine, a broad, discontinuous ridge, rises above the general level of the plain in the vicinity of Buffalo Hart.

The natural drainage is westward. The Sangamon River and its tributaries—the South Fork of the Sangamon River and numerous smaller streams—drain all of the survey area, except for the northeast corner.

Climate

Prepared by the National Water and Climate Center, Natural Resources Conservation Service, and the Illinois State Water Survey.

Table 1 gives data on temperature and precipitation for the survey area as recorded at Springfield in the period 1970 to 1999. Table 2 shows probable dates of the first freeze in fall and the last freeze in spring. Table 3 provides data on length of the growing season.

In winter, the average temperature is 31 degrees F and the average daily minimum temperature is 22 degrees. The lowest temperature on record, which occurred at Springfield on January 20, 1979, is -21 degrees. In summer, the average temperature is 73 degrees and the average daily maximum temperature is 83 degrees. The highest recorded temperature, which occurred on July 14, 1954, is 112 degrees.

Growing degree days are shown in table 1. They are equivalent to “heat units.” During the month, growing degree days accumulate by the amount that the average temperature each day exceeds a base temperature (50 degrees F). The normal monthly accumulation is used to schedule single or successive

plantings of a crop between the last freeze in spring and the first freeze in fall.

The total average annual precipitation is nearly 36 inches. Of this, more than 21 inches, or about 60 percent, usually falls in April through September. The growing season for most crops falls within this period. In 2 years out of 10, the rainfall in April through September is less than 18 inches. The heaviest 1-day rainfall during the period of record was 6.12 inches at Springfield on December 3, 1982. Thunderstorms occur on about 50 days each year. Tornadoes and severe thunderstorms strike occasionally. These storms are usually local in extent and of short duration and cause damage in scattered areas.

The average seasonal snowfall is about 24 inches. The greatest snow depth at any one time during the period of record was 22.7 inches. On the average, 15 days of the year have at least 1 inch of snow on the ground. The number of such days varies greatly from year to year.

The average relative humidity in midafternoon is about 60 percent. Humidity is higher at night, and the average at dawn is about 80 percent. The sun shines 70 percent of the time possible in summer and 50 percent in winter. The prevailing wind is from the south. Average windspeed is highest, 14 miles per hour, in March.

How This Survey Was Made

This survey was made to provide updated information about the soils and miscellaneous areas in Sangamon County, which is a subset of MLRAs 108B and 115C. MLRAs are geographically associated land resource units that share a common land use, elevation, topography, climate, pattern of water and soils, and vegetation (USDA, 1981). Map unit design is based on the occurrence of each soil throughout the MLRAs. In some cases a soil component may be referred to that does not occur in the Sangamon County subset but that has been mapped within the MLRAs.

The information in this survey includes a description of the soils and miscellaneous areas and their location and a discussion of their suitability, limitations, and management for specified uses.

Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They prepared new soil profile descriptions and studied many existing profile descriptions. These descriptions show the sequence of natural layers, or

horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

The soils and miscellaneous areas in the survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind or segment of the landscape. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landscape, soil scientists develop a concept, or model, of how the soils were formed. Thus, during the update, this model enables the soil scientists to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Individual soils on the landscape commonly merge into one another as their characteristics gradually change. To construct an accurate map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they observed. The maximum depth of observation was about 80 inches (6.7 feet). The scientists noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, soil reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

While a soil survey is in progress, samples of some of the soils in the survey area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Interpretations are modified as necessary to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a water table within certain depths in most years, but they cannot predict that the water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.

Aerial photographs used in this update survey were taken between 1993 and 1996. Soil scientists also studied U.S. Geological Survey topographic maps enlarged to a scale of 1:12,000 and orthophotographs to relate land and image features. Specific soil boundaries from the soil maps published in 1980 were drawn on the orthophotographs. Adjustments of soil boundary lines were made to coincide with the U.S. Geological Survey topographic map contour lines and tonal patterns on aerial photographs.

The descriptions, names, and delineations of the soils in this county may not fully agree with those of the soils in adjacent counties. Differences are the result of a better knowledge of soils, modifications in series concepts, or variations in the intensity of mapping or in the extent of the soils in the counties.

Formation and Classification of the Soils

This section relates the soils in the survey area to the major factors and processes of soil formation and describes the system of soil classification.

Formation of the Soils

Steve Suhl, Resource Soil Scientist, Natural Resources Conservation Service, prepared this section.

A soil is a three-dimensional natural body consisting of mineral and organic material that can support plants. The nature of any soil at a given site is the result of the interaction of the major factors of soil formation and their influence on the processes of soil formation.

Factors of Soil Formation

The major factors of soil formation are parent material, climate, plants and animals, topography, and time (Jenny, 1941). Climate and plants and animals act directly on the parent material, which is modified by topography over time. Theoretically, if all these factors were identical at different sites, the soils at these sites would be identical. The diversity among soils is the result of unique combinations of the soil-forming factors. Soils are continually evolving in response to these factors.

Parent Material

Parent material is the unconsolidated geologic material in which a soil forms. It determines the basis for the chemical and mineralogical composition of the soil. The properties of the parent material vary greatly, sometimes within small areas, depending on how the material was deposited. The soils in Sangamon County formed in a variety of parent materials. The majority of the soils formed in loess. Other soils formed in glacial drift, alluvium, sandy eolian deposits, bedrock residuum, or a combination of these.

Glacial drift is glacially deposited sediment. There are two main types of glacial drift—till and outwash. Till is material that was deposited directly by glacial ice with little or no water action. It typically has particles that vary in size, including sand, silt, clay, and some

pebbles, cobbles, and larger rock fragments. The small pebbles in till generally have distinct edges and corners, indicating that they have not been subject to intense washing by water. Till is well graded and unstratified. The till in Sangamon County was deposited during the Illinoian age. Soils that formed in till are of minor extent in the county. Hickory soils are examples.

During the Sangamon interglacial stage, which occurred between the Illinoian and Wisconsinan stages, the relatively flat, stable till surface was exposed to intense weathering. A soil formed on the till surface and was subsequently buried by depositions of loess. In Sangamon County, the loess deposits were thick enough to remove the soil from the influence of the active soil-forming processes. The soils that formed in the till are called paleosols, and they reflect the environmental conditions of their formation period. There are several types of paleosols. The type in Sangamon County is called a buried paleosol. A buried paleosol generally is no longer subject to the soil-forming processes that created it. In some areas where the loess deposits are thinner, however, the current processes of soil formation have extended through the loess and into the upper part of the paleosol. The result is a welded soil profile. Elco soils are examples of soils that formed in these areas.

Outwash includes all sediments deposited by running water from melting glaciers. The size of the particles that can be transported by water, either as bed load or suspended sediments, depends on the gradient, volume, and velocity of the moving water. Water velocity decreases when a stream loses grade or flows into a larger body of water. As the velocity decreases, suspended particles begin to settle out. The coarser materials, such as gravel and cobbles, are deposited nearer to the source; the finer materials, such as fine sands, silts, and clays, are carried farther downstream. The pebbles in outwash generally have rounded edges and corners, indicating that they have been subject to intense washing by water. Outwash is poorly graded, is stratified, and varies in composition because of variations in the flow of water. Outwash is generally permeable. The outwash in Sangamon County was deposited during the Wisconsinan age.

Soils that formed in outwash are of minor extent in the county. Camden soils are examples.

Alluvium is material deposited by running water. There are two major types—stream alluvium and valley-side alluvium.

Stream alluvium is soil material deposited by floodwater along streams. The source of the alluvium generally is material eroded from other parent materials farther upstream in the watershed. Stream alluvium is poorly graded, stratified, and well sorted. The texture of the soil material varies, depending on the speed of the floodwater, the duration of flooding, and the distance from the streambank. Once the faster moving water within the stream channel is outside the channel, it slows quickly as the concentrated channel flow changes to broad overland flow. The capacity of the stream to carry a sediment load decreases as the water velocity decreases. The coarser textured material is deposited first, near the channel. The fine textured material is carried a greater distance from the channel. Ross soils are examples of soils that formed in the coarser textured alluvium close to the stream channel. Tice soils formed in the finer textured alluvium farther from the stream channel.

Valley-side alluvium is poorly graded and stratified, but it generally is not well sorted. The source of the alluvium generally is material eroded from parent material directly upslope. The soils that formed in valley-side alluvium are similar in character to the upslope source material. Drury soils formed in valley-side alluvium.

Eolian sediments are materials transported and deposited by the wind. These sediments were derived from periglacial regions where sparse vegetation and low temperatures and precipitation rates left unconsolidated sediments exposed to wind action. The unconsolidated sediments, primarily outwash, were then stripped of their finer components by the strong wind. Eolian sediments were deposited during the Wisconsinan age and are either loess or windblown sand. Loess is the major parent material in Sangamon County. It is fine grained and poorly graded and averages less than 7 percent sand. The loess is about 15 feet thick in the northern part of the county and thins to less than 7 feet in the southern part. Fayette and Osco soils formed in loess.

Windblown sand is poorly graded and consists primarily of very fine sand and fine sand. It generally is in scattered areas on the south side of the valley of the Sangamon River. The soils that formed in windblown sand are of minor extent in Sangamon County. Alvin soils are examples.

The bedrock residuum in Sangamon County is material weathered from shale and sandstone. It is

generally grayish and unstratified. The bedrock is Pennsylvanian in age. The soils that formed in bedrock residuum are of minor extent in the county. Marseilles soils are examples.

Climate

The climate in Sangamon County has significantly affected the soil-forming processes. The county has a humid, temperate climate. In this climatic environment, physical and chemical weathering of the parent material can occur along with the accumulation of organic matter, the decomposition of minerals, the formation and translocation of clay, the leaching of soluble compounds, and alternating periods of freezing and thawing.

The two climatic factors that have the greatest influence on soil-forming processes are precipitation and temperature. Precipitation supplies the moisture needed for most physical and chemical processes and determines the depth to which these processes occur. The soil moisture regime, which is partially a function of precipitation, determines the processes that occur in the soil. The rate at which these physical and chemical processes proceed depends on the temperature, particularly the soil temperature regime.

Two soil moisture regimes occur in the county—aquic and udic. The aquic moisture regime is a reducing regime in a soil that is virtually free of dissolved oxygen because of saturation by water, including water of the capillary fringe. Biological activity is necessary for the removal of dissolved oxygen from ground water. Therefore, the soil temperature must be above biologic zero (5 degrees C) for some time while the soil is saturated. Sawmill soils have an aquic soil moisture regime. The udic moisture regime is one in which the soil moisture control section is not dry in any part for as long as 90 cumulative days per year. Also required, except for short periods, is a three-phase system, solid-liquid-gas, in part or all of the soil moisture control section when the soil temperature is above biologic zero. Osco soils have a udic soil moisture regime.

The mesic soil temperature regime is the only temperature regime recognized in the county. This regime is one in which the mean annual soil temperature is 8 degrees to less than 15 degrees C and the difference between mean summer and mean winter soil temperatures is more than 5 degrees C at a depth of 20 inches.

Plants and Animals

The vegetation under which a soil forms influences several important soil properties, such as color, structure, reaction, and content and distribution of

organic matter. Vegetation extracts water from the soil, recycles nutrients, and adds organic matter to the soil. Gases derived from root respiration combine with water to form acids that influence the weathering of minerals.

Several different types of vegetation have influenced soil formation in Sangamon County. These include prairie vegetation, upland hardwood forests, forest and prairie vegetation in transitional areas, and vegetation on flood plains.

Prairie vegetation.—The decomposition of the roots of annual prairie grasses provides well distributed subsurface accumulations of organic material, resulting in a thick, dark surface layer. Osco soils formed under prairie vegetation. The average content of organic matter in the surface layer of these soils is 3 to 4 percent.

Upland hardwood forests.—These forests contribute organic matter to the soil primarily when they add leaf litter to the surface each year, resulting in a thin, dark surface layer. Fayette soils formed under this type of vegetation. The average content of organic matter in the surface layer of these soils is 1 to 2 percent.

Forest and prairie vegetation in transitional areas.—Soils that formed in these areas exhibit modified characteristics of both forest and prairie vegetative regimes. An example is Clarksdale soils, which have a surface layer that is thinner than that of the soils that formed under prairie vegetation. The average content of organic matter in the surface layer of Clarksdale soils is 2 to 3 percent.

Vegetation on flood plains.—The soils on these plains formed under a combination of trees and grasses. They have colors that largely reflect those of the sediments in which they formed. Lawson soils are examples.

Bacteria, fungi, and many other micro-organisms decompose organic matter and release nutrients to growing plants. They influence the formation of peds. Soil properties, such as drainage, temperature, and reaction, influence the type of micro-organisms that live in the soil. Fungi are generally more active in the more acid soils, while bacteria are more active in the less acid soils.

Earthworms, crayfish, insects, and small burrowing animals mix the soil and create small channels that influence soil aeration and the percolation of water. Earthworms help to incorporate crop residue or other organic matter into the soil. The organic material improves tilth. In areas that are well populated with earthworms, the leaf litter that accumulates on the soil in the fall is generally incorporated into the soil by the following spring. If the earthworm population is low,

part of the leaf fall can remain on the surface of the soil for several years.

Human activities have significantly influenced soil formation through their effect on soil health. Soil health has been damaged by degradation processes, such as erosion, compaction, contamination, disaggregation, loss of biological activity, and nutrient depletion. Native forests have been cleared and wet soils drained for farming and other uses. The development of land for urban uses has significantly influenced the soils in some areas.

Topography

Topography, or the configuration of the land surface in terms of relief and contour, influences soil formation mainly through its effect on surface water runoff or accumulation and on erosion and deposition. The degree of the effect of topography depends on the type and stability of the land surface.

There are two types of land surfaces—aggrading and degrading—and three levels of stability—stable, metastable, and active. The aggrading surfaces in Sangamon County receive material either through the deposition associated with flooding or through the accumulation of erosional sediments. Ross soils formed on natural levees on flood plains. The natural levees are an example of active-aggrading land surfaces. They receive depositions of sediment during frequent episodes of flooding. Drury soils formed on footslopes that receive runoff with some accumulation of hillslope sediments. Footslopes are an example of metastable-aggrading land surfaces. Sable soils formed in broad, low areas on drainage divides that receive runoff from upslope areas but accumulate little sediment from hillslope erosion. These broad, low areas are an example of stable-aggrading land surfaces.

Degrading surfaces lose material primarily through erosion. Keomah soils formed on the broad summits of interfluves. These broad summits are an example of stable-degrading surfaces, where runoff is limited. Fayette soils are on the shoulders of hillslopes and thus are more susceptible to runoff and erosion than the Keomah soils. Shoulders are metastable-degrading surfaces, where increased runoff leads to higher rates of erosion. Backslopes are an example of active-degrading surfaces. Hickory soils are on backslopes, where runoff and erosion rates are highest.

Time

The length of time that the parent material has been exposed to soil-forming processes influences the degree of genetic horizon development that occurs in

the soil. Evaluation of time as a factor of soil formation is difficult because of the effects of the other soil-forming factors. The influence of time can be modified by erosion, deposition of material, topography, and kind of parent material.

Soils on flood plains receive alluvial material during each flood. This repeated deposition interrupts soil formation. Lawson soils are examples of immature soils that formed in stream alluvium.

Processes of Soil Formation

Soil forms through the complex interaction of four general processes. These processes are *additions*, *transformations*, *removals*, and *translocations*. The degree of interaction of each these processes varies, resulting in the variety of soils on the landscape.

Additions to the soil can occur directly through the deposition of sediment by floodwater or through the accumulation of windblown sediment. They also occur through the accumulation and incorporation of organic matter in the A horizon of mineral soils. The most striking example of this addition is the formation of a mollic epipedon. The mollic epipedon forms in an environment that features optimum amounts of moisture, temperature, and bivalent cations. Such an environment allows grasses to thrive. The grassland vegetation produces large amounts of organic matter. Microbial decomposition of subsurface organic residues and of surface organic residues taken underground by soil fauna results in the most recognizable property of the mollic epipedon, a dark color. Ipava soils are examples of soils that have a mollic epipedon.

Transformations are changes that take place in the soil through the interaction of biological, chemical, and physical processes. An example is the reduction of iron and manganese oxides, which occurs in soils that are saturated with water. Typically, iron oxides coat soil particles and produce brownish, yellowish, or reddish colors, and manganese oxides produce black colors. When a soil becomes saturated with water and the dissolved oxygen is removed, anaerobic conditions develop. These conditions result in changes in the biochemical processes occurring in the soils and in the development of distinctive soil morphological characteristics (redoximorphic features). Reduced iron and manganese can move with the soil water to other parts of the soil or can be removed entirely from the soil by leaching. After the iron and manganese are removed, the leached area, or depletion zone, generally is grayish or whitish. If the reduced iron comes in contact with oxygen, it can reoxidize. The result is the formation of bright colored concentrations

or accumulations. Repeated cycles of saturation and drying result in a mottled soil. Part of the soil is gray because of the loss of iron, and other parts are brown because the iron oxide has accumulated or has not been removed. The somewhat poorly drained Ipava soils are examples of soils in which this process has occurred. Iron may be leached from soils that remain saturated for long periods. Such soils are generally grayish or gleyed. The poorly drained Titus soils are examples.

Removals from the soil can occur as solid mineral and organic particles are lost through erosion of the soil surface. Such losses can be serious because the material lost generally is from the most productive part of the soil profile.

Removals can also occur as a result of leaching. The leaching of calcium carbonate from calcareous loess is an example. The loess was initially high in content of calcium carbonate. Water percolating through the loess dissolved the calcium carbonate and transported it deeper into the solum. Calcium carbonate is relatively soluble and is removed early in the formation of the soil. It is also a powerful flocculent, creating soil particles too large to be transported in suspension in the soil water. Removal of calcium carbonate facilitates the dispersion of clay particles. Translocation of the dispersed clay particles can then occur in percolating soil water.

Translocations are movements of material from one part of the soil to another. An example is the translocation of clay from the A or E horizon, the zone of eluviation or loss, to the B horizon, the zone of illuviation or gain. In Fayette soils, for example, a significant amount of clay has accumulated, forming an illuvial horizon called an argillic horizon. The argillic horizon developed on a relatively old, stable landscape. Water from rain and melting snow transferred fine clay from the A or E horizon downward to the B horizon, where the clay particles were deposited on the faces of peds and along pores.

Soils and Soil-Landscape Units

Soils are natural bodies that are distributed on the landscape in a predictable way in response to a systematic interaction of the five factors of soil formation. The relationship of the landscape to the five factors results in a soil-landscape unit (Hudson, 1992). A soil-landscape unit is similar to a landform that has been modified by one or more of the soil-forming factors. Within a particular soil-landscape unit, the same kind of soil should form. Changes in the interaction of one or more of the five factors leads to a change in the soil-landscape unit, influencing the soil-

forming processes and the soil that forms within this unit.

The following paragraphs describe the relationships and interactions that occur in some of the more common soil-landscape units in Sangamon County and the soils that have formed in these units.

Upland landscapes predominate in Sangamon County. These landscapes range from broad, relatively undissected drainage divides to dissected areas adjacent to river bluffs. The predominant parent material in the more stable landscape positions is loess. Much of the calcium carbonate that was present when the loess was deposited has been leached to a sufficient depth to facilitate soil formation.

Low areas on the broad drainage divides are on stable-aggrading land surfaces that receive water as direct precipitation and as runoff from upslope areas. These conditions result in a wet soil microclimate. A seasonal high water table is at or near the surface much of the year, and at times the areas are ponded. Redoximorphic features associated with prolonged saturated conditions, such as a depleted soil matrix and iron and manganese accumulations along root channels and pores, occur throughout the soil profile.

The native vegetation in this landscape unit consisted of prairie grasses. Additions of large amounts of organic matter from the incorporation of plant residues by soil fauna and from decomposition of the extensive and deep root systems of the grasses resulted in a thick, dark surface horizon, called a mollic epipedon.

The saturated conditions and poor aeration influenced the rate at which organic matter decomposed. This rate is slower in soils that are saturated for prolonged periods, resulting in a thicker mollic epipedon and a higher content of organic matter than is characteristic of the better aerated positions upslope.

The extended periods of saturation impeded the illuviation of clay and the formation of an argillic horizon. A cambic horizon has developed through the aggregation of soil particles into structural units, or peds, and the development of redoximorphic features. Sable soils formed in these low areas.

Upslope from the low areas is a soil-landscape unit that consists of the summits of broad rises on drainage divides. These summits are stable-degrading land surfaces that receive water primarily through direct precipitation. The seasonal high water table and associated redoximorphic features occur at a lower depth in the soils of this unit than in the soils in the adjacent low areas. The redoximorphic features also

indicate a fluctuating water table. The soil microclimate in the upper part of the profile alternates between periods when the soils are unsaturated and periods when they are saturated. A yellowish brown soil matrix indicates an oxidizing environment. Redoximorphic depletions along root channels and pores and iron and manganese accumulations within the matrix, however, indicate short periods of saturation. The lower part of the profile is saturated for extended periods and has a depleted soil matrix.

The native vegetation in this landscape unit consisted of prairie grasses. The soils of this unit are better aerated than the soils in low areas and tend to have a higher rate of organic-matter decomposition. Thus, they generally have a slightly thinner mollic epipedon and a lower content of organic matter than the soils in the low areas.

The fluctuating water table disrupts the soil fabric through wetting and drying cycles, which aid in the dispersal of clay, the movement of clay with percolating water, and the precipitation of clay as films on the faces of peds and as linings of pores. The result is the formation of an illuvial horizon called an argillic horizon. Ipava soils formed in areas of this soil-landscape unit.

The soil-landscape unit in the more dissected areas consists of the broad summits of interfluves. It has characteristics similar to those of the unit on the summits of broad rises on drainage divides. These dissected areas are stable-degrading land surfaces that receive water primarily through direct precipitation. Depth to the seasonal high water table and the associated redoximorphic features are nearly identical to those of the soil-landscape unit on the summits of broad rises.

The native vegetation in this soil-landscape unit is transitional between forest and prairie vegetation. In areas of forest vegetation, most of the organic matter is deposited on the surface. In areas of prairie grasses, extensive root systems provide most of the organic material. The soils in this unit have a dark surface layer, but they do not have a mollic epipedon because the dark surface layer is not thick enough and does not have a sufficient accumulation of organic matter. This type of surface horizon is called an ochric epipedon.

A light colored, eluvial subsurface horizon (called an albic horizon) also has developed in the soils of this unit. This horizon is typical of soils that formed under forest vegetation. In this horizon, much of the clay and free iron oxides have been removed and the color is determined primarily by the uncoated silt particles and sand grains. The translocation of clay from the eluvial

horizon to the illuvial horizon results in the formation of an argillic horizon. Clarksdale soils are in areas of this soil-landscape unit.

Adjacent to this soil-landscape unit is a unit that also consists of the summits of interfluves but that is generally closer to the opposing interfluve drainageways and is on narrower summits. These summits are stable-degrading land surfaces that receive water through direct precipitation. Water that does not infiltrate the soil is lost through surface flow or runoff. Runoff increases the susceptibility to erosion.

The seasonal high water table and associated redoximorphic features occur at a much lower depth than those in the soils on the broad summits. The upper part of the soils formed in an oxidizing environment. The soil matrix is generally yellowish brown or brown and is free of depletions. The matrix in the lower part of the profile is yellowish brown and generally has depletions along pores and root channels, indicating short periods of saturation.

The native vegetation in areas of this soil-landscape unit is broadleaf deciduous forest. Under forest vegetation, most of the addition of organic matter occurs above ground. Organic matter is not incorporated as deep in the soil profile as it is in soils that formed under prairie vegetation, and the content decreases rapidly with increasing depth. Therefore, the dark surface layer, which is generally evident prior to tillage, is thinner than that in the Clarksdale soils. An ochric epipedon and an albic horizon have developed.

The more acid leaching environment that occurs under forest vegetation allows dispersed clay particles to be translocated to a greater depth than is characteristic in similar positions under prairie vegetation. The result is a well developed argillic horizon. Rozetta soils formed in areas of this soil-landscape unit.

Downslope from this soil-landscape unit is a unit that consists of the shoulders of hillslopes. These are metastable-degrading land surfaces that receive water through direct precipitation but also lose some of this water through runoff. Runoff increases the susceptibility to erosion.

The seasonal high water table is at a depth similar to that of the Rozetta soils. The soil matrix is yellowish brown or brown. Depletions in the lower part of the profile are generally along pores and root channels.

The native vegetation in this soil-landscape unit is forest. The soils have an ochric epipedon and albic and argillic horizons. Navlys soils formed in areas of this soil-landscape unit.

Farther downslope is a unit that consists of the backslopes of side slopes. These backslopes are active-degrading land surfaces that receive water through direct precipitation but also lose much of this water through runoff. The loss of water results in a drier soil microclimate. The seasonal high water table is below the depth of the developing soil profile. The entire profile formed in an oxidizing environment. The soil matrix is yellowish brown or brown and is free of depletions.

The native vegetation is forest. Like the Navlys soils, the soils in these areas have an ochric epipedon and albic and argillic horizons. Fayette soils formed in areas of this soil-landscape unit.

On the narrow flood plains between opposing side slopes is an active-aggrading landscape position that receives sediment during frequent episodes of flooding. The nearly continual deposition of sediment interrupts the soil-forming processes. The result is less time for soil formation to proceed and a less developed soil profile. A mollic epipedon is evident in the developing soil profile, but the fine stratification common to recent alluvial deposits remains and no diagnostic subsurface horizons occur. Lawson soils formed in areas of this soil-landscape unit.

Classification of the Soils

The system of soil classification used by the National Cooperative Soil Survey has six categories (Soil Survey Staff, 1999). Beginning with the broadest, these categories are the order, suborder, great group, subgroup, family, and series. Classification is based on soil properties observed in the field or inferred from those observations or from laboratory measurements. Table 4 shows the classification of the soils in the county. The categories are defined in the following paragraphs.

ORDER. Twelve soil orders are recognized. The differences among orders reflect the dominant soil-forming processes and the degree of soil formation. Each order is identified by a word ending in *sol*. An example is Mollisol.

SUBORDER. Each order is divided into suborders primarily on the basis of properties that influence soil genesis and are important to plant growth or properties that reflect the most important variables within the orders. The last syllable in the name of a suborder indicates the order. An example is Aquoll (*Aqu*, meaning water, plus *oll*, from Mollisol).

GREAT GROUP. Each suborder is divided into great groups on the basis of close similarities in kind, arrangement, and degree of development of

pedogenic horizons; soil moisture and temperature regimes; type of saturation; and base status. Each great group is identified by the name of a suborder and by a prefix that indicates a property of the soil. An example is Endoaquolls (*Endo*, meaning within, plus *aquoll*, the suborder of the Mollisols that has an aquatic moisture regime).

SUBGROUP. Each great group has a typic subgroup. Other subgroups are intergrades or extragrades. The typic is the central concept of the great group; it is not necessarily the most extensive. Intergrades are transitions to other orders, suborders, or great groups. Extragrades have some properties that are not representative of the great group but do not indicate transitions to any other known kind of soil. Each subgroup is identified by one or more adjectives preceding the name of the great group. The adjective *Typic* identifies the subgroup that typifies the great group. An example is Typic Endoaquolls.

FAMILY. Families are established within a subgroup on the basis of physical and chemical properties and other characteristics that affect management. Generally, the properties are those of horizons below plow depth where there is much biological activity. Among the properties and characteristics considered are particle-size class, mineralogy class, cation-exchange activity class, soil temperature regime, soil depth, and reaction class. A family name consists of the name of a subgroup preceded by terms that indicate soil properties. An example is fine-silty, mixed, superactive, mesic Typic Endoaquolls.

SERIES. The series consists of soils that have similar horizons in their profile. The horizons are similar in color, texture, structure, reaction, consistence, mineral and chemical composition, and arrangement in the profile. The texture of the surface layer or of the substratum can differ within a series.

Soil Series and Detailed Soil Map Units

In this section each soil series recognized in the survey area is described. Each series description is followed by detailed descriptions of the associated detailed soil map units.

Characteristics of the soil and the material in which it formed are identified for each soil series. A pedon, a small three-dimensional area of soil, that is typical of the series in the survey area is described. The detailed description of each soil horizon follows standards in the "Soil Survey Manual" (Soil Survey Division Staff, 1993). Many of the technical terms used in the descriptions are defined in "Soil Taxonomy" (Soil Survey Staff, 1999). Unless otherwise stated, colors in the descriptions are for moist soil. Following the pedon description is the range of important characteristics of the soils in the series.

The map units on the detailed soil maps in this survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions in this section, along with the maps, can be used to determine the suitability and potential of a unit for specific uses. They also can be used to plan the management needed for those uses. More information about each map unit is given under the headings "Use and Management of the Soils" and "Soil Properties."

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus

they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. The contrasting components are mentioned in the map unit descriptions. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives the principal hazards and limitations to be considered in planning for specific uses.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer or of the underlying layers, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer or of the underlying layers. They also can differ in slope, stoniness, salinity, wetness, degree of erosion, and other characteristics that affect their use.

On the basis of such differences, a soil series is divided into *soil phases*. The name of a soil phase commonly indicates a feature that affects use or management. For example, Fayette silt loam, 5 to 10 percent slopes, eroded, is a phase of the Fayette series.

This survey includes *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Pits, sand, is an example.

Table 5 gives the acreage and proportionate extent of each map unit. Other tables give properties of the soils and the limitations, capabilities, and potentials for many uses. The Glossary defines many of the terms used in describing the soils or miscellaneous areas. The letters "OSD" following the heading "Typical Pedon" mean "Official Series Description."

Alvin Series

Taxonomic classification: Coarse-loamy, mixed, superactive, mesic Typic Hapludalfs

Typical Pedon

Alvin fine sandy loam, 2 to 5 percent slopes, at an elevation of about 660 feet; Vermilion County, Illinois; about 2,320 feet south and 1,760 feet east of the northwest corner of sec. 32, T. 21 N., R. 11 W.; USGS Danville NE topographic quadrangle; lat. 40 degrees 14 minutes 8 seconds N. and long. 87 degrees 36 minutes 58 seconds W., NAD 27:

Ap—0 to 8 inches; brown (10YR 4/3) fine sandy loam, light brownish gray (10YR 6/2) dry; weak fine granular structure; very friable; moderately acid; abrupt smooth boundary.

BE—8 to 11 inches; dark yellowish brown (10YR 4/4) fine sandy loam; weak fine subangular blocky structure; very friable; few distinct grayish brown (10YR 5/2) silt coatings on faces of peds; moderately acid; clear smooth boundary.

Bt1—11 to 15 inches; dark yellowish brown (10YR 4/4) fine sandy loam; moderate fine subangular blocky structure; friable; few distinct brown (10YR 4/3) clay films on faces of peds; moderately acid; clear smooth boundary.

Bt2—15 to 25 inches; dark yellowish brown (10YR 4/4) fine sandy loam; moderate medium subangular blocky structure; friable; common distinct brown (10YR 4/3) clay films on faces of peds; strongly acid; clear smooth boundary.

E and Bt—25 to 74 inches; yellowish brown (10YR 5/4) loamy fine sand (E); weak medium subangular blocky structure; very friable; dark yellowish brown (10YR 4/6) fine sandy loam (Bt);

3 to 10 percent of volume occurring as common or many thin lamellae; moderate medium subangular blocky structure; friable; common distinct brown (10YR 4/3) clay films on faces of peds; strongly acid; clear smooth boundary.

C—74 to 80 inches; 80 percent brown (10YR 4/3) and 20 percent yellowish brown (10YR 5/6), stratified fine sandy loam; massive; friable; moderately acid.

Range in Characteristics

Depth to base of diagnostic horizon: 40 to more than 80 inches

Ap or A horizon(s):

Hue—10YR

Value—3 or 4; value of 3 where the A horizon is less than 6 inches thick

Chroma—1 to 4

Texture—fine sandy loam, sandy loam, or very fine sandy loam

E, EB, or BE horizon(s), where present:

Hue—10YR or 7.5YR

Value—4 to 6

Chroma—2 to 4

Texture—very fine sandy loam, fine sandy loam, sandy loam, or loamy fine sand

Bt horizon(s):

Hue—10YR or 7.5YR

Value—4 to 6

Chroma—3 to 6

Texture—very fine sandy loam, loam, fine sandy loam, or sandy loam; also, thin layers of sandy clay loam

E part of E and Bt or Bt and E horizon(s):

Hue—10YR or 7.5YR

Value—4 to 6

Chroma—2 to 6

Texture—sandy loam, loamy sand, sand, or the fine or very fine analogs of those textures

Bt part of E and Bt or Bt and E horizon(s):

Hue—10YR or 7.5YR

Value—4 to 6

Chroma—3 to 6

Texture—sandy loam, loamy sand, the fine or very fine analogs of those textures, or loam

BC or C horizon(s):

Hue—10YR or 7.5YR

Value—4 to 6

Chroma—3 to 6

Texture—sandy loam, loamy sand, sand, or the fine or very fine analogs of those textures

131C2—Alvin fine sandy loam, 5 to 10 percent slopes, eroded

Setting

Landform: Knolls and stream terraces

Position on landform: Summits, backslopes, and risers

Map Unit Composition

Alvin and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have more clay in the subsoil
- Soils that have less clay in the surface soil

Dissimilar soils:

- The well drained Camden soils on summits

Properties and Qualities of the Alvin Soil

Parent material: Eolian deposits and/or outwash

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:
Moderately rapid

Permeability below a depth of 60 inches: Moderately rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 9.0 inches to a depth of 60 inches

Content of organic matter in the surface layer: 0.5 to 1.0 percent

Shrink-swell potential: Low

Flooding: None

Accelerated erosion: The surface layer has been thinned by erosion.

Potential for frost action: Moderate

Corrosivity: Low for steel and high for concrete

Surface runoff class: Low

Susceptibility to water erosion: Moderate

Susceptibility to wind erosion: Moderately high

Interpretive Groups

Land capability classification: Alvin—3e

Prime farmland status: Alvin—prime farmland in all areas

Hydric soil status: Alvin—not hydric

131D2—Alvin fine sandy loam, 10 to 18 percent slopes, eroded

Setting

Landform: Stream terraces and knolls

Position on landform: Summits, risers, and backslopes

Map Unit Composition

Alvin and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have more clay in the subsoil
- Soils that have less clay in the surface soil

Dissimilar soils:

- The well drained Camden soils on summits
- The well drained Middletown soils on summits and shoulders

Properties and Qualities of the Alvin Soil

Parent material: Eolian deposits and/or outwash

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:
Moderately rapid

Permeability below a depth of 60 inches: Moderately rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 9.0 inches to a depth of 60 inches

Content of organic matter in the surface layer: 0.5 to 1.0 percent

Shrink-swell potential: Low

Flooding: None

Accelerated erosion: The surface layer has been thinned by erosion.

Potential for frost action: Moderate

Corrosivity: Low for steel and high for concrete

Surface runoff class: Low

Susceptibility to water erosion: Moderate

Susceptibility to wind erosion: Moderately high

Interpretive Groups

Land capability classification: Alvin—4e

Prime farmland status: Alvin—not prime farmland

Hydric soil status: Alvin—not hydric

Assumption Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Oxyaquic Argiudolls

Typical Pedon (OSD)

Assumption silt loam, 2 to 5 percent slopes, at an elevation of 720 feet; Henry County, Illinois; 100 feet north and 300 feet east of the southwest corner of sec. 29, T. 15 N., R. 2 E.; USGS Andover topographic quadrangle; lat. 41 degrees 15 minutes 00 seconds N. and long. 90 degrees 17 minutes 57 seconds W., NAD 27:

Ap—0 to 6 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; weak medium granular structure; friable; many fine roots throughout; neutral; abrupt smooth boundary.

A—6 to 13 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; moderate medium granular structure; friable; many fine roots throughout; slightly acid; clear smooth boundary.

AB—13 to 16 inches; very dark grayish brown (10YR 3/2) silt loam mixed with some brown (10YR 4/3) material in the lower 2 inches; grayish brown (10YR 5/2) and brown (10YR 5/3) dry; weak medium subangular blocky structure; friable; many fine roots throughout; neutral; clear wavy boundary.

Bt1—16 to 26 inches; brown (10YR 4/3) silty clay loam; weak medium prismatic structure parting to moderate fine subangular blocky; firm; common fine roots between pedes; many moderately thick brown (10YR 5/3) clay films on faces of pedes; slightly acid; clear wavy boundary.

Bt2—26 to 35 inches; brown (10YR 5/3) silty clay loam; weak medium prismatic structure parting to moderate medium subangular blocky; firm; common fine roots between pedes; many distinct brown (10YR 4/3) clay films on faces of pedes; many medium distinct brownish yellow (10YR 6/6) accumulations of iron and common distinct grayish brown (2.5Y 5/2) iron depletions in the matrix; slightly acid; abrupt wavy boundary.

2Bt3—35 to 51 inches; yellowish brown (10YR 5/4) clay loam; weak medium subangular blocky structure; firm; common fine roots between pedes; common distinct moderately thick dark yellowish brown (10YR 4/3) clay films on faces of pedes; many coarse faint yellowish brown (10YR 5/8) accumulations of iron and common medium prominent light olive gray (5Y 6/2) iron depletions; slightly acid; clear wavy boundary.

2Bt4—51 to 60 inches; brown (10YR 5/3) silty clay loam; weak medium prismatic structure parting to moderate medium subangular blocky; firm; common fine roots between pedes; many moderately thick brown (10YR 4/3) clay films on faces of pedes; many medium distinct brownish yellow (10YR 6/6) accumulations of iron; slightly acid; clear wavy boundary.

2C—60 to 80 inches; brown (10YR 5/3) clay loam; massive; firm; common coarse prominent grayish brown (2.5Y 5/2) iron depletions and common coarse distinct brown (7.5YR 4/4) accumulations

of iron in the matrix; slightly effervescent; slightly alkaline.

Range in Characteristics

Thickness of the mollic epipedon: 10 to 20 inches

Thickness of the loess: 20 to 40 inches

Depth to base of diagnostic horizon: 48 to 70 or more inches

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 to 3

Texture—silt loam or silty clay loam

Bt horizon(s):

Hue—10YR or 2.5Y

Value—4 or 5

Chroma—2 to 6

Texture—silty clay loam or silt loam

2Btg or 2Bt horizon(s):

Hue—7.5YR, 10YR, 2.5Y, or 5Y

Value—3 to 6

Chroma—1 to 6

Texture—clay loam, silty clay loam, clay, or silty clay

2C or 2Cg horizon(s):

Hue—7.5YR, 10YR, 2.5Y, or 5Y

Value—3 to 6

Chroma—1 to 6

Texture—clay loam, silty clay loam, clay, or silty clay

Taxadjunct Feature

The Assumption soils have a dark surface soil that is thinner than is definitive for the series. This difference, however, does not significantly affect the use and management of the soils. The soils are classified as fine-silty, mixed, superactive, mesic Oxyaquic Hapludalfs.

259C2—Assumption silt loam, 5 to 10 percent slopes, eroded

Setting

Landform: Ground moraines

Position on landform: Shoulders and backslopes

Map Unit Composition

Assumption and similar soils: 100 percent

Minor Components

Similar soils:

- Soils that have more clay in the surface layer
- Soils that have less sand and less clay in the lower part of the subsoil and in the underlying material
- Soils that have a thicker dark surface soil

Properties and Qualities of the Assumption Soil

Parent material: Loess over a paleosol that formed in till

Drainage class: Moderately well drained

Slowest permeability within a depth of 40 inches: Slow

Permeability below a depth of 60 inches: Slow or moderately slow

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.2 inches to a depth of 60 inches

Content of organic matter in the surface layer: 2.0 to 3.0 percent

Shrink-swell potential: High

Perched seasonal high water table: 2.0 to 3.5 feet below the surface

Flooding: None

Accelerated erosion: The surface layer has been thinned by erosion.

Potential for frost action: High

Corrosivity: High for steel and moderate for concrete

Surface runoff class: Medium

Susceptibility to water erosion: Moderate

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Assumption—3e

Prime farmland status: Assumption—not prime farmland

Hydric soil status: Assumption—not hydric

259D2—Assumption silt loam, 10 to 18 percent slopes, eroded

Setting

Landform: Ground moraines

Position on landform: Shoulders and backslopes

Map Unit Composition

Assumption and similar soils: 100 percent

Minor Components

Similar soils:

- Soils that have more clay in the surface layer
- Soils that have less sand and less clay in the lower part of the subsoil and in the underlying material

- Soils that have a thicker dark surface soil

Properties and Qualities of the Assumption Soil

Parent material: Loess over a paleosol that formed in till

Drainage class: Moderately well drained

Slowest permeability within a depth of 40 inches: Slow

Permeability below a depth of 60 inches: Slow or moderately slow

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.3 inches to a depth of 60 inches

Content of organic matter in the surface layer: 2.0 to 3.0 percent

Shrink-swell potential: High

Perched seasonal high water table: 2.0 to 3.5 feet below the surface

Flooding: None

Accelerated erosion: The surface layer has been thinned by erosion.

Potential for frost action: High

Corrosivity: High for steel and moderate for concrete

Surface runoff class: Medium

Susceptibility to water erosion: High

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Assumption—4e

Prime farmland status: Assumption—not prime farmland

Hydric soil status: Assumption—not hydric

Blackberry Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Oxyaquic Argiudolls

Typical Pedon

Blackberry silt loam, 2 to 5 percent slopes, at an elevation of about 748 feet; Champaign County, Illinois; about 25 feet north and 450 feet west of the southeast corner of sec. 19, T. 21 N., R. 7 E.; USGS Foosland topographic quadrangle; lat. 40 degrees 15 minutes 10 seconds N. and long. 88 degrees 26 minutes 36 seconds W., NAD 27:

Ap—0 to 10 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; moderate medium granular structure; friable; neutral; abrupt smooth boundary.

A—10 to 16 inches; dark brown (10YR 3/3) silt loam, grayish brown (10YR 5/2) dry; moderate medium

granular structure; friable; many faint very dark grayish brown (10YR 3/2) organic coatings on faces of peds; neutral; clear smooth boundary.

BA—16 to 20 inches; brown (10YR 4/3) silty clay loam; weak very fine subangular blocky structure; friable; many faint dark brown (10YR 3/3) organic coatings on faces of peds; slightly acid; clear smooth boundary.

Bt1—20 to 24 inches; yellowish brown (10YR 5/4) silty clay loam; moderate fine subangular blocky structure; friable; many distinct brown (10YR 4/3) clay films on faces of peds; moderately acid; clear smooth boundary.

Bt2—24 to 34 inches; yellowish brown (10YR 5/4) silty clay loam; moderate medium subangular blocky structure; friable; many distinct brown (10YR 4/3) clay films on faces of peds; few fine distinct grayish brown (10YR 5/2) iron depletions in the matrix; few fine irregular black (7.5YR 2.5/1) very weakly cemented nodules of iron and manganese oxide in the matrix; moderately acid; clear smooth boundary.

Bt3—34 to 47 inches; yellowish brown (10YR 5/4) silty clay loam; moderate medium and coarse prismatic structure; friable; common distinct brown (10YR 4/3) clay films on faces of peds; few fine distinct grayish brown (10YR 5/2) iron depletions in the matrix; few fine distinct yellowish brown (10YR 5/6) accumulations of iron in the matrix; few fine irregular black (7.5YR 2.5/1) very weakly cemented nodules of iron and manganese oxide in the matrix; moderately acid; clear smooth boundary.

2Bt4—47 to 62 inches; yellowish brown (10YR 5/4), stratified silt loam and loam; weak coarse subangular blocky structure; friable; very few faint brown (10YR 4/3) and grayish brown (10YR 5/2) clay films lining pores and on faces of peds; common medium distinct light brownish gray (10YR 6/2) iron depletions in the matrix; common medium distinct yellowish brown (10YR 5/6) accumulations of iron in the matrix; few fine irregular black (7.5YR 2.5/1) very weakly cemented nodules of iron and manganese oxide in the matrix; slightly acid; gradual smooth boundary.

2C—62 to 70 inches; light olive brown (2.5Y 5/4), stratified silt loam, loam, and sandy loam; massive; friable; common medium distinct light brownish gray (10YR 6/2) iron depletions in the matrix; common medium distinct yellowish brown (10YR 5/6) accumulations of iron in the matrix; few fine irregular black (7.5YR 2.5/1) very weakly cemented nodules of iron and manganese oxide in the matrix; neutral.

Range in Characteristics

Thickness of the mollic epipedon: 10 to 20 inches

Thickness of the loess: 40 to 60 inches

Depth to carbonates: More than 40 inches

Depth to base of diagnostic horizon: 45 to 70 inches

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 to 3

Texture—silt loam

BA or AB horizon(s), where present:

Hue—10YR

Value—3 or 4

Chroma—2 to 4

Texture—silt loam or silty clay loam

Bt horizon(s):

Hue—7.5YR or 10YR

Value—4 or 5

Chroma—2 to 4

Texture—silty clay loam or silt loam

2Bt or 2BC horizon(s):

Hue—7.5YR or 10YR

Value—4 to 6

Chroma—2 to 6

Texture—clay loam, loam, silt loam, sandy loam, sandy clay loam, or the gravelly analogs of those textures

2C horizon(s):

Hue—7.5YR, 10YR, or 2.5Y

Value—4 to 6

Chroma—2 to 6

Texture—loam, clay loam, loamy sand, sandy loam, silt loam, or the gravelly analogs of those textures

679B—Blackberry silt loam, 2 to 5 percent slopes

Setting

Landform: Stream terraces

Position on landform: Summits and risers

Map Unit Composition

Blackberry and similar soils: 100 percent

Minor Components

Similar soils:

- Soils that have a seasonal high water table at a depth of 1 to 2 feet
- Soils that have a thinner dark surface soil

- Soils that have more sand in the upper part of the subsoil
- Soils that have less sand in the lower part of the subsoil and in the underlying material

Properties and Qualities of the Blackberry Soil

Parent material: Loess over outwash

Drainage class: Moderately well drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderate or moderately rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.8 inches to a depth of 60 inches

Content of organic matter in the surface layer: 3.0 to 5.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: 2.0 to 3.5 feet below the surface

Flooding: None

Potential for frost action: High

Corrosivity: High for steel and moderate for concrete

Surface runoff class: Low

Susceptibility to water erosion: Moderate

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Blackberry—2e

Prime farmland status: Blackberry—prime farmland in all areas

Hydric soil status: Blackberry—not hydric

Broadwell Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Typic Argiudolls

Typical Pedon (OSD)

Broadwell silt loam, 2 to 5 percent slopes, at an elevation of 625 feet; Christian County, Illinois; 2,500 feet north and 460 feet west of the center of sec. 11, T. 15 N., R. 2 W.; USGS Mount Auburn topographic quadrangle; lat. 39 degrees 46 minutes 17 seconds N. and long. 84 degrees 16 minutes 51 seconds W., NAD 27:

Ap—0 to 9 inches; very dark grayish brown (10YR 3/2) silt loam, brown (10YR 5/3) dry; weak fine granular structure; friable; common very fine and few fine roots; moderately acid; clear smooth boundary.

A—9 to 15 inches; very dark grayish brown (10YR 3/2)

silt loam, brown (10YR 5/3) dry; moderate fine granular structure; friable; few very fine and fine roots; slightly acid; clear smooth boundary.

Bt1—15 to 18 inches; dark brown (10YR 3/3) silty clay loam, yellowish brown (10YR 5/4) dry; weak very fine and fine subangular blocky structure; friable; few very fine and fine roots; many faint very dark grayish brown (10YR 3/2) organo-clay films on faces of peds; slightly acid; clear smooth boundary.

Bt2—18 to 25 inches; brown (10YR 4/3) silty clay loam; moderate fine subangular blocky structure; firm; few very fine and fine roots; common faint very dark grayish brown (10YR 3/2) organo-clay films on faces of peds; moderately acid; clear smooth boundary.

Bt3—25 to 31 inches; brown (10YR 4/3) silty clay loam; moderate fine subangular blocky structure; firm; few very fine and fine roots; common faint very dark grayish brown (10YR 3/2) clay films in root channels and/or pores and common faint brown (10YR 4/3) clay films on faces of peds; moderately acid; clear smooth boundary.

Bt4—31 to 41 inches; dark yellowish brown (10YR 4/4) silty clay loam; moderate medium subangular blocky structure; firm; few very fine and fine roots; few distinct very dark grayish brown (10YR 3/2) clay films in root channels and/or pores, common faint brown (10YR 4/3) clay films on faces of peds, and few distinct light gray (10YR 7/2) silt coatings on faces of peds and in pores; moderately acid; clear smooth boundary.

Bt5—41 to 50 inches; dark yellowish brown (10YR 4/4) silt loam; weak medium and coarse subangular blocky structure; friable; few very fine roots; common faint brown (10YR 4/3) clay films on faces of peds; few distinct light gray (10YR 7/2) silt coatings on faces of peds and in pores; few fine faint brown (10YR 5/3) accumulations of iron along micropores; moderately acid; abrupt smooth boundary.

2BC1—50 to 55 inches; dark yellowish brown (10YR 4/4), stratified loamy fine sand and sandy loam; weak coarse subangular blocky structure; very friable; slightly acid; clear smooth boundary.

2BC2—55 to 76 inches; stratified yellowish brown (10YR 5/4) and brown (7.5YR 4/4) fine sand and loamy sand; single grain; loose; few distinct dark brown (7.5YR 3/2) organic coatings in pores; 3-inch band of yellowish brown (10YR 5/6) silt loam starting at a depth of 75 inches; neutral; clear smooth boundary.

2C—76 to 80 inches; yellowish brown (10YR 5/4) fine sand; single grain; slightly acid.

Range in Characteristics

Thickness of the loess: 40 to 60 inches

Thickness of the mollic epipedon: 10 to 24 inches

Depth to base of diagnostic horizon: 45 to 80 inches

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 to 3

Texture—silt loam

Bt horizon(s):

Hue—10YR or 7.5YR

Value—3 to 5

Chroma—3 to 6

Texture—silt loam or silty clay loam

2Bt or 2BC horizon(s):

Hue—10YR or 7.5YR

Value—4 or 5

Chroma—4 to 6

Texture—loamy sand, loamy fine sand, fine sand, sandy loam, or loam

2C horizon(s):

Hue—10YR or 7.5YR

Value—4 or 5

Chroma—4 to 6

Texture—fine sand, sand, or loamy sand

Taxadjunct Feature

Broadwell silt loam, 5 to 10 percent slopes, eroded, has a dark surface soil that is thinner than is definitive for the series. This difference, however, does not significantly affect the use and management of the soil. The soil is classified as a fine-silty, mixed, superactive, mesic Mollic Hapludalf.

684B—Broadwell silt loam, 2 to 5 percent slopes

Setting

Landform: Ground moraines and knolls

Position on landform: Summits and shoulders

Map Unit Composition

Broadwell and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a thinner dark surface soil
- Soils that have more sand in the upper part of the subsoil
- Soils that have more sand in the surface layer

Dissimilar soils:

- The poorly drained Elburn soils on footslopes

Properties and Qualities of the Broadwell Soil

Parent material: Loess over sandy eolian material

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:

Moderate

Permeability below a depth of 60 inches: Rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.6 inches to a depth of 60 inches

Content of organic matter in the surface layer: 3.0 to 4.0 percent

Shrink-swell potential: Moderate

Flooding: None

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Low

Susceptibility to water erosion: Moderate

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Broadwell—2e

Prime farmland status: Broadwell—prime farmland in all areas

Hydric soil status: Broadwell—not hydric

684C2—Broadwell silt loam, 5 to 10 percent slopes, eroded

Setting

Landform: Knolls and ground moraines

Position on landform: Backslopes

Map Unit Composition

Broadwell and similar soils: 100 percent

Minor Components

Similar soils:

- Soils that have a thicker dark surface soil
- Soils that have more sand in the upper part of the subsoil
- Soils that have more sand in the surface layer
- Soils that have more clay in the surface layer

Properties and Qualities of the Broadwell Soil

Parent material: Loess over sandy eolian material

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:

Moderate

Permeability below a depth of 60 inches: Rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.1 inches to a depth of 60 inches

Content of organic matter in the surface layer: 2.0 to 3.0 percent

Shrink-swell potential: Moderate

Flooding: None

Accelerated erosion: The surface layer has been thinned by erosion.

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Medium

Susceptibility to water erosion: Moderate

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Broadwell—3e

Prime farmland status: Broadwell—not prime farmland

Hydric soil status: Broadwell—not hydric

Buckhart Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Oxyaquic Argiudolls

Typical Pedon (OSD)

Buckhart silt loam, 2 to 5 percent slopes, at an elevation of about 603 feet; Christian County, Illinois; approximately 360 feet west and 540 feet north of the southeast corner of sec. 24, T. 14 N., R. 3 W.; USGS Grove City, Illinois, topographic quadrangle; lat. 39 degrees 33 minutes 53 seconds N. and long. 89 degrees 22 minutes 6 seconds W., NAD 27:

Ap—0 to 8 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; moderate medium granular structure; friable; few very fine roots; moderately acid; clear smooth boundary.

A—8 to 15 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; moderate fine subangular blocky structure parting to moderate medium granular; friable; few very fine roots; moderately acid; clear smooth boundary.

Bt1—15 to 26 inches; dark yellowish brown (10YR 4/4) silty clay loam; moderate medium subangular blocky structure parting to moderate medium granular; friable; few very fine roots; many distinct brown (10YR 4/3) clay films on faces of peds and few distinct very dark grayish brown (10YR 3/2) organic coatings in root channels and/or pores; slightly acid; clear smooth boundary.

Bt2—26 to 37 inches; dark yellowish brown (10YR 4/4) silty clay loam; moderate medium subangular blocky structure; friable; few very fine roots; many distinct dark grayish brown (10YR 4/2) clay films

on faces of peds; few fine irregular prominent strong brown (7.5YR 5/6) accumulations of iron and manganese along pores and few fine irregular prominent light brownish gray (2.5Y 6/2) iron depletions along pores; neutral; clear smooth boundary.

Bt3—37 to 52 inches; brown (10YR 5/3) silt loam; weak medium subangular blocky structure; friable; few very fine roots; common distinct dark grayish brown (10YR 4/2) clay films on faces of peds; common fine irregular prominent strong brown (7.5YR 5/6) accumulations of iron and manganese along pores, few fine rounded prominent black (7.5YR 2/1) iron-manganese nodules throughout, and common fine distinct irregular light brownish gray (2.5Y 6/2) iron depletions along pores; slightly acid; clear smooth boundary.

BCt—52 to 67 inches; light olive brown (2.5Y 5/3) silt loam; weak coarse subangular blocky structure; friable; few very fine roots; few distinct dark grayish brown (10YR 4/2) clay films in root channels and/or pores; common fine irregular prominent strong brown (7.5YR 5/6) accumulations of iron and manganese along pores, common fine irregular light brownish gray (2.5Y 6/2) iron depletions along pores, and few fine rounded prominent black (7.5YR 2/1) iron-manganese nodules throughout; neutral; gradual smooth boundary.

C—67 to 80 inches; yellowish brown (10YR 5/4) silt loam; massive; friable; common medium irregular distinct strong brown (7.5YR 5/6) accumulations of iron and manganese throughout, common medium irregular prominent light brownish gray (2.5Y 6/2) iron depletions throughout, and few fine rounded prominent black (7.5YR 2/1) iron-manganese nodules throughout; neutral.

Range in Characteristics

Thickness of the mollic epipedon: 10 to 20 inches

Depth to base of diagnostic horizon: 40 to 55 inches

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 to 3

Texture—silt loam or silty clay loam

Bt or Btg horizon(s):

Hue—10YR or 2.5Y

Value—4 to 6

Chroma—2 to 6

Texture—silt loam or silty clay loam

BC, BCt, or BCg horizon(s):

Hue—10YR or 2.5Y

Value—5 or 6
 Chroma—2 to 4
 Texture—silt loam or silty clay loam

C or Cg horizon(s):

Hue—10YR or 2.5Y
 Value—5 or 6
 Chroma—2 to 6
 Texture—silt loam or silty clay loam

705A—Buckhart silt loam, 0 to 2 percent slopes

Setting

Landform: Ground moraines and knolls
Position on landform: Summits

Map Unit Composition

Buckhart and similar soils: 90 percent
 Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have carbonates within a depth of 40 inches
- Soils that have more clay in the subsoil
- Soils that have a seasonal high water table at a depth of 1 to 2 feet

Dissimilar soils:

- The poorly drained Denny, Sable, and Virden soils in depressions

Properties and Qualities of the Buckhart Soil

Parent material: Loess

Drainage class: Moderately well drained

Slowest permeability within a depth of 40 inches:
 Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.2 inches to a depth of 60 inches

Content of organic matter in the surface layer: 3.0 to 4.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: 2.0 to 3.5 feet below the surface

Flooding: None

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Low

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Buckhart—1

Prime farmland status: Buckhart—prime farmland in all areas

Hydric soil status: Buckhart—not hydric

705B—Buckhart silt loam, 2 to 5 percent slopes

Setting

Landform: Knolls (fig. 2)

Position on landform: Summits and backslopes

Map Unit Composition

Buckhart and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a thinner dark surface soil
- Soils that have carbonates within a depth of 40 inches
- Soils that have a seasonal high water table at a depth of more than 3.5 feet
- Soils that have more clay in the subsoil

Dissimilar soils:

- The poorly drained Sable and Virden soils in depressions

Properties and Qualities of the Buckhart Soil

Parent material: Loess

Drainage class: Moderately well drained

Slowest permeability within a depth of 40 inches:
 Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.0 inches to a depth of 60 inches

Content of organic matter in the surface layer: 3.0 to 4.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: depths 2.0 to 3.5 feet below the surface

Flooding: None

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Low

Susceptibility to water erosion: Moderate

Susceptibility to wind erosion: Low




Figure 2.—Typical pattern of upland prairie soils that formed in loess in nearly level to gently sloping areas.

Interpretive Groups

Land capability classification: Buckhart—2e

Prime farmland status: Buckhart—prime farmland in all areas

Hydric soil status: Buckhart—not hydric

Camden Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Typic Hapludalfs

Typical Pedon

Camden silt loam, 0 to 2 percent slopes, at an elevation of 560 feet; Bureau County, Illinois; 1,280 feet west and 1,740 feet south of the northeast corner of sec. 12, T. 15 N., R. 8 E.; USGS Wyanet topographic quadrangle; lat. 41 degrees 18 minutes 05 seconds N. and long. 89 degrees 30 minutes 52 seconds W., NAD 27:

Ap—0 to 7 inches; dark brown (10YR 4/3) silt loam, pale brown (10YR 6/3) dry; moderate fine granular structure; friable; few fine roots; slightly acid; abrupt smooth boundary.

E—7 to 12 inches; yellowish brown (10YR 5/4) silt loam; weak medium platy structure parting to weak fine subangular blocky; friable; few fine roots; neutral; clear smooth boundary.

Bt1—12 to 18 inches; yellowish brown (10YR 5/6) silty

clay loam; moderate fine subangular blocky structure; friable; few fine roots; common distinct yellowish brown (10YR 5/4) clay films on faces of peds; neutral; clear smooth boundary.

Bt2—18 to 26 inches; yellowish brown (10YR 5/6) silty clay loam; moderate medium prismatic structure parting to moderate medium subangular blocky; friable; few fine roots; many distinct dark yellowish brown (10YR 4/4) clay films on faces of peds; moderately acid; clear smooth boundary.

2Bt3—26 to 34 inches; yellowish brown (10YR 5/6) clay loam; weak medium prismatic structure parting to moderate medium subangular blocky; friable; few fine roots; many distinct dark yellowish brown (10YR 4/4) clay films on faces of peds; moderately acid; clear smooth boundary.

2Bt4—34 to 37 inches; strong brown (7.5YR 5/6) clay loam; weak medium subangular blocky structure; friable; few fine roots; many distinct brown (7.5YR 4/4) clay films on faces of peds; about 7 percent gravel; slightly acid; clear smooth boundary.

2Bt5—37 to 48 inches; strong brown (7.5YR 5/6) sandy clay loam; 1-inch strata of yellowish brown (10YR 5/6) sandy loam; weak medium subangular blocky structure; friable; common distinct brown (7.5YR 4/4) clay films on faces of peds; about 5 percent gravel; slightly acid; clear smooth boundary.

2Bt6—48 to 53 inches; strong brown (7.5YR 5/6)

sandy loam; weak medium subangular blocky; friable; common distinct brown (7.5YR 4/4) clay films bridging sand grains; about 2 percent gravel; neutral; clear wavy boundary.

2C—53 to 60 inches; brown (7.5YR 4/4) sandy loam that has thin strata of loamy sand; single grain; loose; about 5 percent gravel; neutral.

Range in Characteristics

Thickness of the loess: 24 to 40 inches

Depth to base of diagnostic horizon: 30 to 65 inches

Ap horizon(s):

Hue—10YR

Value—3 to 5; value of 3 in horizons less than 6 inches thick

Chroma—2 or 3

Texture—silt loam

E horizon(s):

Hue—10YR

Value—4 to 6

Chroma—2 to 4

Texture—silt loam

Bt horizon(s):

Hue—10YR or 7.5YR

Value—4 or 5

Chroma—3 to 6

Texture—silty clay loam or silt loam

2Bt or 2BC horizon(s):

Hue—10YR, 7.5YR, or 2.5Y

Value—4 to 6

Chroma—3 to 6

Texture—silty clay loam, clay loam, loam, sandy loam, sandy clay loam, or silt loam

2C horizon(s):

Hue—10YR or 7.5YR

Value—4 to 6

Chroma—3 to 6

Texture—stratified sandy loam, loam, silt loam, loamy sand, sandy clay loam, or clay loam

134A—Camden silt loam, 0 to 2 percent slopes

Setting

Landform: Stream terraces

Position on landform: Summits

Map Unit Composition

Camden and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a darker surface layer
- Soils that have less sand in the lower part of the subsoil and in the underlying material
- Soils that have more sand in the upper part of the subsoil
- Soils that are subject to rare flooding

Dissimilar soils:

- The poorly drained Vesser soils on flood plains
- The somewhat poorly drained Kendall soils on summits and footslopes

Properties and Qualities of the Camden Soil

Parent material: Loess over outwash

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderate or moderately rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.9 inches to a depth of 60 inches

Content of organic matter in the surface layer: 1.0 to 3.0 percent

Shrink-swell potential: Moderate

Flooding: None

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Low

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Camden—1

Prime farmland status: Camden—prime farmland in all areas

Hydric soil status: Camden—not hydric

134B—Camden silt loam, 2 to 5 percent slopes

Setting

Landform: Stream terraces (fig. 3)

Position on landform: Summits and riser

Map Unit Composition

Camden and similar soils: 90 percent

Dissimilar soils: 10 percent




Figure 3.—Typical pattern of soils along the Sangamon River.

Minor Components

Similar soils:

- Soils that have more sand in the upper part of the subsoil
- Soils that have less sand in the lower part of the subsoil and in the underlying material
- Soils that are subject to rare flooding
- Soils that have a darker surface layer

Dissimilar soils:

- The somewhat poorly drained Kendall soils on toeslopes

Properties and Qualities of the Camden Soil

Parent material: Loess over outwash

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderate or moderately rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.3 inches to a depth of 60 inches

Content of organic matter in the surface layer: 1.0 to 3.0 percent

Shrink-swell potential: Moderate

Flooding: None

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Low

Susceptibility to water erosion: Moderate

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Camden—2e

Prime farmland status: Camden—prime farmland in all areas

Hydric soil status: Camden—not hydric

134C2—Camden silt loam, 5 to 10 percent slopes, eroded

Setting

Landform: Stream terraces

Position on landform: Shoulders and risers

Map Unit Composition

Camden and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have more sand in the upper part of the subsoil
- Soils that have more sand in the surface soil

- Soils that have less sand in the lower part of the subsoil and in the underlying material
- Soils that have more clay in the surface layer

Dissimilar soils:

- The somewhat poorly drained Kendall soils on summits and footslopes

Properties and Qualities of the Camden Soil

Parent material: Loess over outwash

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:

Moderate

Permeability below a depth of 60 inches: Moderate or moderately rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 10.4 inches to a depth of 60 inches

Content of organic matter in the surface layer: 1.0 to 2.0 percent

Shrink-swell potential: Moderate

Flooding: None

Accelerated erosion: The surface layer has been thinned by erosion.

Potential for frost action: High

Corrosivity: Low for steel and moderate for concrete

Surface runoff class: Medium

Susceptibility to water erosion: High

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Camden—3e

Prime farmland status: Camden—not prime farmland

Hydric soil status: Camden—not hydric

Clarksdale Series

Taxonomic classification: Fine, smectitic, mesic Udollic Endoaqualfs

Typical Pedon (OSD)

Clarksdale silt loam, 0 to 2 percent slopes, at an elevation of 650 feet; Adams County, Illinois; 800 feet south and 550 feet east of the northwest corner of sec. 16, T. 2 N., R. 7 W.; USGS Lorraine, Illinois, topographic quadrangle; lat. 40 degrees 9 minutes 55.1 seconds N. and long. 91 degrees 13 minutes 18 seconds W., NAD 27:

Ap—0 to 8 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; moderate medium granular structure; friable; common fine roots throughout; neutral; abrupt smooth boundary.

E—8 to 12 inches; dark grayish brown (10YR 4/2) silt loam; moderate medium platy structure parting to weak very fine subangular blocky; friable; common very fine and fine roots throughout; many faint very dark grayish brown (10YR 3/2) organic coatings on faces of peds and in pores; few fine prominent yellowish brown (10YR 5/6) accumulations of iron lining root channels and/or pores, few fine distinct black (2.5Y 2/1) accumulations of iron and manganese throughout, and many fine distinct light gray (10YR 7/1 and 7/2) clay depletions between peds; neutral; clear smooth boundary.

BE—12 to 16 inches; grayish brown (10YR 5/2) silt loam; moderate fine subangular blocky structure; friable; few fine roots throughout; common distinct very dark grayish brown (10YR 3/2) organo-clay films on faces of peds and in pores; few fine prominent black (2.5Y 2/1) accumulations of iron and manganese throughout, common fine prominent yellowish brown (10YR 5/6) accumulations of iron throughout, and common fine faint light gray (10YR 7/1) clay depletions between peds; moderately acid; clear smooth boundary.

Bt1—16 to 23 inches; brown (10YR 5/3) silty clay loam; moderate medium prismatic structure parting to moderate medium subangular blocky; firm; few very fine and fine roots throughout; many prominent dark grayish brown (10YR 4/2) clay films on faces of peds and many prominent very dark gray (10YR 3/1) organo-clay films on faces of peds and in pores; common fine prominent black (2.5Y 2/1) accumulations of iron and manganese and common fine distinct yellowish brown (10YR 5/6) accumulations of iron throughout; moderately acid; clear smooth boundary.

Bt2—23 to 31 inches; brown (10YR 5/3) silty clay loam; moderate medium prismatic structure parting to moderate medium subangular blocky; firm; few fine roots throughout; many faint grayish brown (10YR 5/2) clay films on faces of peds and many prominent very dark gray (10YR 3/1) organo-clay films on faces of peds and in pores; many fine distinct yellowish brown (10YR 5/6) accumulations of iron throughout, few fine prominent strong brown (7.5YR 5/6) accumulations of iron throughout, common fine prominent black (2.5Y 2/1) accumulations of iron and manganese throughout, and common fine faint light brownish gray (10YR 6/2) iron depletions throughout; moderately acid; gradual wavy boundary.

Btg1—31 to 47 inches; light brownish gray (2.5Y 6/2) silty clay loam; moderate coarse prismatic structure parting to moderate coarse subangular blocky; firm; few fine roots throughout; common prominent grayish brown (10YR 5/2) clay films on faces of peds and many prominent very dark gray (10YR 3/1) organo-clay films on faces of peds and in pores; many fine and medium prominent strong brown (7.5YR 5/6) accumulations of iron throughout, few fine prominent black (2.5Y 2/1) accumulations of iron and manganese throughout, and few fine faint light brownish gray (10YR 6/2) iron depletions lining root channels and/or pores; neutral; gradual wavy boundary.

Btg2—47 to 57 inches; light brownish gray (2.5Y 6/2) silt loam; weak coarse prismatic structure; firm; few fine roots throughout; common prominent dark grayish brown (10YR 4/2) clay films in root channels and/or pores; many medium prominent strong brown (7.5YR 5/6) accumulations of iron and few fine prominent black (2.5Y 2/1) accumulations of iron and manganese throughout; neutral; clear wavy boundary.

BCg—57 to 67 inches; light brownish gray (2.5Y 6/2) silt loam; weak coarse subangular blocky structure; firm; common prominent dark grayish brown (10YR 4/2) clay films in root channels and/or pores; common medium prominent strong brown (7.5YR 5/6) accumulations of iron and common medium prominent yellowish red (5YR 5/6) accumulations of iron throughout; neutral; clear wavy boundary.

Cg—67 to 80 inches; light brownish gray (10YR 6/2) silt loam; massive; friable; few distinct dark grayish brown (10YR 4/2) clay films in root channels and/or pores; many medium prominent yellowish red (5YR 4/6) accumulations of iron and common medium prominent strong brown (7.5YR 5/6) accumulations of iron throughout; neutral.

Range in Characteristics

Depth to carbonates, where present: More than 40 inches

Depth to base of diagnostic horizon: 40 to 60 inches

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 or 2

Texture—silt loam

E or BE horizon(s):

Hue—10YR

Value—4 to 6

Chroma—1 or 2

Texture—silt loam

Bt or Btg horizon(s):

Hue—10YR, 2.5Y, or 5Y

Value—4 to 6

Chroma—1 to 6

Texture—silty clay loam, silty clay, or silt loam

C or Cg horizon(s):

Hue—10YR, 2.5Y, or 5Y

Value—4 to 6

Chroma—1 to 6

Texture—silt loam

257A—Clarksdale silt loam, 0 to 2 percent slopes

Setting

Landform: Ground moraines (fig. 4)

Position on landform: Summits

Map Unit Composition

Clarksdale and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have less clay in the subsoil
- Soils that have more sand in the lower part of the subsoil and in the underlying material
- Soils that have a lighter colored surface layer

Dissimilar soils:

- The well drained Fayette, Osco, and Rozetta soils on shoulders
- The poorly drained Denny soils in depressions

Properties and Qualities of the Clarksdale Soil

Parent material: Loess

Drainage class: Somewhat poorly drained

Slowest permeability within a depth of 40 inches:

Moderately slow

Permeability below a depth of 60 inches: Moderately slow

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.3 inches to a depth of 60 inches

Content of organic matter in the surface layer: 2.0 to 3.0 percent

Shrink-swell potential: High

Apparent seasonal high water table: 0.5 foot to 2.0 feet below the surface

Flooding: None




Figure 4.—Typical pattern of upland prairie and forest soils that formed in loess or in loess and glacial till in gently sloping to strongly sloping areas.

Potential for frost action: High
Corrosivity: High for steel and moderate for concrete
Surface runoff class: Medium
Susceptibility to water erosion: Low
Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Clarksdale—1
Prime farmland status: Clarksdale—prime farmland where drained
Hydric soil status: Clarksdale—not hydric

Cowden Series

Taxonomic classification: Fine, smectitic, mesic Mollic Albaqualfs

Typical Pedon (OSD)

Cowden silt loam, 0 to 2 percent slopes, at an elevation of about 665 feet; Montgomery County, Illinois; approximately 1,980 feet west and 30 feet north of the southeast corner of sec. 8, T. 9 N., R. 4 W.;

USGS Butler, Illinois, topographic quadrangle; lat. 39 degrees 13 minutes 55 seconds N. and long. 89 degrees 33 minutes 18 seconds W., NAD 27:

Ap—0 to 8 inches; very dark gray (10YR 3/1) silt loam, gray (10YR 5/1) dry; weak fine granular structure; friable; common very fine and few fine roots; few fine continuous tubular pores; few fine irregular dark brown (10YR 3/3) accumulations of iron and manganese; moderately acid; abrupt smooth boundary.

Eg1—8 to 14 inches; dark gray (10YR 4/1) silt loam, gray (10YR 6/1) dry; weak medium platy structure parting to weak fine subangular blocky; friable; few very fine roots; common fine and medium tubular and vesicular pores; many distinct very dark gray (10YR 3/1) organic coatings on faces of peds and filling pores; few fine irregular dark brown (10YR 3/3) accumulations of iron and manganese; moderately acid; clear smooth boundary.

Eg2—14 to 19 inches; gray (10YR 5/1) silt loam, light gray (10YR 7/1) dry; weak medium platy structure parting to weak fine subangular blocky;

friable; few very fine roots; common fine and medium continuous tubular pores; common fine faint grayish brown (10YR 5/2) accumulations of iron in the matrix; common fine irregular dark brown (10YR 3/3) accumulations of iron and manganese; strongly acid; abrupt smooth boundary.

Btg1—19 to 26 inches; grayish brown (10YR 5/2) silty clay loam; moderate fine and medium prismatic structure parting to moderate medium angular and subangular blocky; firm; common very fine roots; few fine continuous tubular pores; common distinct light gray (10YR 7/1 dry) clay depletions on faces of peds in the upper 2 inches; many prominent very dark gray (10YR 3/1) organo-clay films on faces of peds; few fine distinct yellowish brown (10YR 5/4 and 5/6) accumulations of iron in the matrix; common fine and medium irregular black (10YR 2/1) iron-manganese nodules with sharp boundaries; strongly acid; clear smooth boundary.

Btg2—26 to 43 inches; grayish brown (10YR 5/2) silty clay loam; moderate medium prismatic structure parting to moderate medium and coarse angular blocky; firm; few very fine roots; many prominent very dark grayish brown (10YR 3/2) organo-clay films on faces of peds; many medium distinct yellowish brown (10YR 5/6) accumulations of iron in the matrix; common fine and medium irregular black (10YR 2/1) and dark reddish brown (5YR 3/4) iron-manganese nodules with sharp boundaries; moderately acid; gradual smooth boundary.

Btg3—43 to 50 inches; light brownish gray (10YR 6/2) silty clay loam; weak coarse angular blocky structure; firm; few very fine roots; few fine vesicular and tubular pores; few prominent black (10YR 2/1) organo-clay films lining root channels and pores; common distinct dark grayish brown (10YR 4/2) clay films on vertical faces of peds; common coarse prominent yellowish brown (10YR 5/8) accumulations of iron in the matrix; few medium and coarse irregular black (10YR 2/1) iron-manganese nodules with clear strong brown (7.5YR 5/6) boundaries; slightly acid; gradual smooth boundary.

BCtg—50 to 58 inches; gray (10YR 6/1) silt loam; weak medium and coarse angular blocky structure; friable; few very fine roots; few fine vesicular and tubular pores; few prominent very dark gray (10YR 3/1) organo-clay films lining root channels and pores; few distinct dark grayish brown (10YR 4/2) clay films on vertical faces of peds; common coarse prominent strong brown

(7.5YR 5/8) accumulations of iron in the matrix; few fine and medium irregular black (10YR 2/1) iron-manganese nodules with clear strong brown (7.5YR 4/6) boundaries; neutral; clear smooth boundary.

Cg—58 to 69 inches; grayish brown (10YR 5/2) silt loam; massive; friable; few fine and medium vesicular and tubular pores; few prominent very dark gray (10YR 3/1) organo-clay films lining root channels and pores; many medium and coarse prominent strong brown (7.5YR 5/6) accumulations of iron in the matrix; common fine and medium irregular black (5YR 2.5/1) iron-manganese nodules with diffuse yellowish red (5YR 5/6) boundaries; about 8 percent sand; neutral; clear smooth boundary.

2Btgb—69 to 80 inches; grayish brown (10YR 5/2) silty clay loam; moderate fine and medium prismatic structure parting to weak medium angular blocky; firm; common medium and coarse vesicular and tubular pores; few prominent very dark gray (10YR 3/1) organo-clay films lining root channels and pores; common distinct dark grayish brown (10YR 4/2) clay films on faces of peds; common fine and medium distinct yellowish brown (10YR 5/6) accumulations of iron in the matrix; few medium and coarse irregular black (5YR 2.5/1) and yellowish red (5YR 4/6) iron-manganese nodules with clear boundaries; about 15 percent sand and 2 percent pebbles; neutral.

Range in Characteristics

Depth to base of diagnostic horizon: 40 to 65 inches

Thickness of the loess: 55 to more than 80 inches

Profile feature: A B/E horizon in some pedons

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 or 2

Texture—silt loam

Eg horizon(s):

Hue—10YR

Value—4 to 6

Chroma—1 or 2

Texture—silt loam

Btg horizon(s):

Hue—10YR, 2.5Y, or 5Y

Value—4 to 6

Chroma—1 or 2

Texture—silty clay loam, silty clay, or silt loam

Cg horizon and BCtg or BCg horizon(s):

Hue—10YR, 2.5Y, 5Y, or neutral

Value—4 to 6
 Chroma—0 to 2
 Texture—silt loam or silty clay loam

2Cg, 2Ab, 2Btg, or 2Bb horizon(s):

Hue—7.5YR, 10YR, 2.5Y, 5Y, or neutral
 Value—3 to 6
 Chroma—0 to 2
 Texture—silt loam, loam, silty clay loam, or clay loam

112A—Cowden silt loam, 0 to 2 percent slopes

Setting

Landform: Ground moraines
Position on landform: Toeslopes

Map Unit Composition

Cowden and similar soils: 90 percent
 Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a thicker dark surface soil
- Soils that have a lighter colored surface layer
- Soils that have less clay in the subsoil

Dissimilar soils:

- The moderately well drained Harrison soils on summits and backslopes

Properties and Qualities of the Cowden Soil

Parent material: Loess

Drainage class: Poorly drained

Slowest permeability within a depth of 40 inches: Slow

Permeability below a depth of 60 inches: Moderately slow

Depth to restrictive feature: More than 80 inches

Available water capacity: About 10.9 inches to a depth of 60 inches

Content of organic matter in the surface layer: 2.0 to 3.0 percent

Shrink-swell potential: High

Apparent seasonal high water table: At the surface to 1 foot below the surface

Ponding: At the surface to 0.5 foot above the surface

Flooding: None

Potential for frost action: High

Corrosivity: High for steel and moderate for concrete

Surface runoff class: Negligible

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Cowden—2w

Prime farmland status: Cowden—prime farmland where drained

Hydric soil status: Cowden—hydric

Denny Series

Taxonomic classification: Fine, smectitic, mesic Mollic Albaqualfs

Typical Pedon (OSD)

Denny silt loam, 0 to 2 percent slopes, at an elevation of 720 feet; McDonough County, Illinois; 225 feet north and 1,680 feet east of the southwest corner of sec. 25, T. 7 N., R. 3 W.; USGS Good Hope topographic quadrangle; lat. 40 degrees 33 minutes 31 seconds N. and long. 90 degrees 41 minutes 14 seconds W., NAD 27:

Ap—0 to 8 inches; very dark gray (10YR 3/1) silt loam, gray (10YR 5/1) dry; weak fine granular structure; very friable; few very fine roots throughout; moderately acid; abrupt smooth boundary.

Eg1—8 to 14 inches; dark grayish brown (10YR 4/2) silt loam, light brownish gray (10YR 6/2) dry; weak thick platy structure parting to weak thin platy; very friable; few very fine roots throughout; few very fine vesicular pores throughout; few distinct very dark gray (10YR 3/1) organic coatings in root channels; common faint grayish brown (10YR 5/2) clay depletions on faces of peds; common fine prominent dark yellowish brown (10YR 3/6) accumulations of iron and manganese throughout; few fine black (N 2/0) concretions of iron and manganese in the matrix; moderately acid; clear smooth boundary.

Eg2—14 to 21 inches; grayish brown (10YR 5/2) silt loam, light gray (10YR 7/2) dry; weak thick platy structure parting to moderate medium platy; friable; few very fine roots throughout; few fine tubular and few very fine vesicular pores throughout; few distinct very dark gray (10YR 3/1) organic coatings in root channels; common fine faint dark brown (10YR 3/3) accumulations of iron and manganese throughout; common fine black (N 2/0) concretions of iron and manganese in the matrix; moderately acid; abrupt smooth boundary.

Btg1—21 to 29 inches; grayish brown (10YR 5/2) silty clay loam; moderate fine and medium subangular blocky structure; firm; few very fine roots between peds; common distinct dark grayish brown (10YR

4/2) clay films on faces of peds; few distinct very dark gray (10YR 3/1) organic coatings in root channels; many fine prominent dark yellowish brown (10YR 4/6) and common fine distinct yellowish brown (10YR 5/4) accumulations of iron and manganese throughout; common fine black (N 2/0) concretions of iron and manganese in the matrix; moderately acid; clear smooth boundary.

Btg2—29 to 38 inches; grayish brown (10YR 5/2) silty clay loam; moderate medium prismatic structure parting to moderate medium subangular blocky; firm; few very fine roots between peds; common faint dark grayish brown (10YR 4/2) clay films on faces of peds; few distinct very dark gray (10YR 3/1) organic coatings in root channels; many fine prominent dark yellowish brown (10YR 4/6) and common fine prominent yellowish brown (10YR 5/8) accumulations of iron and manganese throughout; common fine (N 2/0) concretions of iron and manganese in the matrix; moderately acid; gradual smooth boundary.

Btg3—38 to 46 inches; light brownish gray (2.5Y 6/2) silty clay loam; moderate coarse prismatic structure parting to moderate coarse subangular blocky; firm; very few fine roots between peds; common distinct dark grayish brown (10YR 4/2) clay films on faces of peds; few distinct very dark gray (10YR 3/1) organic coatings in root channels; many fine prominent dark yellowish brown (10YR 4/6) and common fine prominent strong brown (7.5YR 5/6) accumulations of iron and manganese throughout; common fine black (N 2/0) concretions of iron and manganese in the matrix; moderately acid; gradual wavy boundary.

Cg1—46 to 63 inches; light brownish gray (2.5Y 6/2) silty clay loam; massive; firm; few very fine roots between peds; few very fine vesicular pores throughout; very few distinct very dark gray (10YR 3/1) organic coatings in root channels; many fine prominent dark yellowish brown (10YR 4/6) and common fine prominent strong brown (7.5YR 5/6) accumulations of iron and manganese throughout; few medium black (N 2/0) concretions of iron and manganese in the matrix; slightly acid; diffuse wavy boundary.

Cg2—63 to 80 inches; light brownish gray (2.5Y 6/2) silt loam; massive; firm; many very fine vesicular pores throughout; very few distinct very dark gray (10YR 3/1) organic coatings in root channels; many fine prominent dark yellowish brown (10YR 4/6) and common fine prominent strong brown (7.5YR 5/6) accumulations of iron and manganese throughout; few medium black (N 2/0)

concretions of iron and manganese in the matrix; slightly acid.

Range in Characteristics

Depth to base of diagnostic horizon: 40 to 65 inches

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 or 2

Texture—silt loam

Eg horizon(s):

Hue—10YR or 2.5Y

Value—4 to 6

Chroma—1 or 2

Texture—silt loam

Btg horizon(s):

Hue—10YR, 2.5Y, or 5Y

Value—4 to 6

Chroma—1 or 2

Texture—silty clay loam or silty clay

Cg horizon(s):

Hue—10YR, 2.5Y, or 5Y

Value—4 to 6

Chroma—1 or 2

Texture—silt loam or silty clay loam

45A—Denny silt loam, 0 to 2 percent slopes

Setting

Landform: Depressions (fig. 5)

Map Unit Composition

Denny and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a thicker dark surface soil
- Soils that have a lighter colored surface layer

Dissimilar soils:

- The moderately well drained Buckhart soils on summits and backslopes
- The well drained Osco soils on shoulders and summits

Properties and Qualities of the Denny Soil

Parent material: Loess

Drainage class: Poorly drained

Slowest permeability within a depth of 40 inches: Slow




Figure 5.—Typical pattern of upland prairie soils that formed in loess in gently sloping and moderately sloping areas.

Permeability below a depth of 60 inches: Moderately slow
Depth to restrictive feature: More than 80 inches
Available water capacity: About 11.6 inches to a depth of 60 inches
Content of organic matter in the surface layer: 2.0 to 3.0 percent
Shrink-swell potential: High
Apparent seasonal high water table: At the surface to 1 foot below the surface
Ponding: At the surface to 1 foot above the surface
Flooding: None
Potential for frost action: High
Corrosivity: High for steel and moderate for concrete
Surface runoff class: Negligible
Susceptibility to water erosion: Low
Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Denny—3w
Prime farmland status: Denny—prime farmland where drained
Hydric soil status: Denny—hydric

Drury Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Dystric Eutrudepts

Typical Pedon (OSD)

Drury silt loam, 0 to 2 percent slopes, at an elevation of about 465 feet; Monroe County, Illinois; approximately 2,380 feet southeast of intersection of Bluff Road and railroad crossing and 820 feet northeast of railroad tracks, parcel S. 701, C. 495, T. 3 S., R. 11 W.; USGS Selma, Illinois-Missouri, topographic quadrangle; lat. 38 degrees 13 minutes 52 seconds N. and long. 90 degrees 16 minutes 54 seconds W., NAD 27:

Ap—0 to 7 inches; brown (10YR 4/3) silt loam, pale brown (10YR 6/3) dry; moderate medium granular structure; friable; common very fine and few fine roots; few fine continuous tubular pores; neutral; abrupt smooth boundary.

Bw1—7 to 12 inches; dark yellowish brown (10YR 4/4) silt loam; moderate fine subangular blocky structure; friable; few very fine and fine roots; few

medium continuous tubular pores; many faint dark brown (10YR 3/3) organo-clay films on faces of peds and lining vertical tubular pores; neutral; clear smooth boundary.

Bw2—12 to 19 inches; dark yellowish brown (10YR 4/4) silt loam; weak fine prismatic structure parting to moderate fine subangular blocky; friable; few very fine and fine roots; common fine continuous tubular pores; common faint dark brown (10YR 3/3) organo-clay films on faces of peds and lining vertical tubular pores; neutral; gradual smooth boundary.

Bw3—19 to 26 inches; dark yellowish brown (10YR 4/4) silt loam; weak medium prismatic structure parting to moderate fine and medium subangular blocky; friable; few very fine and fine roots; common fine continuous tubular pores; common faint dark brown (10YR 3/3) organo-clay films on faces of peds and lining vertical tubular pores; neutral; gradual smooth boundary.

Bw4—26 to 36 inches; dark yellowish brown (10YR 4/4) silt loam; weak medium prismatic structure parting to moderate medium subangular blocky; friable; few very fine and fine roots; common fine continuous tubular pores; few faint dark brown (10YR 3/3) organo-clay films on faces of peds and lining vertical tubular pores; neutral; gradual smooth boundary.

Bw5—36 to 43 inches; dark yellowish brown (10YR 4/4) silt loam; weak medium prismatic structure parting to weak medium subangular blocky; very friable; few very fine roots; common fine continuous tubular pores; few faint dark brown (10YR 3/3) organo-clay films on faces of peds and lining vertical tubular pores; neutral; gradual smooth boundary.

C1—43 to 70 inches; dark yellowish brown (10YR 4/4) silt loam; massive; very friable; few very fine and fine continuous tubular pores; few fine distinct yellowish brown (10YR 5/6) accumulations of iron in the matrix; few fine rounded black (N 2.5/0) accumulations of iron and manganese; neutral; gradual smooth boundary.

C2—70 to 80 inches; dark yellowish brown (10YR 4/4) silt loam; massive; friable; few very fine continuous pores; few fine rounded black (N 2.5/0) accumulations of iron and manganese; neutral.

Range in Characteristics

Depth to base of diagnostic horizon: 26 to 55 inches

Ap or A horizon(s):

Hue—10YR

Value—3 or 4

Chroma—2 to 4

Texture—silt loam or silt

E horizon(s), where present:

Hue—10YR

Value—4 or 5 (6 or 7 dry)

Chroma—3 or 4

Texture—silt loam or silt

Bw horizon(s):

Hue—7.5YR or 10YR

Value—4 or 5

Chroma—3 to 6 in the upper part, 2 to 6 in the lower part

Texture—silt loam

C horizon(s):

Hue—10YR

Value—3 to 6

Chroma—2 to 4

Texture—silt loam

7075B—Drury silt loam, 2 to 5 percent slopes, rarely flooded

Setting

Landform: Alluvial fans

Map Unit Composition

Drury and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have more clay in the subsoil
- Soils that have more than 10 percent gravel throughout
- Soils that have a darker surface layer

Dissimilar soils:

- The poorly drained Vesser soils on flood plains

Properties and Qualities of the Drury Soil

Parent material: Silty local alluvium

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:

Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.0 inches to a depth of 60 inches

Content of organic matter in the surface layer: 1.0 to 3.0 percent

Shrink-swell potential: Low

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Low

Susceptibility to water erosion: Moderate

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Drury—2e

Prime farmland status: Drury—prime farmland in all areas

Hydric soil status: Drury—not hydric

536—Dumps, mine

Setting

This map unit is in areas where fly ash from the coal-fired power plant in the southeast corner of the city of Springfield has accumulated.

Map Unit Composition

Dumps, mine: 95 percent

Dissimilar components: 15 percent

Minor Components

Dissimilar components:

- The well drained, silty Orthents along the edge of the refuse area
- Areas of undisturbed soils along the edge of the refuse area
- Areas of water less than 3 acres in size

Interpretive Groups

Land capability classification: Dumps—none assigned

Prime farmland status: Dumps—not prime farmland

Hydric soil status: Dumps—unranked

Edinburg Series

Taxonomic classification: Fine, smectitic, mesic Vertic Argiaquolls

Typical Pedon (OSD)

Edinburg silty clay loam, 0 to 2 percent slopes, at an elevation of 615 feet; Sangamon County, Illinois; 1,200 feet south and 276 feet east of the center of sec. 22, T. 14 N., R. 6 W.; USGS Chatham, Illinois, topographical quadrangle; lat. 39 degrees 38 minutes 37 seconds N. and long. 89 degrees 45 minutes 00 seconds W., NAD 27:

Ap—0 to 8 inches; very dark gray (10YR 3/1) silty clay loam, gray (10YR 5/1) dry; moderate fine and medium granular structure; friable; common fine and very fine roots; neutral; abrupt smooth boundary.

A—8 to 10 inches; very dark gray (10YR 3/1) silty clay loam, gray (10YR 5/1) dry; moderate medium granular structure with some moderate very fine subangular blocky; firm; common fine and very fine roots; neutral; clear smooth boundary.

BE—10 to 16 inches; very dark gray (10YR 3/1) silty clay loam, gray (10YR 5/1) dry; weak and moderate fine subangular blocky structure; firm; common very fine and few fine roots; few fine faint very dark grayish brown (2.5Y 3/2) and dark grayish brown (2.5Y 4/2) iron depletions; few fine prominent olive brown (2.5Y 4/4) accumulations of iron and manganese; few distinct light brownish gray (10YR 6/2) clay depletions on faces of peds; few fine yellowish brown (10YR 5/8) iron and manganese concretions; neutral; clear smooth boundary.

Btg1—16 to 20 inches; dark gray (10YR 4/1) silty clay loam; moderate fine angular blocky structure; firm; few very fine roots; many distinct very dark gray (10YR 3/1) organo-clay films on faces of peds; few fine prominent olive brown (2.5Y 4/4) accumulations of iron and manganese; few fine iron and manganese concretions; neutral; gradual smooth boundary.

Btg2—20 to 26 inches; dark gray (10YR 4/1) silty clay; moderate medium prismatic structure parting to moderate medium and coarse angular blocky; firm; few very fine roots; common distinct dark gray (10YR 4/1) clay films on faces of peds; few fine prominent olive brown (2.5Y 4/4) accumulations of iron and manganese; common fine iron and manganese concretions; slightly acid; gradual smooth boundary.

Btg3—26 to 34 inches; dark gray (10YR 4/1) silty clay loam; moderate medium prismatic structure parting to moderate coarse angular blocky; firm; few very fine roots; common distinct dark gray (10YR 4/1) clay films and few faint very dark gray (10YR 3/1) organo-clay films on faces of peds; common fine prominent yellowish brown (10YR 5/6 and 5/8) accumulations of iron and manganese; few fine iron and manganese concretions; neutral; gradual smooth boundary.

Btg4—34 to 41 inches; olive gray (5Y 5/2) silty clay loam; moderate medium prismatic structure parting to moderate coarse angular blocky; firm; few very fine roots; few distinct dark gray (10YR 4/1) clay films and very dark gray (10YR 3/1) organo-clay films on vertical faces of peds; many fine and medium prominent yellowish brown (10YR 5/6 and 5/8) accumulations of iron and manganese; few fine iron and manganese concretions; neutral; gradual smooth boundary.

BCg—41 to 55 inches; olive gray (5Y 5/2) silty clay loam; weak medium prismatic structure parting to weak coarse angular blocky; friable; few faint dark gray (10YR 4/1) clay films and very dark gray (10YR 3/1) organo-clay films on vertical faces of peds; many fine and medium prominent yellowish brown (10YR 5/6 and 5/8) accumulations of iron and manganese; few fine iron and manganese concretions; neutral; gradual smooth boundary.

C—55 to 60 inches; mottled yellowish brown (10YR 5/6) and light olive gray (5Y 6/2) silt loam; massive; friable; dark gray (10YR 4/1) clay films in root channels; neutral.

Range in Characteristics

Thickness of the mollic epipedon: 10 to 24 inches

Depth to base of diagnostic horizon: 40 to 65 inches

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 or 2

Texture—silty clay loam or silt loam

BE, BEg, or Eg horizon(s):

Hue—10YR

Value—3 or 4

Chroma—1 or 2

Texture—silty clay loam or silt loam

Btg horizon(s):

Hue—10YR, 2.5Y, or 5Y

Value—3 to 6

Chroma—1 or 2

Texture—silty clay loam or silty clay

C or Cg horizon(s):

Hue—10YR, 2.5Y, or 5Y

Value—5 or 6

Chroma—1 or 2

Texture—silt loam or silty clay loam

249A—Edinburg silty clay loam, 0 to 2 percent slopes

Setting

Landform: Depressions

Map Unit Composition

Edinburg and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have less clay in the subsoil

- Soils that have a dark surface soil more than 24 inches thick
- Soils that have a gray subsurface layer

Dissimilar soils:

- The moderately well drained Buckhart soils on summits

Properties and Qualities of the Edinburg Soil

Parent material: Loess

Drainage class: Poorly drained

Slowest permeability within a depth of 40 inches:
Slow

Permeability below a depth of 60 inches: Moderately slow or moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.3 inches to a depth of 60 inches

Content of organic matter in the surface layer: 3.0 to 6.0 percent

Shrink-swell potential: High

Apparent seasonal high water table: At the surface to 1 foot below the surface

Ponding: At the surface to 0.5 foot above the surface

Flooding: None

Potential for frost action: High

Corrosivity: High for steel and moderate for concrete

Surface runoff class: Negligible

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Very low

Interpretive Groups

Land capability classification: Edinburg—3w

Prime farmland status: Edinburg—prime farmland where drained

Hydric soil status: Edinburg—hydric

Elburn Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Aquic Argiudolls

Typical Pedon (OSD)

Typical pedon Elburn silt loam, 0 to 2 percent slopes, at an elevation of about 617 feet; Christian County, Illinois; 2,716 feet north and 1,300 feet west of the southeast corner of sec. 36, T. 14 N. R. 1 E.; USGS Assumption, Illinois, topographic quadrangle; lat. 39 degrees 37 minutes 4.7 seconds N., and long. 89 degrees 1 minute 45.8 seconds W., NAD 27:

Ap—0 to 6 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; weak fine granular structure; friable; few very fine roots; many distinct very dark gray (10YR 3/1) organic

- coatings on faces of peds; slightly acid; abrupt smooth boundary.
- A—6 to 16 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; moderate fine granular structure; friable; few very fine roots; many distinct very dark gray (10YR 3/1) organic coatings on faces of peds; neutral; clear smooth boundary.
- Bt1—16 to 21 inches; brown (10YR 4/3) silty clay loam; moderate fine subangular blocky structure; friable; few very fine roots; many distinct very dark gray (10YR 3/1) organo-clay films and dark gray (10YR 4/1) clay films on faces of peds; few fine prominent yellowish brown (10YR 5/8) accumulations of iron and few fine faint brown (10YR 5/3) accumulations of iron and manganese in the matrix; few fine prominent concretions of iron and manganese throughout; slightly acid; clear smooth boundary.
- Bt2—21 to 28 inches; brown (10YR 5/3) silty clay loam; moderate fine subangular blocky structure; firm; few very fine roots; common distinct very dark gray (10YR 3/1) organo-clay films and common faint dark grayish brown (10YR 4/2) clay films on faces of peds; few fine faint grayish brown (10YR 5/2) iron depletions and few fine distinct yellowish brown (10YR 5/6) accumulations of iron in the matrix; few fine prominent concretions of iron and manganese throughout; neutral; clear smooth boundary.
- Bt3—28 to 36 inches; brown (10YR 5/3) silty clay loam; moderate medium subangular blocky structure; firm; few very fine roots; common distinct very dark gray (10YR 3/1) organo-clay films and dark gray (10YR 4/1) clay films on faces of peds; common fine faint grayish brown (10YR 5/2) iron depletions and common fine distinct yellowish brown (10YR 5/6) accumulations of iron in the matrix; few fine prominent concretions of iron and manganese throughout; neutral; clear smooth boundary.
- Bt4—36 to 43 inches; light olive brown (2.5Y 5/4) silty clay loam; moderate medium subangular blocky structure; friable; few very fine roots; few prominent very dark gray (10YR 3/1) organo-clay films and few distinct brown (10YR 5/3) clay films on faces of peds; common medium distinct yellowish brown (10YR 5/6) and brownish yellow (10YR 6/6) accumulations of iron in the matrix; few fine prominent concretions of iron and manganese throughout; slightly alkaline; clear smooth boundary.
- Btg1—43 to 49 inches; grayish brown (2.5Y 5/2) silty clay loam; weak coarse subangular blocky

structure; friable; few very fine roots; few distinct very dark gray (10YR 3/1) organo-clay films and dark grayish brown (10YR 4/2) clay films on faces of peds; many medium prominent brownish yellow (10YR 6/8) and few fine prominent yellowish brown (10YR 5/8) accumulations of iron in the matrix; few fine prominent concretions of iron and manganese throughout; slightly alkaline; clear smooth boundary.

2BCtg—49 to 58 inches; grayish brown (2.5Y 5/2), stratified silt loam, loam, and sandy loam; weak coarse subangular blocky structure; friable; few very fine roots; few distinct very dark grayish brown (10YR 3/2) organo-clay films and dark grayish brown (10YR 4/2) clay films lining pores; common medium prominent brownish yellow (10YR 6/8) and few fine prominent yellowish brown (10YR 5/8) accumulations of iron in the matrix; few very fine concretions of iron and manganese throughout; slightly alkaline; clear smooth boundary.

2Cg—58 to 62 inches; grayish brown (2.5Y 5/2), stratified sandy loam and loamy sand; massive; very friable; common medium prominent yellowish brown (10YR 5/8) and brownish yellow (10YR 6/8) accumulations of iron in the matrix; slightly alkaline.

Range in Characteristics

Thickness of the mollic epipedon: 10 to 19 inches

Thickness of the loess: 40 to 60 inches

Depth to base of diagnostic horizon: 40 to 70 inches

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 or 2

Texture—silt loam

Bt or Btg horizon(s):

Hue—10YR, 2.5Y, or 5Y

Value—4 or 5

Chroma—2 to 4

Texture—silty clay loam or silt loam

2Btg, 2Bt, 2Bg, 2BC, 2BCtg, or 2BCg horizon(s):

Hue—7.5YR, 10YR, 2.5Y, or 5Y

Value—4 to 6

Chroma—2 to 8

Texture—sandy loam, clay loam, loam, silty clay loam, or silt loam

2C or 2Cg horizon(s):

Hue—7.5YR, 10YR, 2.5Y, or 5Y

Value—4 to 6

Chroma—2 to 8

Texture—sandy loam, loam, loamy sand, sand, or silt loam

198A—Elburn silt loam, 0 to 2 percent slopes

Setting

Landform: Stream terraces (fig. 3)

Position on landform: Summits and footslopes

Map Unit Composition

Elburn and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have more sand in the upper part of the subsoil
- Soils that have a thinner dark surface layer
- Soils that have less sand in the lower part of the subsoil and in the underlying material

Dissimilar soils:

- The poorly drained Vesser soils on flood plains
- The well drained Plano soils on summits

Properties and Qualities of the Elburn Soil

Parent material: Loess over outwash

Drainage class: Somewhat poorly drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderately rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.9 inches to a depth of 60 inches

Content of organic matter in the surface layer: 3.5 to 5.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: 1 to 2 feet below the surface

Flooding: None

Potential for frost action: High

Corrosivity: High for steel and moderate for concrete

Surface runoff class: Low

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Elburn—1

Prime farmland status: Elburn—prime farmland in all areas

Hydric soil status: Elburn—not hydric

Elco Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Oxyaquic Hapludalfs

Typical Pedon (OSD)

Elco silt loam, 10 to 18 percent slopes, at an elevation of about 575 feet; Sangamon County, Illinois; 2,520 feet east and 2,200 feet south of the northwest corner of sec. 35, T. 15 N., R. 4 W.; USGS New City, Illinois, topographic quadrangle; lat. 39 degrees 42 minutes 30 seconds N. and long. 89 degrees 30 minutes 28 seconds W., NAD 27:

Ap—0 to 4 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; strong very fine granular structure; friable; many roots throughout; slightly acid; clear smooth boundary.

E—4 to 12 inches; brown (10YR 4/3) silt loam; weak thin platy structure parting to moderate very fine granular; friable; many distinct light gray (10YR 7/1 dry) clay depletions on faces of peds; few faint very dark grayish brown (10YR 3/2) organic coatings on faces of peds and in pores; few distinct yellowish brown (10YR 5/4) flecks and fragments of subsoil material; slightly acid; clear smooth boundary.

BE—12 to 15 inches; yellowish brown (10YR 5/4) silt loam; moderate very fine and fine subangular blocky structure; friable; few faint dark brown (10YR 3/3) organo-clay films and very few faint dark grayish brown (10YR 4/2) clay films on faces of peds; few distinct light gray (10YR 7/1 dry) clay depletions on faces of peds; few fine black (5YR 2.5/1) concretions of iron and manganese throughout; slightly acid; clear smooth boundary.

Bt—15 to 26 inches; yellowish brown (10YR 5/4) silty clay loam; moderate fine and medium subangular blocky structure; firm; many distinct brown (10YR 4/3) clay films on faces of peds; few distinct light gray (10YR 7/1 dry) silt coatings on faces of peds; few fine distinct yellowish brown (10YR 5/6) accumulations of iron in the matrix; few fine distinct grayish brown (10YR 5/2) iron depletions along micropores; few fine black (5YR 2.5/1) concretions of iron and manganese throughout; slightly acid; clear smooth boundary.

2Btg1—26 to 39 inches; grayish brown (2.5Y 5/2) and yellowish brown (10YR 5/6) silty clay loam; moderate medium and coarse subangular and angular blocky structure; firm; common distinct olive brown (2.5Y 4/4) and brown (10YR 4/3) clay films on faces of peds; few fine prominent strong brown (7.5YR 5/8) accumulations of iron in the

matrix; common very fine black (5YR 2.5/1) concretions of iron and manganese throughout; slightly acid; gradual smooth boundary.

3Btg2—39 to 55 inches; grayish brown (2.5Y 5/2) and yellowish brown (10YR 5/6) silty clay; weak medium prismatic structure parting to moderate coarse subangular and angular blocky; firm; many distinct gray (5Y 5/1) clay films on faces of peds; few fine prominent strong brown (7.5YR 5/8) accumulations of iron in the matrix; few fine black (5YR 2.5/1) concretions of iron and manganese throughout; slightly acid; clear smooth boundary.

3Btg3—55 to 70 inches; grayish brown (2.5Y 5/2) silty clay; moderate fine and medium subangular and angular blocky structure; friable; common distinct gray (5Y 5/1) clay films on faces of peds and in pores; common fine prominent strong brown (7.5YR 5/6) accumulations of iron in the matrix; few fine black (5YR 2.5/1) concretions of iron and manganese throughout; slightly acid; clear smooth boundary.

3Btg4—70 to 80 inches; gray (5Y 5/1) silty clay; moderate coarse subangular blocky structure; firm; common distinct greenish gray (5GY 5/1) clay films on faces of peds; few distinct black (10YR 2/1) organic coatings in root channels and pores; many fine strong brown (7.5YR 4/6) accumulations of iron in the matrix; few fine black (5YR 2.5/1) concretions of iron and manganese throughout; slightly alkaline.

Range in Characteristics

Thickness of the loess: 20 to 40 inches

Depth to base of diagnostic horizon: More than 48 inches

Ap or A horizon(s):

Hue—10YR

Value—3 or 4

Chroma—1 or 2

Texture—silt loam or silty clay loam

E horizon(s):

Hue—10YR

Value—4 or 5

Chroma—3 or 4

Texture—silt loam

BE horizon(s):

Hue—10YR

Value—4 or 5

Chroma—3 to 6

Texture—silt loam or silty clay loam

Bt horizon(s):

Hue—10YR or 7.5YR

Value—4 or 5

Chroma—2 to 6

Texture—silty clay loam or silt loam

2Btg horizon or, where present, 2Bt horizon(s):

Hue—5Y, 2.5Y, 10YR, or 7.5YR

Value—3 to 6

Chroma—1 to 6

Texture—loam, clay loam, silty clay loam, or silt loam

3Btg horizon or, where present, 3Bt horizon(s):

Hue—5Y, 2.5Y, 10YR, or 7.5YR

Value—3 to 6

Chroma—1 to 6

Texture—loam, clay loam, silty clay loam, silty clay, or clay

119C2—Elco silt loam, 5 to 10 percent slopes, eroded

Setting

Landform: Ground moraines (fig. 4)

Position on landform: Shoulders and backslopes

Map Unit Composition

Elco and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have less sand in the lower part of the subsoil
- Soils that have more sand in the upper part of the subsoil
- Soils that have more clay in the surface layer

Dissimilar soils:

- Somewhat poorly drained soils that are on backslopes and have more clay in the upper part of the subsoil

Properties and Qualities of the Elco Soil

Parent material: Loess over a paleosol that formed in till

Drainage class: Moderately well drained

Slowest permeability within a depth of 40 inches: Slow

Permeability below a depth of 60 inches: Slow or moderately slow

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.3 inches to a depth of 60 inches

Content of organic matter in the surface layer: 1.0 to 2.0 percent

Shrink-swell potential: High
Perched seasonal high water table: 2.0 to 3.5 feet below the surface
Flooding: None
Accelerated erosion: The surface layer has been thinned by erosion.
Potential for frost action: High
Corrosivity: High for steel and moderate for concrete
Surface runoff class: Medium
Susceptibility to water erosion: High
Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Elco—3e
Prime farmland status: Elco—not prime farmland
Hydric soil status: Elco—not hydric

119D—Elco silt loam, 10 to 18 percent slopes

Setting

Landform: Ground moraines
Position on landform: Shoulders and backslopes

Map Unit Composition

Elco and similar soils: 90 percent
 Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have less sand in the lower part of the subsoil
- Soils that have more sand in the upper part of the subsoil
- Soils that have more clay in the surface layer

Dissimilar soils:

- Somewhat poorly drained soils that are on backslopes and have more clay in the upper part of the subsoil

Properties and Qualities of the Elco Soil

Parent material: Loess over a paleosol that formed in till
Drainage class: Moderately well drained
Slowest permeability within a depth of 40 inches: Slow
Permeability below a depth of 60 inches: Slow or moderately slow
Depth to restrictive feature: More than 80 inches
Available water capacity: About 11.3 inches to a depth of 60 inches
Content of organic matter in the surface layer: 1.0 to 3.0 percent
Shrink-swell potential: High

Perched seasonal high water table: 2.0 to 3.5 feet below the surface
Flooding: None
Potential for frost action: High
Corrosivity: High for steel and moderate for concrete
Surface runoff class: Medium
Susceptibility to water erosion: High
Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Elco—3e
Prime farmland status: Elco—not prime farmland
Hydric soil status: Elco—not hydric

119D2—Elco silt loam, 10 to 18 percent slopes, eroded

Setting

Landform: Ground moraines
Position on landform: Shoulders and backslopes

Map Unit Composition

Elco and similar soils: 90 percent
 Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have less sand in the lower part of the subsoil
- Soils that have more sand in the upper part of the subsoil
- Soils that have more clay in the surface layer

Dissimilar soils:

- Somewhat poorly drained soils that are on backslopes and have more clay in the upper part of the subsoil
- The well drained Thebes soils on backslopes

Properties and Qualities of the Elco Soil

Parent material: Loess over a paleosol that formed in till
Drainage class: Moderately well drained
Slowest permeability within a depth of 40 inches: Slow
Permeability below a depth of 60 inches: Slow or moderately slow
Depth to restrictive feature: More than 80 inches
Available water capacity: About 11.2 inches to a depth of 60 inches
Content of organic matter in the surface layer: 1.0 to 2.0 percent
Shrink-swell potential: High
Perched seasonal high water table: 2.0 to 3.5 feet below the surface

Flooding: None

Accelerated erosion: The surface layer has been thinned by erosion.

Potential for frost action: High

Corrosivity: High for steel and moderate for concrete

Surface runoff class: Medium

Susceptibility to water erosion: High

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Elco—3e

Prime farmland status: Elco—not prime farmland

Hydric soil status: Elco—not hydric

119D3—Elco silty clay loam, 10 to 18 percent slopes, severely eroded

Setting

Landform: Ground moraines

Position on landform: Shoulders and backslopes

Map Unit Composition

Elco and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have less sand in the lower part of the subsoil
- Soils that have more sand in the upper part of the subsoil
- Soils that have less clay in the surface layer

Dissimilar soils:

- Somewhat poorly drained soils that are on backslopes and have more clay in the upper part of the subsoil
- The well drained Marseilles soils on backslopes

Properties and Qualities of the Elco Soil

Parent material: Loess over a paleosol that formed in till

Drainage class: Moderately well drained

Slowest permeability within a depth of 40 inches: Slow

Permeability below a depth of 60 inches: Slow or moderately slow

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.0 inches to a depth of 60 inches

Content of organic matter in the surface layer: 0.5 to 1.0 percent

Shrink-swell potential: High

Perched seasonal high water table: 2.0 to 3.5 feet below the surface

Flooding: None

Accelerated erosion: The surface layer is mostly subsoil material.

Potential for frost action: High

Corrosivity: High for steel and moderate for concrete

Surface runoff class: Medium

Susceptibility to water erosion: High

Susceptibility to wind erosion: Very low

Interpretive Groups

Land capability classification: Elco—4e

Prime farmland status: Elco—not prime farmland

Hydric soil status: Elco—not hydric

Elkhart Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Typic Argiudolls

Typical Pedon (OSD)

Elkhart silt loam, 5 to 10 percent slopes, eroded, at an elevation of 570 feet; Logan County, Illinois; 2,060 feet south and 1,248 feet west of the northeast corner of sec. 32, T. 19 N., R. 3 W.; USGS Broadwell topographic quadrangle; lat. 40 degrees 03 minutes 26 seconds N. and long. 89 degrees 26 minutes 58 seconds W., NAD 27:

Ap—0 to 8 inches; very dark grayish brown (10YR 3/2) silt loam, dark grayish brown (10YR 4/2) dry; weak fine and medium granular structure; friable; common very fine roots; slightly acid; abrupt smooth boundary.

A—8 to 10 inches; very dark grayish brown (10YR 3/2) silt loam, dark grayish brown (10YR 4/2) dry; moderate fine granular structure; friable; common very fine roots; slightly acid; clear smooth boundary.

BA—10 to 15 inches; dark brown (10YR 3/3) silty clay loam, brown (10YR 4/3) dry; moderate very fine and fine subangular blocky structure; friable; common very fine roots; common faint very dark grayish brown (10YR 3/2) organic coatings on faces of peds; slightly acid; clear smooth boundary.

Bt1—15 to 22 inches; dark yellowish brown (10YR 4/4) silty clay loam; moderate fine subangular blocky structure; firm; few very fine roots; common distinct very dark grayish brown (10YR 3/2) organo-clay films on faces of peds; slightly acid; clear smooth boundary.

Bt2—22 to 28 inches; dark yellowish brown (10YR 4/4) silty clay loam; moderate fine and medium subangular blocky structure; firm; few very fine

roots; few distinct dark brown (10YR 3/3) organo-clay films on faces of peds; slightly acid; clear smooth boundary.

BCt—28 to 31 inches; yellowish brown (10YR 5/4) silty clay loam; weak medium and coarse subangular blocky structure; friable; few very fine roots; few faint brown (10YR 4/3) clay films on faces of peds; few fine black (5YR 2.5/1) manganese concretions with diffuse boundaries in ped interiors; neutral; clear smooth boundary.

C—31 to 60 inches; yellowish brown (10YR 5/4) silt loam; massive; friable; few very fine roots in the upper 10 inches; common fine prominent strong brown (7.5YR 5/8) accumulations of iron in ped interiors; common medium distinct gray (10YR 6/1) iron depletions along root channels and pores; strongly effervescent; moderately alkaline.

Range in Characteristics

Thickness of the mollic epipedon: 10 to 20 inches

Depth to base of diagnostic horizon: 20 to 40 inches

Depth to carbonates: 20 to 40 inches

Ap, A, or AB horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 to 3

Texture—silt loam or silty clay loam

BA or Bt horizon(s):

Hue—10YR or 7.5YR

Value—3 to 5

Chroma—3 to 6

Texture—silty clay loam or silt loam

BC or BCt horizon(s):

Hue—7.5YR, 10YR, or 2.5Y

Value—4 to 6

Chroma—3 to 6

Texture—silt loam or silty clay loam

C horizon(s):

Hue—10YR, 2.5Y, or 5Y

Value—4 to 6

Chroma—1 to 6

Texture—silt or silt loam

Taxadjunct Feature

The Elkhart soils have a dark surface soil that is thinner than is definitive for the series. This difference, however, does not significantly affect the use and management of the soils. The soils are classified as fine-silty, mixed, superactive, mesic Mollic Hapludalfs.

567C2—Elkhart silt loam, 5 to 10 percent slopes, eroded

Setting

Landform: Ground moraines

Position on landform: Shoulders and backslopes

Map Unit Composition

Elkhart and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a lighter colored surface layer
- Soils that have carbonates at a depth of more than 40 inches
- Soils that have more clay in the surface layer
- Soils that have more sand in the lower part of the subsoil and in the underlying material

Dissimilar soils:

- The somewhat poorly drained Ipava soils on summits

Properties and Qualities of the Elkhart Soil

Parent material: Loess

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:

Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.2 inches to a depth of 60 inches

Content of organic matter in the surface layer: 2.0 to 3.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: 4 to 6 feet below the surface

Flooding: None

Accelerated erosion: The surface layer has been thinned by erosion.

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Medium

Susceptibility to water erosion: Moderate

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Elkhart—3e

Prime farmland status: Elkhart—not prime farmland

Hydric soil status: Elkhart—not hydric

567D2—Elkhart silt loam, 10 to 18 percent slopes, eroded

Setting

Landform: Ground moraines

Position on landform: Backslopes

Map Unit Composition

Elkhart and similar soils: 100 percent

Minor Components

Similar soils:

- Soils that have a lighter colored surface layer
- Soils that have carbonates at a depth of more than 40 inches
- Soils that have more clay in the surface layer
- Soils that have more sand in the lower part of the subsoil and in the underlying material
- Soils that have carbonates within a depth of 20 inches

Properties and Qualities of the Elkhart Soil

Parent material: Loess

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:

Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.2 inches to a depth of 60 inches

Content of organic matter in the surface layer: 2.0 to 3.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: 4 to 6 feet below the surface

Flooding: None

Accelerated erosion: The surface layer has been thinned by erosion.

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Medium

Susceptibility to water erosion: High

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Elkhart—3e

Prime farmland status: Elkhart—not prime farmland

Hydric soil status: Elkhart—not hydric

Fayette Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Typic Hapludalfs

Typical Pedon

Fayette silt loam, 10 to 18 percent slopes, eroded (fig. 6), at an elevation of 685 feet; Warren County, Illinois; 2,100 feet north and 1,700 feet west of the southeast corner of sec. 31, T. 12 N., R. 3 W.; USGS Rozetta topographic quadrangle; lat. 40 degrees 59 minutes 13 seconds N. and long. 90 degrees 46 minutes 18 seconds W., NAD 27:

Ap—0 to 5 inches; mixed dark grayish brown (10YR 4/2) and yellowish brown (10YR 5/4) silt loam, light brownish gray (10YR 6/2) dry; moderate medium granular structure; friable; common fine roots throughout; moderately acid; clear smooth boundary.

EB—5 to 9 inches; mixed brown (10YR 5/3) and yellowish brown (10YR 5/4) silt loam; weak medium platy structure parting to moderate fine subangular blocky; friable; common fine roots between peds; few faint dark yellowish brown (10YR 4/4) clay films on faces of peds; moderately acid; clear smooth boundary.

Bt1—9 to 13 inches; dark yellowish brown (10YR 4/4) silt loam; moderate fine and medium subangular blocky structure; friable; few fine roots between peds; common faint brown (10YR 4/3) clay films on faces of peds; moderately acid; clear smooth boundary.

Bt2—13 to 27 inches; yellowish brown (10YR 5/4) silty clay loam; moderate medium subangular blocky structure; friable; few fine roots between peds; common faint dark yellowish brown (10YR 4/4) clay films on faces of peds; moderately acid; gradual smooth boundary.

Bt3—27 to 38 inches; yellowish brown (10YR 5/4) silty clay loam; weak coarse prismatic structure parting to moderate medium subangular blocky; friable; common faint dark yellowish brown (10YR 4/4) clay films on faces of peds; common distinct light gray (10YR 7/2 dry) clay depletions on faces of peds; few distinct dark brown (7.5YR 3/2) accumulations of iron and manganese on faces of peds; moderately acid; gradual wavy boundary.

BC—38 to 55 inches; yellowish brown (10YR 5/4) silt loam; moderate medium and coarse subangular blocky structure; friable; common faint dark yellowish brown (10YR 4/4) clay films on faces of peds; common distinct light gray (10YR 7/2 dry) clay depletions on faces of peds; few distinct dark brown (7.5YR 3/2) accumulations of iron and manganese on faces of peds; moderately acid; clear wavy boundary.

C—55 to 60 inches; yellowish brown (10YR 5/4) silt loam; massive; friable; few distinct dark brown




Figure 6.—Typical profile of Fayette silt loam.

(7.5YR 3/2) concretions of iron and manganese in the matrix; moderately acid.

Range in Characteristics

Depth to base of diagnostic horizon: 36 to 70 inches

Ap or A horizon(s):

Hue—10YR

Value—2 to 4

Chroma—1 to 3

Texture—silt loam or silty clay loam

E horizon(s), where present:

Hue—10YR

Value—4 or 5

Chroma—1 to 4

Texture—silt loam

Bt horizon(s):

Hue—10YR

Value—4 or 5

Chroma—3 to 6

Texture—silty clay loam or silt loam

C horizon(s):

Hue—10YR

Value—4 or 5

Chroma—4 to 6

Texture—silt loam or silty clay loam

280C2—Fayette silt loam, 5 to 10 percent slopes, eroded

Setting

Landform: Ground moraines

Position on landform: Shoulders and backslopes

Map Unit Composition

Fayette and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a seasonal high water table at a depth of 4 to 6 feet
- Soils that have more clay in the surface layer
- Soils that have carbonates within a depth of 40 inches
- Soils that have more sand in the lower part of the subsoil and in the underlying material

Dissimilar soils:

- The somewhat poorly drained Keomah soils on summits
- The well drained Thebes soils on backslopes and summits

Properties and Qualities of the Fayette Soil

Parent material: Loess

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:

Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.4 inches to a depth of 60 inches

Content of organic matter in the surface layer: 1.0 to 2.0 percent

Shrink-swell potential: Moderate
Flooding: None
Accelerated erosion: The surface layer has been thinned by erosion.
Potential for frost action: High
Corrosivity: Moderate for steel and concrete
Surface runoff class: Medium
Susceptibility to water erosion: High
Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Fayette—3e
Prime farmland status: Fayette—not prime farmland
Hydric soil status: Fayette—not hydric

280D2—Fayette silt loam, 10 to 18 percent slopes, eroded

Setting

Landform: Ground moraines
Position on landform: Backslopes

Map Unit Composition

Fayette and similar soils: 90 percent
 Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a seasonal high water table at a depth of 4 to 6 feet
- Soils that have more sand in the lower part of the subsoil and in the underlying material
- Soils that have more clay in the surface layer
- Soils that have carbonates within a depth of 40 inches

Dissimilar soils:

- The somewhat poorly drained Keomah soils on summits
- The well drained Thebes soils on backslopes and summits

Properties and Qualities of the Fayette Soil

Parent material: Loess
Drainage class: Well drained
Slowest permeability within a depth of 40 inches: Moderate
Permeability below a depth of 60 inches: Moderate
Depth to restrictive feature: More than 80 inches
Available water capacity: About 11.4 inches to a depth of 60 inches
Content of organic matter in the surface layer: 1.0 to 2.0 percent
Shrink-swell potential: Moderate

Flooding: None
Accelerated erosion: The surface layer has been thinned by erosion.
Potential for frost action: High
Corrosivity: Moderate for steel and concrete
Surface runoff class: Medium
Susceptibility to water erosion: High
Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Fayette—3e
Prime farmland status: Fayette—not prime farmland
Hydric soil status: Fayette—not hydric

280D3—Fayette silty clay loam, 10 to 18 percent slopes, severely eroded

Setting

Landform: Ground moraines
Position on landform: Backslopes

Map Unit Composition

Fayette and similar soils: 90 percent
 Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a seasonal high water table at a depth of 4 to 6 feet
- Soils that have more sand in the lower part of the subsoil and in the underlying material
- Soils that have less clay in the surface layer
- Soils that have carbonates within a depth of 40 inches

Dissimilar soils:

- The well drained Thebes soils on backslopes
- The moderately well drained Elco soils on backslopes
- Somewhat poorly drained soils that are on backslopes and have more clay in the upper part of the subsoil

Properties and Qualities of the Fayette Soil

Parent material: Loess
Drainage class: Well drained
Slowest permeability within a depth of 40 inches: Moderate
Permeability below a depth of 60 inches: Moderate
Depth to restrictive feature: More than 80 inches
Available water capacity: About 11.4 inches to a depth of 60 inches
Content of organic matter in the surface layer: 0.5 to 1.0 percent

Shrink-swell potential: Moderate

Flooding: None

Accelerated erosion: The surface layer is mostly subsoil material.

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Medium

Susceptibility to water erosion: High

Susceptibility to wind erosion: Very low

Interpretive Groups

Land capability classification: Fayette—4e

Prime farmland status: Fayette—not prime farmland

Hydric soil status: Fayette—not hydric

Harrison Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Oxyaquic Argiudolls

Typical Pedon (OSD)

Harrison silt loam, 2 to 5 percent slopes, at an elevation of 665 feet; Christian County, Illinois; 228 feet north and 1,350 feet west of the southeast corner of sec. 24, T. 12 N., R. 2 W.; USGS Clarksdale topographic quadrangle; lat. 39 degrees 27 minutes 59 seconds N. and long. 89 degrees 15 minutes 17 seconds W., NAD 27:

Ap—0 to 10 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; weak fine granular structure; friable; common very fine and few fine roots; slightly acid; abrupt smooth boundary.

BA—10 to 14 inches; brown (10YR 4/3) silt loam; weak very fine and fine subangular blocky structure; friable; few very fine roots; many distinct very dark grayish brown (10YR 3/2) organic coatings on faces of peds; moderately acid; clear smooth boundary.

Bt1—14 to 20 inches; brown (10YR 4/3) silt loam; moderate fine subangular blocky structure; friable; few very fine roots; common distinct very dark grayish brown (10YR 3/2) and few distinct very dark gray (10YR 3/1) organo-clay films on faces of peds; few fine faint dark yellowish brown (10YR 4/4) accumulations of iron along micropores; few fine accumulations of manganese in the matrix; moderately acid; clear smooth boundary.

Bt2—20 to 27 inches; brown (10YR 4/3) silty clay loam; weak fine prismatic structure parting to moderate medium subangular blocky; firm; few very fine roots; common distinct very dark grayish brown (10YR 3/2) organo-clay films on faces of

peds; few fine faint brown (7.5YR 4/4 and 10YR 5/3) and dark yellowish brown (10YR 4/4) accumulations of iron along micropores; few fine accumulations of manganese in the matrix; moderately acid; clear smooth boundary.

Bt3—27 to 35 inches; brown (10YR 5/3) silty clay loam; moderate medium subangular blocky structure; firm; few very fine roots; few distinct dark grayish brown (10YR 4/2) clay films on faces of peds; many fine faint dark yellowish brown (10YR 4/4) and few fine faint brown (7.5YR 4/4) accumulations of iron along micropores; few fine accumulations of manganese in the matrix; moderately acid; clear smooth boundary.

Bt4—35 to 45 inches; yellowish brown (10YR 5/4) silt loam; weak medium and coarse subangular blocky structure; firm; few very fine roots; few distinct dark grayish brown (10YR 4/2) clay films on faces of peds; many fine distinct grayish brown (10YR 5/2) iron depletions along micropores; common fine faint dark yellowish brown (10YR 4/4) and few fine faint brown (7.5YR 4/4) accumulations of iron in the matrix; few fine accumulations of manganese in the matrix; moderately acid; clear smooth boundary.

2Btg—45 to 65 inches; grayish brown (10YR 5/2) silty clay loam; weak medium and coarse subangular blocky structure; firm; few distinct gray (10YR 5/1) clay films on faces of peds; few fine faint brown (10YR 5/3), common fine and medium distinct dark yellowish brown (10YR 4/4), and few fine distinct brown (7.5YR 4/4) accumulations of iron in the matrix; few fine accumulations of manganese in the matrix; about 15 percent sand; about 1 percent gravel; slightly acid; abrupt smooth boundary.

3Btgb—65 to 80 inches; grayish brown (2.5Y 5/2) clay loam; moderate coarse subangular blocky structure; firm; common distinct olive gray (5Y 4/2) clay films on faces of peds; common medium distinct yellowish brown (10YR 5/4) accumulations of iron in the matrix; few fine accumulations of manganese in the matrix; about 5 percent gravel; neutral.

Range in Characteristics

Thickness of the mollic epipedon: 10 to 19 inches

Thickness of the loess: 40 to 60 inches

Depth to base of diagnostic horizon: More than 45 inches

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 to 3

Texture—silt loam

AB or BA horizon(s):

Hue—10YR

Value—3 or 4

Chroma—2 to 4

Texture—silt loam or silty clay loam

Bt horizon(s):

Hue—10YR

Value—4 to 6

Chroma—2 to 6

Texture—silty clay loam or silt loam

2Btg or 2BCg horizon(s):

Hue—10YR, 7.5YR, or 2.5Y

Value—4 to 6

Chroma—2 to 6

Texture—silty clay loam, silt loam, loam, or clay loam

3Btgb horizon(s):

Hue—10YR, 2.5Y, 5Y, or neutral

Value—4 to 6

Chroma—0 to 3

Texture—clay loam, clay, silty clay, or silty clay loam

Taxadjunct Feature

Harrison silt loam, 5 to 10 percent slopes, eroded, has a dark surface soil that is thinner than is definitive for the series. This difference, however, does not significantly affect the use and management of the soil. The soil is classified as a fine-silty, mixed, superactive, mesic Oxyaquic Hapludalf.

127B—Harrison silt loam, 2 to 5 percent slopes

Setting

Landform: Ground moraines

Position on landform: Summits and backslopes

Map Unit Composition

Harrison and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a thinner dark surface layer
- Soils that have less sand in the lower part of the subsoil and in the underlying material
- Soils that have a lighter colored surface layer
- Soils that have a seasonal high water table at a depth of more than 3.5 feet

Dissimilar soils:

- The poorly drained Cowden and Virden soils on toeslopes

Properties and Qualities of the Harrison Soil

Parent material: Loess over pedisediment

Drainage class: Moderately well drained

Slowest permeability within a depth of 40 inches:

Moderate

Permeability below a depth of 60 inches: Slow

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.8 inches to a depth of 60 inches

Content of organic matter in the surface layer: 3.0 to 4.0 percent

Shrink-swell potential: High

Perched seasonal high water table: 2.0 to 3.5 feet below the surface

Flooding: None

Potential for frost action: High

Corrosivity: High for steel and moderate for concrete

Surface runoff class: Low

Susceptibility to water erosion: Moderate

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Harrison—2e

Prime farmland status: Harrison—prime farmland in all areas

Hydric soil status: Harrison—not hydric

127C2—Harrison silt loam, 5 to 10 percent slopes, eroded

Setting

Landform: Ground moraines

Position on landform: Backslopes

Map Unit Composition

Harrison and similar soils: 100 percent

Minor Components

Similar soils:

- Soils that have less sand in the lower part of the subsoil and in the underlying material
- Soils that have a thicker dark surface soil
- Soils that have a lighter colored surface layer
- Soils that have a seasonal high water table at a depth of more than 3.5 feet

Properties and Qualities of the Harrison Soil

Parent material: Loess over pedisediment

Drainage class: Moderately well drained

Slowest permeability within a depth of 40 inches:

Moderate

Permeability below a depth of 60 inches: Slow

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.8 inches to a depth of 60 inches

Content of organic matter in the surface layer: 2.0 to 3.0 percent

Shrink-swell potential: High

Perched seasonal high water table: 2.0 to 3.5 feet below the surface

Flooding: None

Accelerated erosion: The surface layer has been thinned by erosion.

Potential for frost action: High

Corrosivity: High for steel and moderate for concrete

Surface runoff class: Medium

Susceptibility to water erosion: Moderate

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Harrison—3e

Prime farmland status: Harrison—not prime farmland

Hydric soil status: Harrison—not hydric

Hartsburg Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Typic Endoaquolls

Typical Pedon (OSD)

Hartsburg silty clay loam, 0 to 2 percent slopes, at an elevation of 562 feet; Logan County, Illinois; 660 feet west and 40 feet north of the southeast corner of sec. 23, T. 21 N., R. 4 W.; USGS New Holland topographic quadrangle; lat. 40 degrees 14 minutes 58 seconds N. and long. 89 degrees 31 minutes 28 seconds W., NAD 27:

Ap—0 to 7 inches; black (10YR 2/1) silty clay loam, dark gray (10YR 4/1) dry; moderate medium granular structure; friable; slightly acid; abrupt smooth boundary.

A1—7 to 12 inches; black (10YR 2/1) silty clay loam, dark gray (10YR 4/1) dry; moderate medium granular structure; friable; few very fine roots; slightly acid; clear smooth boundary.

A2—12 to 17 inches; very dark gray (10YR 3/1) silty clay loam, gray (10YR 5/1) dry; moderate medium granular structure; firm; few very fine roots; few fine rounded black (7.5YR 2.5/1) weakly cemented concretions of iron and manganese with diffuse boundaries along root channels and pores; few

fine faint dark grayish brown (2.5Y 4/2) iron depletions in the matrix; neutral; clear smooth boundary.

Bg—17 to 21 inches; dark grayish brown (2.5Y 4/2) silty clay loam; weak fine and medium subangular blocky structure; firm; few very fine roots; common distinct very dark gray (10YR 3/1) organic coatings on faces of peds; common very dark gray (10YR 3/1) krotovinas; few fine rounded black (7.5YR 2.5/1) weakly cemented concretions of iron and manganese with diffuse boundaries lining root channels and pores; common fine prominent yellowish brown (10YR 5/6) accumulations of iron in the matrix; neutral; clear smooth boundary.

Bkg—21 to 30 inches; gray (5Y 5/1) silty clay loam; moderate medium subangular blocky structure; firm; few very fine roots; common distinct dark grayish brown (10YR 4/2) and grayish brown (2.5Y 5/2) pressure faces on peds; common very dark gray (10YR 3/1) krotovinas; few fine rounded black (7.5YR 2.5/1) weakly cemented concretions of iron and manganese with diffuse boundaries lining root channels and pores; few fine and medium rounded white (10YR 8/1) weakly cemented calcium carbonate concretions throughout; common medium prominent yellowish brown (10YR 5/8) and strong brown (7.5YR 5/8) accumulations of iron in the matrix; slightly effervescent; slightly alkaline; abrupt wavy boundary.

BCkg—30 to 34 inches; light brownish gray (2.5Y 6/2) silty clay loam; weak coarse subangular blocky structure; firm; many distinct gray (N 5/0) and grayish brown (2.5Y 5/2) linings in pores and root channels; common very dark gray (10YR 3/1) krotovinas; few fine rounded black (7.5YR 2.5/1) weakly cemented concretions of iron and manganese with diffuse boundaries lining pores; many medium and coarse rounded white (10YR 8/1) weakly cemented calcium carbonate concretions throughout; many medium prominent yellowish brown (10YR 5/8) accumulations of iron in the matrix; violently effervescent among concretions, slightly effervescent in the matrix; slightly alkaline; clear wavy boundary.

Cg—34 to 60 inches; light brownish gray (2.5Y 6/2) silt loam; massive; friable; common very dark gray (10YR 3/1) krotovinas; few medium rounded white (10YR 8/1) weakly cemented calcium carbonate concretions throughout; many medium prominent strong brown (7.5YR 5/8) accumulations of iron with diffuse boundaries lining pores; strongly effervescent; moderately alkaline.

Range in Characteristics

Thickness of the mollic epipedon: 10 to 20 inches
Depth to carbonates: 15 to 35 inches
Depth to base of diagnostic horizon: 24 to 50 inches

Ap, A, or AB horizon(s):

Hue—10YR or neutral
 Value—2 or 3
 Chroma—0 to 2
 Texture—silty clay loam

BA, Bg, Bkg, Btg, BCK, or BCg horizon(s):

Hue—10YR, 2.5Y, or 5Y
 Value—3 to 6
 Chroma—1 or 2
 Texture—silty clay loam or silt loam

Cg horizon(s):

Hue—10YR, 2.5Y, or 5Y
 Value—5 or 6
 Chroma—1 or 2
 Texture—silt loam

244A—Hartsburg silty clay loam, 0 to 2 percent slopes

Setting

Landform: Ground moraines
Position on landform: Toeslopes

Map Unit Composition

Hartsburg and similar soils: 90 percent
 Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have more clay in the subsoil
- Soils that have carbonates at a depth of more than 35 inches
- Soils that have carbonates within a depth of 15 inches

Dissimilar soils:

- The somewhat poorly drained Ipava soils on summits and footslopes
- The moderately well drained Buckhart soils on backslopes and summits

Properties and Qualities of the Hartsburg Soil

Parent material: Loess

Drainage class: Poorly drained

Slowest permeability within a depth of 40 inches:
 Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches
Available water capacity: About 12.6 inches to a depth of 60 inches

Content of organic matter in the surface layer: 3.0 to 5.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: At the surface to 1 foot below the surface

Ponding: At the surface to 0.5 foot above the surface

Flooding: None

Potential for frost action: High

Corrosivity: High for steel and low for concrete

Surface runoff class: Negligible

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Very low

Interpretive Groups

Land capability classification: Hartsburg—2w

Prime farmland status: Hartsburg—prime farmland where drained

Hydric soil status: Hartsburg—hydric

Herrick Series

Taxonomic classification: Fine, smectitic, mesic Aquic Argiudolls

Typical Pedon (OSD)

Herrick silt loam, 0 to 2 percent slopes, at an elevation of 635 feet; Christian County, Illinois; 1,260 feet south and 60 feet west of the northeast corner of sec. 1, T. 11 N., R. 3 W.; USGS Clarksdale Topographic quadrangle; lat. 39 degrees 26 minutes 0 seconds N. and long. 89 degrees 22 minutes 0 seconds W., NAD 27:

Ap1—0 to 7 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; moderate very fine granular structure; very friable; common fine and very fine roots; common wormcasts and worm channels; common very dark gray (10YR 3/1) organic coatings on faces of peds and lining channels; light brownish gray (10YR 6/2 dry) silt coatings and common very fine concretions of iron and manganese on the soil surface; slightly acid; clear wavy boundary.

Ap2—7 to 11 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; weak coarse subangular blocky structure parting to weak fine granular; very friable; few fine and medium roots; common wormcasts and worm channels; common distinct very dark gray (10YR 3/1) organic coatings lining channels; few very fine

- iron and manganese accumulations; slightly acid; abrupt smooth boundary.
- E—11 to 15 inches; very dark grayish brown (10YR 3/2) silt loam, gray (10YR 5/1) dry; discontinuous weak thick platy structure parting to weak medium granular; friable; few fine roots; few medium worm channels; many distinct grayish brown (10YR 5/2) clay depletions on the surface of peds, distinct light gray (10YR 7/1) dry; few fine concretions of iron and manganese and common soft yellowish brown iron accumulations that streak with tools on cut faces; moderately acid; clear smooth boundary.
- Btg—15 to 19 inches; dark grayish brown (10YR 4/2) silty clay loam; weak very fine and fine prismatic structure parting to moderate fine and medium subangular blocky; friable; few fine roots; few distinct very dark grayish brown (10YR 3/2) organo-clay films on horizontal and vertical faces of peds; few very fine distinct yellowish brown (10YR 5/4) accumulations of iron; many distinct light gray (10YR 7/1) dry clay depletions on faces of peds; few fine iron and manganese concretions and stains; moderately acid; clear smooth boundary.
- Bt1—19 to 25 inches; brown (10YR 5/3) silty clay loam; moderate medium prismatic structure parting to moderate medium subangular and angular blocky; firm; few fine and very fine roots, dominantly between peds; many distinct very dark grayish brown (10YR 3/2) and common prominent very dark gray (10YR 3/1) organo-clay films and brown (10YR 4/3) clay films on vertical and horizontal faces of peds; common fine distinct yellowish brown (10YR 5/6) and common fine faint brown (7.5YR 4/4) accumulations of iron and few fine faint grayish brown (10YR 5/2) iron depletions; few fine iron and manganese concretions; moderately acid; gradual smooth boundary.
- Bt2—25 to 35 inches; yellowish brown (10YR 5/6) silty clay loam; moderate medium and coarse prismatic structure parting to weak coarse subangular blocky; very firm; few fine and very fine roots, dominantly in the cracks between peds; many distinct very dark gray (10YR 3/1) organo-clay films on vertical and horizontal faces of peds; common medium distinct brown (7.5YR 4/4) accumulations of iron and few fine prominent grayish brown (10YR 5/2) iron depletions; few iron and manganese concretions; moderately acid; gradual smooth boundary.
- Bt3—35 to 47 inches; yellowish brown (10YR 5/6) silty clay loam; moderate coarse prismatic structure parting to weak coarse subangular blocky; firm; few very fine roots; few very fine continuous vertical tubular pores; many distinct very dark grayish brown (10YR 3/2) organo-clay films on vertical faces of peds and lining pores and common distinct clay films on horizontal faces of peds; common fine prominent light brownish gray (2.5Y 5/2) iron depletions; slightly acid; gradual wavy boundary.
- Bt4—47 to 58 inches; yellowish brown (10YR 5/4) silty clay loam; weak very coarse prismatic structure; firm; few very fine roots; many very fine continuous vertical tubular pores; few distinct very dark grayish brown (10YR 3/2) organo-clay films on vertical faces of peds and lining channels; many medium distinct yellowish brown (10YR 5/6) accumulations of iron and common medium distinct light brownish gray (2.5Y 6/2) iron depletions; slightly acid; gradual smooth boundary.
- C—58 to 62 inches; mottled yellowish brown (10YR 5/6) and light brownish gray (2.5Y 6/2) silt loam; massive; friable; few very fine and fine continuous vertical tubular pores; very dark grayish brown clay lining pores; few iron and manganese accumulations and stains; an increase in the content of coarse silt and very fine sand; neutral.

Range in Characteristics

Thickness of the mollic epipedon: 10 to 21 inches
Depth to base of diagnostic horizon: 45 to 60 inches

Ap or A horizon(s):

Hue—10YR
 Value—2 or 3
 Chroma—1 or 2
 Texture—silt loam

E horizon(s):

Hue—10YR
 Value—3 or 4
 Chroma—1 or 2
 Texture—silt loam

Btg or Bt horizon(s):

Hue—10YR, 2.5Y, or 5Y
 Value—4 to 6
 Chroma—2 to 6
 Texture—silty clay loam, silty clay, or silt loam

C or Cg horizon(s):

Hue—10YR, 2.5Y, or 5Y
 Value—4 to 6

Chroma—2 to 6
Texture—silt loam or silty clay loam

46A—Herrick silt loam, 0 to 2 percent slopes

Setting

Landform: Ground moraines
Position on landform: Summits

Map Unit Composition

Herrick and similar soils: 90 percent
Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a seasonal high water table at a depth of 2.0 to 3.5 feet
- Soils that have less clay in the subsoil
- Soils that do not have a light colored subsurface layer

Dissimilar soils:

- The poorly drained Cowden and Virden soils on toeslopes

Properties and Qualities of the Herrick Soil

Parent material: Loess

Drainage class: Somewhat poorly drained

Slowest permeability within a depth of 40 inches:
Moderately slow

Permeability below a depth of 60 inches: Moderately slow

Depth to restrictive feature: More than 80 inches

Available water capacity: About 10.8 inches to a depth of 60 inches

Content of organic matter in the surface layer: 3.0 to 4.0 percent

Shrink-swell potential: High

Apparent seasonal high water table: 1 to 2 feet below the surface

Flooding: None

Potential for frost action: High

Corrosivity: High for steel and concrete

Surface runoff class: Medium

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Herrick—2w

Prime farmland status: Herrick—prime farmland in all areas

Hydric soil status: Herrick—not hydric

Hickory Series

Taxonomic classification: Fine-loamy, mixed, active, mesic Typic Hapludalfs

Typical Pedon

Hickory silt loam, 35 to 60 percent slopes, at an elevation of 565 feet; Cass County, Illinois; 1,935 feet north and 2,130 feet west of the southeast corner of sec. 27, T. 18 N., R. 9 W.; USGS Ashland, Illinois, topographic quadrangle; lat. 39 degrees 58 minutes 47.5 seconds N. and long. 90 degrees 5 minutes 38 seconds W., NAD 27:

A1—0 to 1 inch; very dark grayish brown (10YR 3/2) silt loam, light brownish gray (10YR 6/2) dry; moderate medium granular structure; friable; many very fine roots; slightly acid; abrupt smooth boundary.

A2—1 to 4 inches; 90 percent dark grayish brown (10YR 4/2) and 10 percent brown (10YR 4/3) silt loam, pale brown (10YR 6/3) dry; weak fine subangular blocky and granular structure; friable; many very fine roots; moderately acid; abrupt smooth boundary.

E—4 to 8 inches; brown (10YR 5/3) loam, light gray (10YR 7/2) dry; moderate thin platy structure; friable; few very fine and fine roots; few distinct very dark grayish brown (10YR 3/2) organic coatings in root channels and/or pores; common fine distinct very pale brown (10YR 8/2) clay depletions between peds; 3 percent gravel; strongly acid; abrupt smooth boundary.

BE—8 to 12 inches; yellowish brown (10YR 5/4) loam, light gray (10YR 7/2) dry; moderate very fine and fine subangular blocky structure; friable; few very fine roots; very few faint brown (10YR 5/3) and very few distinct dark grayish brown (10YR 4/2) organic coatings in root channels and/or pores; common fine distinct very pale brown (10YR 8/2) clay depletions between peds; 3 percent gravel; strongly acid; clear smooth boundary.

Bt1—12 to 22 inches; yellowish brown (10YR 5/4) clay loam; moderate fine and medium subangular blocky structure; firm; few very fine roots; common faint dark yellowish brown (10YR 4/4) clay films and common distinct very pale brown (10YR 7/3) silt coatings on faces of peds; 5 percent gravel; very strongly acid; clear smooth boundary.

Bt2—22 to 29 inches; yellowish brown (10YR 5/4) clay loam; moderate fine and medium subangular blocky structure; firm; few very fine roots; many faint dark yellowish brown (10YR 4/4) clay films and few distinct very pale brown (10YR 7/3) silt

coatings on faces of peds; 5 percent gravel; strongly acid; clear smooth boundary.

Bt3—29 to 40 inches; yellowish brown (10YR 5/4) clay loam; moderate medium prismatic and subangular blocky structure; firm; few very fine roots; many distinct brown (7.5YR 4/4) clay films and very few distinct very pale brown (10YR 7/3) silt coatings on faces of peds; 5 percent gravel; moderately acid; clear smooth boundary.

Bt4—40 to 53 inches; yellowish brown (10YR 5/6) clay loam; weak medium prismatic and weak medium and coarse subangular blocky structure; firm; few very fine roots; many distinct brown (7.5YR 4/4) clay films on faces of peds; few prominent fine black (10YR 2/1) accumulations of iron and manganese throughout; 5 percent gravel; moderately acid; gradual smooth boundary.

BCt—53 to 58 inches; yellowish brown (10YR 5/6) loam; weak medium prismatic and weak medium and coarse subangular blocky structure; firm; few very fine roots; common distinct brown (7.5YR 4/4) clay films on faces of peds; few fine prominent black (10YR 2/1) accumulations of iron and manganese and common distinct brown (10YR 5/3) iron depletions throughout; 5 percent gravel; neutral; gradual smooth boundary.

C—58 to 63 inches; yellowish brown (10YR 5/6) loam; massive; firm; very few distinct brown (7.5YR 4/4) clay films in root channels and/or pores; few prominent fine black (10YR 2/1) accumulations of iron and manganese and many fine prominent light brownish gray (2.5Y 6/2) iron depletions throughout; 3 percent gravel; slightly alkaline.

Range in Characteristics

Thickness of the loess: 0 to 20 inches

Depth to carbonates, where present: More than 40 inches

Depth to base of diagnostic horizon: More than 40 inches

Ap or A horizon(s):

Hue—10YR or 7.5YR

Value—2 to 5

Chroma—2 to 4

Texture—silt loam, loam, clay loam, or silty clay loam

Content of rock fragments—0 to 5 percent

E horizon(s):

Hue—10YR

Value—4 to 6

Chroma—2 to 4

Texture—silt loam or loam

Content of rock fragments—0 to 5 percent

Bt horizon(s):

Hue—10YR, 7.5YR, or 2.5Y

Value—4 to 6

Chroma—3 to 6

Texture—clay loam, silty clay loam, loam, or gravelly clay loam

Content of rock fragments—0 to 20 percent

C horizon(s):

Hue—7.5YR, 10YR, or 2.5Y

Value—5 to 7

Chroma—1 to 8

Texture—loam, clay loam, sandy loam, or the gravelly analogs of those textures

Content of rock fragments—2 to 20 percent

8D—Hickory silt loam, 10 to 18 percent slopes

Setting

Landform: Ground moraines

Position on landform: Backslopes

Map Unit Composition

Hickory and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have more sand in the surface layer
- Soils that have more clay in the subsoil
- Soils that have more clay in the surface layer
- Soils that have carbonates within a depth of 40 inches

Dissimilar soils:

- The well drained Rozetta soils on summits
- The well drained Fayette soils on shoulders
- The well drained Marseilles soils on the lower backslopes
- Somewhat poorly drained soils that are on backslopes and have more clay in the upper part of the subsoil

Properties and Qualities of the Hickory Soil

Parent material: Till

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches
Available water capacity: About 10.3 inches to a depth of 60 inches
Content of organic matter in the surface layer: 1.0 to 3.0 percent
Shrink-swell potential: Moderate
Flooding: None
Potential for frost action: Moderate
Corrosivity: Moderate for steel and concrete
Surface runoff class: Medium
Susceptibility to water erosion: High
Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Hickory—3e
Prime farmland status: Hickory—not prime farmland
Hydric soil status: Hickory—not hydric

8D2—Hickory loam, 10 to 18 percent slopes, eroded

Setting

Landform: Ground moraines (fig. 4)
Position on landform: Backslopes

Map Unit Composition

Hickory and similar soils: 90 percent
 Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have more clay in the surface layer
- Soils that have carbonates within a depth of 40 inches
- Soils that have more clay in the subsoil
- Soils that have less sand in the surface layer

Dissimilar soils:

- The well drained Fayette soils on shoulders
- The well drained Marseilles soils on the lower backslopes
- The well drained Rozetta soils on summits
- Somewhat poorly drained soils that are on backslopes and have more clay in the upper part of the subsoil

Properties and Qualities of the Hickory Soil

Parent material: Till
Drainage class: Well drained
Slowest permeability within a depth of 40 inches: Moderate
Permeability below a depth of 60 inches: Moderate
Depth to restrictive feature: More than 80 inches
Available water capacity: About 10.1 inches to a depth of 60 inches

Content of organic matter in the surface layer: 1.0 to 2.0 percent
Shrink-swell potential: Moderate
Flooding: None
Accelerated erosion: The surface layer has been thinned by erosion.
Potential for frost action: Moderate
Corrosivity: Moderate for steel and concrete
Surface runoff class: Medium
Susceptibility to water erosion: High
Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Hickory—3e
Prime farmland status: Hickory—not prime farmland
Hydric soil status: Hickory—not hydric

8D3—Hickory clay loam, 10 to 18 percent slopes, severely eroded

Setting

Landform: Ground moraines
Position on landform: Backslopes

Map Unit Composition

Hickory and similar soils: 90 percent
 Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have less clay in the surface layer
- Soils that have carbonates within a depth of 40 inches
- Soils that have more clay in the subsoil
- Soils that have less sand in the surface layer

Dissimilar soils:

- The well drained Rozetta soils on summits
- The well drained Fayette soils on shoulders
- The well drained Marseilles soils on the lower backslopes
- Somewhat poorly drained soils that are on backslopes and have more clay in the upper part of the subsoil

Properties and Qualities of the Hickory Soil

Parent material: Till
Drainage class: Well drained
Slowest permeability within a depth of 40 inches: Moderate
Permeability below a depth of 60 inches: Moderate
Depth to restrictive feature: More than 80 inches
Available water capacity: About 9.7 inches to a depth of 60 inches

Content of organic matter in the surface layer: 0.5 to 1.0 percent

Shrink-swell potential: Moderate

Flooding: None

Accelerated erosion: The surface layer is mostly subsoil material.

Potential for frost action: Moderate

Corrosivity: Moderate for steel and concrete

Surface runoff class: Medium

Susceptibility to water erosion: High

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Hickory—4e

Prime farmland status: Hickory—not prime farmland

Hydric soil status: Hickory—not hydric

8F—Hickory silt loam, 18 to 35 percent slopes

Setting

Landform: Ground moraines (fig. 3)

Position on landform: Backslopes

Map Unit Composition

Hickory and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have more clay in the surface layer
- Soils that have more clay in the subsoil
- Soils that have more sand in the surface layer
- Soils that have carbonates within a depth of 40 inches

Dissimilar soils:

- The well drained Rozetta soils on summits
- The well drained Fayette soils on shoulders
- The well drained Marseilles soils on the lower backslopes
- Somewhat poorly drained soils that are on backslopes and have more clay in the upper part of the subsoil

Properties and Qualities of the Hickory Soil

Parent material: Till

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 10.3 inches to a depth of 60 inches

Content of organic matter in the surface layer: 1.0 to 2.0 percent

Shrink-swell potential: Moderate

Flooding: None

Potential for frost action: Moderate

Corrosivity: Moderate for steel and concrete

Surface runoff class: High

Susceptibility to water erosion: High

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Hickory—6e

Prime farmland status: Hickory—not prime farmland

Hydric soil status: Hickory—not hydric

Huntsville Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Cumulic Hapludolls

Typical Pedon (OSD)

Huntsville silt loam, 0 to 2 percent slopes, occasionally flooded, at an elevation of 667 feet; Knox County, Illinois; 2,475 feet east and 495 feet south of the northwest corner of sec. 1, T. 12 N., R. 4 E.; USGS La Fayette topographic quadrangle; lat. 41 degrees 03 minutes 37 seconds N. and long. 89 degrees 59 minutes 42 seconds W., NAD 27:

Ap—0 to 10 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; weak fine and medium subangular blocky structure; friable; slightly acid; clear smooth boundary.

A1—10 to 16 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; weak fine granular structure; friable; common faint very dark gray (10YR 3/1) organic coatings on faces of peds; neutral; clear smooth boundary.

A2—16 to 27 inches; dark brown (10YR 3/3) silt loam, brown (10YR 5/3) dry; weak fine granular structure; friable; common distinct very dark gray (10YR 3/1) organic coatings on faces of peds; neutral; clear smooth boundary.

AC—27 to 52 inches; brown (10YR 4/3) silt loam, pale brown (10YR 6/3) dry; weak medium subangular blocky structure; friable; common distinct very dark grayish brown (10YR 3/2) organic coatings on faces of peds; neutral; clear smooth boundary.

C—52 to 60 inches; dark brown (10YR 3/3) silt loam; massive; friable; slightly acid.

Range in Characteristics

Thickness of the mollic epipedon: 24 to 57 inches

Ap or A horizon(s):

Hue—10YR
 Value—2 or 3
 Chroma—1 to 3
 Texture—silt loam

AC horizon:

Hue—10YR
 Value—4 or 5
 Chroma—3 or 4
 Texture—silt loam or loam below a depth of 40 inches

C horizon(s):

Hue—10YR
 Value—3 to 5
 Chroma—3 or 4
 Texture—silt loam with loam or strata of very fine sandy loam to fine sand below a depth of 40 inches in some pedons

3077A—Huntsville silt loam, 0 to 2 percent slopes, frequently flooded

Setting

Landform: Flood plains

Map Unit Composition

Huntsville and similar soils: 90 percent
 Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a thinner dark surface soil
- Soils that have more sand throughout

Dissimilar soils:

- The somewhat poorly drained Lawson and Tice soils on flood plains
- The poorly drained Sawmill soils on flood plains

Properties and Qualities of the Huntsville Soil

Parent material: Alluvium

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:
 Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 13.4 inches to a depth of 60 inches

Content of organic matter in the surface layer: 2.0 to 4.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: 4 to 6 feet below the surface

Potential for frost action: High

Corrosivity: Low for steel and concrete

Surface runoff class: Low

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Huntsville—2w

Prime farmland status: Huntsville—prime farmland where protected from flooding or not frequently flooded during the growing season

Hydric soil status: Huntsville—not hydric

Ipava Series

Taxonomic classification: Fine, smectitic, mesic Aquic Argiudolls

Typical Pedon (OSD)

Ipava silt loam, 0 to 2 percent slopes, at an elevation of 804 feet; Knox County, Illinois; 2,046 feet west and 594 feet north of the southeast corner of sec. 25, T. 13 N., R. 2 E.; USGS Oneida topographic quadrangle; lat. 41 degrees 04 minutes 40 seconds N. and long. 90 degrees 13 minutes 03 seconds W., NAD 27:

Ap—0 to 10 inches; black (10YR 2/1) silt loam, dark gray (10YR 4/1) dry; moderate fine and medium subangular blocky structure; friable; moderately acid; abrupt smooth boundary.

A—10 to 18 inches; very dark grayish brown (10YR 3/2) silty clay loam, grayish brown (10YR 5/2) dry; weak fine and medium subangular blocky structure; friable; common distinct black (10YR 2/1) organic coatings on faces of peds; moderately acid; clear smooth boundary.

BA—18 to 24 inches; brown (10YR 4/3) silty clay loam; moderate fine and medium subangular blocky structure; friable; common distinct very dark gray (10YR 3/1) organic coatings on faces of peds; few fine distinct light brownish gray (2.5Y 6/2) iron depletions in the matrix; few distinct yellowish brown (10YR 5/6) accumulations of iron in the matrix; moderately acid; clear smooth boundary.

Btg1—24 to 31 inches; dark grayish brown (10YR 4/2) silty clay; moderate fine prismatic structure parting to moderate fine subangular blocky; friable; common faint dark gray (10YR 4/1) clay films on faces of peds; few fine distinct light brownish gray (2.5Y 6/2) iron depletions in the matrix; common fine prominent yellowish brown (10YR 5/8)

accumulations of iron in the matrix; slightly acid; clear smooth boundary.

Btg₂—31 to 37 inches; grayish brown (2.5Y 5/2) silty clay loam; moderate medium prismatic structure parting to moderate medium angular blocky; friable; common distinct dark gray (10YR 4/1) clay films on faces of peds; common fine faint light brownish gray (2.5Y 6/2) iron depletions in the matrix; common medium prominent strong brown (7.5YR 5/8) accumulations of iron in the matrix; few fine black (7.5YR 2.5/1) very weakly cemented concretions of iron and manganese throughout; few fine black (7.5YR 2.5/1) iron and manganese stains on faces of peds; slightly alkaline; gradual smooth boundary.

BCg—37 to 50 inches; grayish brown (2.5Y 5/2) silty clay loam; moderate medium prismatic structure parting to moderate medium angular blocky; friable; few distinct very dark grayish brown (10YR 3/2) organo-clay films lining pores and on a few vertical faces of peds; common fine faint light brownish gray (2.5Y 6/2) iron depletions in the matrix; common fine prominent strong brown (7.5YR 5/8) accumulations of iron in the matrix; few fine black (7.5YR 2.5/1) very weakly cemented concretions of iron and manganese throughout; common fine black (7.5YR 2.5/1) iron and manganese stains on faces of peds; slightly alkaline; clear smooth boundary.

Cg—50 to 60 inches; light brownish gray (2.5Y 6/2) silt loam; massive; friable; few faint very dark grayish brown (10YR 3/2) organo-clay films lining pores; common fine prominent yellowish brown (10YR 5/8) accumulations of iron in the matrix; few fine black (7.5YR 2.5/1) very weakly cemented concretions of iron and manganese throughout; few fine black (7.5YR 2.5/1) iron and manganese stains on faces of vertical cracks; moderately alkaline.

Range in Characteristics

Thickness of the mollic epipedon: 10 to 24 inches

Depth to carbonates: More than 40 inches

Depth to base of diagnostic horizon: 35 to 55 inches

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 or 2

Texture—silt loam or silty clay loam

Bt or Btg horizon(s):

Hue—10YR or 2.5Y

Value—3 to 6

Chroma—2 to 4

Texture—silty clay loam, silty clay, or silt loam

Cg or C horizon(s):

Hue—10YR or 2.5Y

Value—5 or 6

Chroma—1 to 4

Texture—silt loam or silty clay loam

43A—Ipava silt loam, 0 to 2 percent slopes

Setting

Landform: Ground moraines (fig. 2)

Position on landform: Summits and footslopes

Map Unit Composition

Ipava and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a seasonal high water table at a depth of 2.0 to 3.5 feet
- Soils that have less clay in the subsoil
- Soils that have more sand in the lower part of the subsoil and in the underlying material
- Soils that have a lighter colored subsurface layer

Dissimilar soils:

- The poorly drained Sable soils in depressions
- The well drained Osco soils on summits
- The poorly drained Virden soils on toeslopes
- The poorly drained Denny soils in depressions

Properties and Qualities of the Ipava Soil

Parent material: Loess

Drainage class: Somewhat poorly drained

Slowest permeability within a depth of 40 inches:

Moderately slow

Permeability below a depth of 60 inches: Moderately slow

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.0 inches to a depth of 60 inches

Content of organic matter in the surface layer: 4.0 to 5.0 percent

Shrink-swell potential: High

Apparent seasonal high water table: 1 to 2 feet below the surface

Flooding: None

Potential for frost action: High

Corrosivity: High for steel and moderate for concrete

Surface runoff class: Medium

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: lpava—1

Prime farmland status: lpava—prime farmland in all areas

Hydric soil status: lpava—not hydric

Kendall Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Aeric Endoaqualfs

Typical Pedon (OSD)

Kendall silt loam, 0 to 2 percent slopes, at an elevation of about 650 feet; Douglas County, Illinois; about 1,160 feet north and 400 feet west of center of sec. 36, T. 15 N., R. 10 E.; USGS Oakland topographic quadrangle; lat. 39 degrees 42 minutes 24 seconds N. and long. 88 degrees 2 minutes 17 seconds W., NAD 27:

Ap—0 to 7 inches; dark grayish brown (10YR 4/2) silt loam, light grayish brown (10YR 6/2) dry; weak medium granular structure; friable; many very fine and fine roots; few fine and medium rounded black (7.5YR 2.5/1) weakly cemented nodules of iron and manganese throughout; neutral; abrupt smooth boundary.

E—7 to 11 inches; grayish brown (10YR 5/2) silt loam; moderate fine and medium granular structure; friable; many very fine and fine roots; common fine and medium rounded black (7.5YR 2.5/1) weakly cemented nodules of iron and manganese throughout; slightly acid; clear smooth boundary.

BE—11 to 14 inches; brown (10YR 5/3) silty clay loam; moderate fine subangular blocky structure; firm; many very fine and fine roots; common fine and medium rounded black (7.5YR 2.5/1) weakly cemented nodules of iron and manganese throughout; slightly acid; clear smooth boundary.

Btg1—14 to 25 inches; grayish brown (10YR 5/2) silty clay loam; moderate fine and medium prismatic structure parting to moderate fine and medium subangular blocky; firm; few very fine and fine roots; common distinct brown (10YR 4/3) clay films on faces of peds; few medium rounded black (7.5YR 2.5/1) weakly cemented nodules of iron and manganese throughout; common fine faint brown (10YR 5/3) accumulations of iron in the matrix; strongly acid; clear smooth boundary.

Btg2—25 to 41 inches; grayish brown (2.5Y 5/2) silty clay loam; moderate medium prismatic structure parting to moderate medium and coarse subangular blocky; firm; few very fine and fine

roots; common distinct dark grayish brown (10YR 4/2) clay films on faces of peds; few medium rounded black (7.5YR 2.5/1) weakly cemented nodules of iron and manganese throughout; common medium prominent yellowish brown (10YR 5/6) accumulations of iron in the matrix; moderately acid; clear smooth boundary.

Btg3—41 to 51 inches; 55 percent yellowish brown (10YR 5/6) and 45 percent gray (5Y 5/1) silty clay loam; weak medium prismatic structure parting to weak coarse subangular blocky; firm; few very fine and fine roots; common distinct gray (10YR 5/1) clay films on faces of peds; few medium rounded black (7.5YR 2.5/1) weakly cemented nodules of iron and manganese throughout; slightly acid; clear smooth boundary.

2Btg4—51 to 58 inches; 40 percent strong brown (7.5YR 5/6), 30 percent yellowish brown (10YR 5/6), and 30 percent gray (5Y 5/1) loam; weak coarse subangular blocky structure; friable; few distinct dark gray (10YR 4/1) clay films on faces of peds; common fine and medium rounded black (7.5YR 2.5/1) weakly cemented nodules of iron and manganese throughout; about 5 percent fine gravel; neutral; clear smooth boundary.

2Cg1—58 to 74 inches; 45 percent yellowish brown (10YR 5/6), 45 percent gray (5Y 5/1), and 10 percent strong brown (7.5YR 5/6), stratified loam, sandy loam, and silt loam; massive; friable; about 5 percent fine gravel; slightly alkaline; abrupt smooth boundary.

2Cg2—74 to 80 inches; 60 percent grayish brown (10YR 5/2), 30 percent gray (10YR 5/1), and 10 percent yellowish brown (10YR 5/6), stratified gravelly loam, gravelly sandy loam, and silt loam; massive; friable; slightly effervescent; slightly alkaline.

Range in Characteristics

Thickness of the loess: 40 to 60 inches

Depth to carbonates: 40 inches or more

Depth to base of diagnostic horizon: 40 to more than 60 inches

Ap or A horizon(s):

Hue—10YR

Value—2 to 5; 2 or 3 where the A horizon is less than 7 inches thick

Chroma—1 to 3

Texture—silt loam

E or Eg horizon(s):

Hue—10YR or 2.5Y

Value—4 to 7

Chroma—2 or 3

Texture—silt loam

BE horizon, where present:

Hue—7.5YR, 10YR, or 2.5Y

Value—4 to 6

Chroma—2 to 6

Texture—silt loam or silty clay loam

Btg or Bt horizon(s):

Hue—7.5YR, 10YR, 2.5Y, or 5Y

Value—4 to 6

Chroma—1 to 8

Texture—silty clay loam

2Btg, 2Bt, 2BCg, or 2BC horizon(s):

Hue—7.5YR, 10YR, 2.5Y, or 5Y

Value—4 to 6

Chroma—1 to 8

Texture—loam, clay loam, silt loam, or sandy loam

2Cg or 2C horizon(s):

Hue—7.5YR, 10YR, 2.5Y, or 5Y

Value—4 to 6

Chroma—1 to 8

Texture—stratified silt loam, loam, sandy loam, clay loam, silty clay loam, sandy clay loam, or the gravelly analogs of those textures

242A—Kendall silt loam, 0 to 2 percent slopes

Setting

Landform: Stream terraces

Position on landform: Summits and footslopes

Map Unit Composition

Kendall and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a darker surface soil
- Soils that have less sand in the lower part of the subsoil and in the underlying material
- Soils that are subject to rare flooding

Dissimilar soils:

- The well drained Camden soils on summits and risers
- The poorly drained Vesser soils on flood plains

Properties and Qualities of the Kendall Soil

Parent material: Loess over outwash

Drainage class: Somewhat poorly drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderate or moderately rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.6 inches to a depth of 60 inches

Content of organic matter in the surface layer: 1.0 to 3.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: 0.5 foot to 2.0 feet below the surface

Flooding: None

Potential for frost action: High

Corrosivity: High for steel and concrete

Surface runoff class: Low

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Kendall—2w

Prime farmland status: Kendall—prime farmland where drained

Hydric soil status: Kendall—not hydric

7242A—Kendall silt loam, 0 to 2 percent slopes, rarely flooded

Setting

Landform: Flood-plain steps

Map Unit Composition

Kendall and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a darker surface soil
- Soils that have less sand in the lower part of the subsoil and in the underlying material

Dissimilar soils:

- The well drained Camden soils on summits and risers
- The poorly drained Vesser soils on flood plains

Properties and Qualities of the Kendall Soil

Parent material: Loess or other silty material

Drainage class: Somewhat poorly drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderate or moderately rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.6 inches to a depth of 60 inches

Content of organic matter in the surface layer: 1.0 to 3.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: 0.5 foot to 2.0 feet below the surface

Potential for frost action: High

Corrosivity: High for steel and moderate for concrete

Surface runoff class: Low

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Kendall—2w

Prime farmland status: Kendall—prime farmland where drained

Hydric soil status: Kendall—not hydric

Keomah Series

Taxonomic classification: Fine, smectitic, mesic Aeric Endoaqualfs

Typical Pedon

Keomah silt loam, 0 to 2 percent slopes, at an elevation of 655 feet; Adams County, Illinois; 2,495 feet south and 300 feet west of the northeast corner of sec. 4, T. 2 N., R. 7 W.; USGS Loraine topographic quadrangle; lat. 40 degrees 11 minutes 22 seconds N. and long. 91 degrees 12 minutes 11 seconds W., NAD 27:

Ap1—0 to 6 inches; dark grayish brown (10YR 4/2) silt loam, light brownish gray (10YR 6/2) dry; weak thick platy structure parting to weak fine subangular blocky; friable; many very fine and fine roots; moderately acid; abrupt smooth boundary.

Ap2—6 to 11 inches; dark grayish brown (10YR 4/2) silt loam, light brownish gray (10YR 6/2) dry; weak medium platy structure parting to weak fine subangular blocky; friable; common very fine and fine roots; few distinct brown (7.5YR 4/4) accumulations of iron throughout; moderately acid; abrupt smooth boundary.

E—11 to 18 inches; grayish brown (10YR 5/2) silt loam, light gray (10YR 7/2) dry; weak medium platy structure parting to weak fine subangular blocky; friable; common fine roots; few faint dark grayish brown (10YR 4/2) organic coatings on faces of peds and in pores; few distinct black (2.5Y 2/1) accumulations of iron and manganese throughout, few prominent strong brown (7.5YR 5/6) accumulations of iron throughout, and few faint light gray (10YR 7/2) clay depletions throughout; slightly acid; clear smooth boundary.

Bt1—18 to 25 inches; brown (10YR 5/3) silty clay loam; weak medium prismatic structure parting to moderate fine subangular blocky; firm; common fine roots; many distinct grayish brown (10YR 5/2) clay films on faces of peds; many prominent strong brown (7.5YR 5/6) accumulations of iron throughout, common distinct black (2.5Y 2/1) accumulations of iron and manganese throughout, and few faint grayish brown (10YR 5/2) iron depletions throughout; strongly acid; clear smooth boundary.

Bt2—25 to 33 inches; brown (10YR 5/3) silty clay loam; weak medium prismatic structure parting to moderate medium subangular blocky; firm; few fine roots; many distinct grayish brown (10YR 5/2) clay films on faces of peds and few faint pressure faces; common distinct black (2.5Y 2/1) accumulations of iron and manganese and many prominent strong brown (7.5YR 5/6) accumulations of iron throughout; strongly acid; clear smooth boundary.

Bt3—33 to 44 inches; brown (10YR 5/3) silty clay loam; weak medium prismatic structure parting to moderate medium subangular blocky; firm; few very fine roots; common distinct grayish brown (10YR 5/2) clay films on faces of peds; many prominent strong brown (7.5YR 5/6) accumulations of iron throughout, common prominent black (2.5Y 2/1) accumulations of iron and manganese throughout, and common faint light brownish gray (10YR 6/2) iron depletions throughout; moderately acid; clear smooth boundary.

Bt4—44 to 51 inches; light brownish gray (10YR 6/2) silty clay loam; weak coarse prismatic structure; firm; few fine roots; few distinct dark grayish brown (10YR 4/2) clay films in root channels and/or pores; few prominent black (2.5Y 2/1) accumulations of iron and manganese and many prominent strong brown (7.5YR 5/6) accumulations of iron throughout; moderately acid; clear smooth boundary.

BC1—51 to 63 inches; light brownish gray (10YR 6/2) silt loam; weak coarse prismatic structure; friable; few very fine roots; common prominent very dark grayish brown (10YR 3/2) organo-clay films in root channels and/or pores; many prominent strong brown (7.5YR 5/6) accumulations of iron and few prominent black (2.5Y 2/1) accumulations of iron and manganese throughout; slightly acid; clear smooth boundary.

BC2—63 to 76 inches; light brownish gray (10YR 6/2) silt loam; weak coarse prismatic structure; friable; common prominent very dark grayish brown

(10YR 3/2) organo-clay films in root channels and/or pores; few prominent black (2.5Y 2/1) accumulations of iron and manganese and many prominent strong brown (7.5YR 5/6) accumulations of iron throughout; slightly acid; clear smooth boundary.

C—76 to 89 inches; yellowish brown (10YR 5/4) silt loam; massive; friable; few distinct strong brown (7.5YR 5/6) accumulations of iron throughout, few prominent black (2.5Y 2/1) accumulations of iron and manganese throughout, and common distinct light brownish gray (10YR 6/2) iron depletions throughout; slightly acid.

Range in Characteristics

Depth to base of diagnostic horizon: 40 to 76 inches

Ap or A horizon(s):

Hue—10YR

Value—3 or 4; 3 where the horizon is less than 3 inches thick

Chroma—1 or 2

Texture—silt loam

E horizon(s):

Hue—10YR

Value—4 or 5

Chroma—1 to 3

Texture—silt loam

Bt horizon(s):

Hue—10YR, 2.5Y, or 5Y

Value—4 to 6

Chroma—2 to 4

Texture—silty clay loam or silty clay

C horizon(s):

Hue—10YR, 2.5Y, or 5Y

Value—4 to 6

Chroma—2 to 6

Texture—silty clay loam or silt loam

17A—Keomah silt loam, 0 to 2 percent slopes

Setting

Landform: Ground moraines (fig. 4)

Position on landform: Summits

Map Unit Composition

Keomah and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a darker surface soil
- Soils that have less clay in the subsoil

Dissimilar soils:

- The poorly drained Denny soils in depressions
- The well drained Fayette soils on shoulders
- The well drained Rozetta soils on summits and shoulders

Properties and Qualities of the Keomah Soil

Parent material: Loess

Drainage class: Somewhat poorly drained

Slowest permeability within a depth of 40 inches: Slow

Permeability below a depth of 60 inches: Moderately slow

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.8 inches to a depth of 60 inches

Content of organic matter in the surface layer: 1.0 to 3.0 percent

Shrink-swell potential: High

Apparent seasonal high water table: 0.5 foot to 2.0 feet below the surface

Flooding: None

Potential for frost action: High

Corrosivity: High for steel and moderate for concrete

Surface runoff class: Medium

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Keomah—2w

Prime farmland status: Keomah—prime farmland where drained

Hydric soil status: Keomah—not hydric

830—Landfills

Setting

This map unit consists of areas where garbage and other refuse and rubble from the demolition of buildings and pavement typically are covered by a layer of compacted earth. Slopes vary widely. Some landfills are active, but some have been abandoned.

Map Unit Composition

Landfills: 85 percent

Dissimilar components: 15 percent

Minor Components

Dissimilar components:

- The well drained, silty Orthents in areas adjacent to the landfills

Interpretive Groups

Land capability classification: Landfills—none assigned

Prime farmland status: Landfills—not prime farmland

Hydric soil status: Landfills—unranked

Lawson Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Aquic Cumulic Hapludolls

Typical Pedon

Lawson silt loam, 0 to 2 percent slopes, frequently flooded, at an elevation of 685 feet; Adams County, Illinois; 1,900 feet east and 265 feet south of the northwest corner of sec. 3, T. 1 S., R. 5 W.; USGS Clayton, Illinois, topographic quadrangle; lat. 40 degrees 1 minute 5 seconds N. and long. 90 degrees 57 minutes 53 seconds W., NAD 27:

Ap—0 to 6 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; moderate fine granular structure; friable; many fine roots; neutral; abrupt smooth boundary.

A1—6 to 14 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; moderate fine granular structure; friable; common fine roots; neutral; clear smooth boundary.

A2—14 to 22 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; weak fine granular structure; friable; common fine roots; common fine faint brown (10YR 4/3) accumulations of iron throughout; neutral; clear smooth boundary.

A3—22 to 33 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; weak fine granular structure; friable; common fine roots; common fine faint brown (10YR 4/3) accumulations of iron throughout; neutral; clear smooth boundary.

C1—33 to 40 inches; 70 percent very dark grayish brown (10YR 3/2) and 20 percent dark brown (10YR 3/3), stratified silt loam; massive; friable; common fine roots; common fine and medium distinct yellowish brown (10YR 5/6) accumulations of iron and common fine and medium faint dark grayish brown (10YR 4/2) iron depletions throughout; slightly acid; clear smooth boundary.

C2—40 to 56 inches; 60 percent very dark grayish

brown (10YR 3/2) and 30 percent dark brown (10YR 3/3), stratified silt loam; massive; friable; few fine roots; common fine and medium distinct yellowish brown (10YR 5/6) accumulations of iron and common medium faint dark grayish brown (10YR 4/2) iron depletions throughout; slightly acid; clear smooth boundary.

C3—56 to 75 inches; 80 percent very dark grayish brown (10YR 3/2) and 10 percent dark brown (10YR 3/3), stratified silt loam; massive; friable; few fine roots; common fine and medium distinct yellowish brown (10YR 5/6) accumulations of iron between peds, common medium prominent strong brown (7.5YR 5/8) accumulations of iron between peds, and many medium faint dark grayish brown (10YR 4/2) iron depletions throughout; slightly acid; clear smooth boundary.

C4—75 to 80 inches; 80 percent dark grayish brown (10YR 4/2) and 10 percent very dark grayish brown (10YR 3/2), stratified silt loam; massive; friable; common medium and coarse prominent yellowish brown (10YR 5/6) and common fine prominent strong brown (7.5YR 5/8) accumulations of iron throughout and common fine faint dark gray (10YR 4/1) iron depletions throughout; neutral.

Range in Characteristics

Thickness of the mollic epipedon: 24 to 36 inches

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 or 2

Texture—silt loam or silty clay loam

C horizon(s):

Hue—10YR or 2.5Y

Value—3 to 6

Chroma—1 to 3

Texture—stratified silt loam or silty clay loam; strata with more sand below a depth of 40 inches in some pedons

3451A—Lawson silt loam, 0 to 2 percent slopes, frequently flooded

Setting

Landform: Flood plains

Map Unit Composition

Lawson and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a thinner dark surface soil
- Soils that have a lighter colored surface soil
- Soils that have more sand throughout
- Soils that have a buried soil at a depth of 20 to 40 inches

Dissimilar soils:

- The well drained Huntsville soils on flood plains
- The poorly drained Sawmill and Zook soils on flood plains

Properties and Qualities of the Lawson Soil

Parent material: Alluvium

Drainage class: Somewhat poorly drained

Slowest permeability within a depth of 40 inches:

Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.1 inches to a depth of 60 inches

Content of organic matter in the surface layer: 2.0 to 4.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: 1 to 2 feet below the surface

Potential for frost action: High

Corrosivity: Moderate for steel and low for concrete

Surface runoff class: Negligible

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Lawson—3w

Prime farmland status: Lawson—prime farmland where protected from flooding or not frequently flooded during the growing season

Hydric soil status: Lawson—not hydric

Marseilles Series

Taxonomic classification: Fine-silty, mixed, active, mesic Typic Hapludalfs

Typical Pedon (OSD)

Marseilles silt loam, 35 to 60 percent slopes (fig. 7), at an elevation of about 669 feet; Bureau County, Illinois; 2,200 feet west and 1,180 feet south of the northeast corner of sec. 14, T. 15 N., R. 8 E.; USGS Wyandot topographic quadrangle; lat. 41 degrees 17 minutes 20 seconds N. and long. 89 degrees 32 minutes 13 seconds W., NAD 27:



Figure 7.—Typical profile of Marseilles silt loam.

A—0 to 4 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; moderate medium granular structure; friable; many very fine and few fine roots; slightly acid; abrupt smooth boundary.

BE—4 to 9 inches; yellowish brown (10YR 5/4) silt loam; moderate medium platy structure parting to weak very fine subangular blocky; friable; many very fine roots; common distinct light gray (10YR 7/2 dry) clay depletions on faces of peds; few fine accumulations of iron and manganese; neutral; clear smooth boundary.

Bt1—9 to 15 inches; yellowish brown (10YR 5/4) silty clay loam; moderate very fine and fine subangular blocky structure; friable; many very fine and few fine roots; common distinct light gray (10YR 7/2 dry) clay depletions and common distinct brown (10YR 4/3) clay films on faces of peds; few fine accumulations of iron and manganese; slightly acid; clear smooth boundary.

Bt2—15 to 23 inches; brown (10YR 5/3) silty clay loam; strong fine and medium subangular blocky structure; friable; many very fine and few fine roots; few distinct light gray (10YR 7/2 dry) clay depletions and many distinct brown (10YR 4/3) clay films on faces of peds; few fine accumulations of iron and manganese; slightly acid; clear smooth boundary.

2Bt3—23 to 36 inches; grayish brown (2.5Y 5/2) silty clay loam; moderate medium prismatic structure parting to moderate medium subangular blocky; firm; common very fine and few medium roots; many prominent brown (10YR 4/3) clay films on faces of peds; few medium prominent yellowish red (5YR 5/8) redoximorphic concentrations; few fine accumulations of iron and manganese; slightly acid; gradual wavy boundary.

2Cr1—36 to 51 inches; olive gray (5Y 5/2), soft shale; firm; common very fine and few fine roots; common prominent brown (10YR 4/3) clay films on shale fragments; few medium prominent yellowish red (5YR 5/8) redoximorphic concentrations; few fine accumulations of iron and manganese; slightly acid; gradual wavy boundary.

2Cr2—51 to 60 inches; olive (5Y 5/3), soft shale; very firm; few very fine roots; few prominent brown (10YR 4/3) clay films on shale fragments; few fine accumulations of iron and manganese; neutral.

Range in Characteristics

Depth to residuum: 0 to 30 inches

Depth to paralithic contact: 20 to 40 inches

Ap or A horizon(s):

Hue—10YR

Value—2 to 5

Chroma—2 or 3

Texture—silt loam or silty clay loam

BE or E horizon(s), where present:

Hue—10YR

Value—4 or 5

Chroma—2 to 4

Texture—silt loam

Bt horizon(s):

Hue—10YR

Value—4 or 5

Chroma—3 to 6

Texture—silty clay loam or silt loam

2Bt horizon(s):

Hue—7.5YR, 10YR, 2.5Y, or 5Y

Value—4 to 6

Chroma—2 to 4

Texture—clay loam, silt loam, silty clay loam, or silty clay

2Cr horizon(s):

Hue—10YR, 2.5Y, 5Y, or neutral

Value—4 to 6

Chroma—0 to 4

549G—Marseilles silt loam, 35 to 60 percent slopes

Setting

Landform: Ground moraines

Position on landform: Backslopes

Map Unit Composition

Marseilles and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have more clay in the upper part of the subsoil
- Soils that are underlain by sandstone bedrock
- Soils that have shale bedrock within a depth of 20 inches

Dissimilar soils:

- The well drained Hickory soils on the upper backslopes

Properties and Qualities of the Marseilles Soil

Parent material: Thin layer of loess and the underlying residuum weathered from shale

Drainage class: Well drained
Slowest permeability within a depth of 40 inches: Very slow
Permeability below a depth of 60 inches: Very slow or slow
Depth to restrictive feature: 36 inches to paralithic bedrock
Available water capacity: About 5.7 inches to a depth of 60 inches
Content of organic matter in the surface layer: 1.0 to 3.0 percent
Shrink-swell potential: High
Flooding: None
Potential for frost action: High
Corrosivity: High for steel and moderate for concrete
Surface runoff class: Very high
Susceptibility to water erosion: High
Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Marseilles—7e
Prime farmland status: Marseilles—not prime farmland
Hydric soil status: Marseilles—not hydric

Middletown Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Typic Hapludalfs

Typical Pedon (OSD)

Middletown silt loam, 2 to 5 percent slopes, at an elevation of 605 feet; Sangamon County, Illinois; 20 feet west and 1,145 feet south of the northeast corner of sec. 26, T. 17 N., R. 6 W.; USGS Athens topographic quadrangle; lat. 39 degrees 53 minutes 57 seconds N. and long. 89 degrees 43 minutes 51 seconds W., NAD 27:

Ap—0 to 9 inches; dark grayish brown (10YR 4/2) silt loam; moderate fine and medium granular structure; friable; common fine roots; neutral; abrupt smooth boundary.
 E—9 to 12 inches; yellowish brown (10YR 5/4) silt loam; weak medium platy structure; friable; common fine roots; common distinct dark grayish brown (10YR 4/2) organic coatings on faces of peds; neutral; clear smooth boundary.
 Bt1—12 to 17 inches; dark yellowish brown (10YR 4/4) silty clay loam; moderate fine subangular blocky structure; firm; common fine and medium roots; common distinct brown (10YR 4/3) clay films on faces of peds; slightly acid; clear smooth boundary.
 Bt2—17 to 35 inches; dark yellowish brown (10YR 4/4)

silty clay loam; moderate medium subangular blocky structure; firm; common distinct brown (10YR 4/3) clay films on faces of peds; few fine rounded black (5YR 2/1) concretions of iron and manganese in the matrix; strongly acid; gradual smooth boundary.

Bt3—35 to 44 inches; dark yellowish brown (10YR 4/4) silty clay loam; moderate coarse subangular blocky structure; friable; few distinct brown (10YR 4/3) clay films on vertical faces of peds; few fine black (5YR 2/1) concretions of iron and manganese in the matrix; moderately acid; clear smooth boundary.

2Bt4—44 to 47 inches; dark yellowish brown (10YR 4/4) clay loam; weak coarse subangular blocky structure; friable; few distinct brown (10YR 4/3) clay films on vertical faces of peds; moderately acid; abrupt smooth boundary.

2BC1—47 to 52 inches; dark yellowish brown (10YR 4/4) loamy fine sand; weak coarse subangular blocky structure; very friable; moderately acid; gradual smooth boundary.

2BC2—52 to 75 inches; stratified yellowish brown (10YR 5/6) and strong brown (7.5YR 4/6) sand and loamy sand; single grain; loose; 2-inch band of brown (7.5YR 4/4) sandy loam starting at a depth of 64 inches; moderately acid; gradual smooth boundary.

2C—75 to 80 inches; strong brown (7.5YR 4/6) sand; single grain; loose; slightly acid.

Range in Characteristics

Thickness of the loess: 40 to 60 inches

Depth to base of diagnostic horizon: 45 to 80 inches

Ap horizon(s):

Hue—10YR

Value—4 or 5

Chroma—2 or 3

Texture—silt loam

E or BE horizon(s), where present:

Hue—10YR

Value—4 or 5

Chroma—2 to 4

Texture—silt loam

Bt horizon(s):

Hue—10YR or 7.5YR

Value—4 or 5

Chroma—3 to 5

Texture—silty clay loam or silt loam

2Bt horizon(s):

Hue—10YR or 7.5YR

Value—4 or 5

Chroma—3 to 5

Texture—clay loam, fine sandy loam, or loam

2BC horizon(s), where present:

Hue—10YR or 7.5YR

Value—4 or 5

Chroma—4 to 6

Texture—loamy fine sand, loamy sand, or fine sand

2C horizon(s):

Hue—10YR or 7.5YR

Value—4 or 5

Chroma—4 to 6

Texture—fine sand, sand, loamy fine sand, or loamy sand

685B—Middletown silt loam, 2 to 5 percent slopes

Setting

Landform: Ground moraines and knolls

Position on landform: Summits and shoulders

Map Unit Composition

Middletown and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have more sand in the surface layer
- Soils that have less sand in the lower part of the subsoil and in the underlying material
- Soils that have more sand in the upper part of the subsoil

Dissimilar soils:

- The well drained Alvin soils on summits and backslopes

Properties and Qualities of the Middletown Soil

Parent material: Loess over sandy eolian material

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderately rapid or rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 10.7 inches to a depth of 60 inches

Content of organic matter in the surface layer: 1.0 to 3.0 percent

Shrink-swell potential: Moderate

Flooding: None

Potential for frost action: High

Corrosivity: High for steel and concrete

Surface runoff class: Low

Susceptibility to water erosion: Moderate

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Middletown—2e

Prime farmland status: Middletown—prime farmland in all areas

Hydric soil status: Middletown—not hydric

685C2—Middletown silt loam, 5 to 10 percent slopes, eroded

Setting

Landform: Knolls and ground moraines

Position on landform: Backslopes

Map Unit Composition

Middletown and similar soils: 100 percent

Minor Components

Similar soils:

- Soils that have more clay in the surface layer
- Soils that have more sand in the upper part of the subsoil
- Soils that have more sand in the surface layer
- Soils that have less sand in the lower part of the subsoil and in the underlying material

Properties and Qualities of the Middletown Soil

Parent material: Loess over sandy eolian material

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderately rapid or rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.3 inches to a depth of 60 inches

Content of organic matter in the surface layer: 1.0 to 2.0 percent

Shrink-swell potential: Moderate

Flooding: None

Accelerated erosion: The surface layer has been thinned by erosion.

Potential for frost action: High

Corrosivity: High for steel and concrete

Surface runoff class: Medium

Susceptibility to water erosion: High

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Middletown—3e

Prime farmland status: Middletown—not prime farmland

Hydric soil status: Middletown—not hydric

Navlys Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Typic Hapludalfs

Typical Pedon (OSD)

Navlys silty clay loam, 5 to 10 percent slopes, severely eroded, at an elevation of 650 feet; Fulton County, Illinois, 1,411 feet south and 255 feet east of the northwest corner of sec. 11, T. 4 N., R. 2 E.; USGS Ipava topographic quadrangle; lat. 40 degrees 19 minutes 36 seconds N. and long. 90 degrees 14 minutes 50 seconds W., NAD 27:

Ap—0 to 6 inches; 70 percent dark grayish brown (10YR 4/2) and 30 percent yellowish brown (10YR 5/4) silty clay loam, pale brown (10YR 6/3) dry; moderate fine subangular blocky structure parting to weak fine granular; friable; common very fine roots; moderately acid; clear smooth boundary.

Bt1—6 to 15 inches; yellowish brown (10YR 5/4) silty clay loam; strong fine and medium subangular blocky structure; firm; common very fine roots; common distinct brown (10YR 4/3) clay films on faces of peds; common fine distinct yellowish brown (10YR 5/6) accumulations of iron with diffuse boundaries in ped interiors; moderately acid; gradual smooth boundary.

Bt2—15 to 22 inches; 90 percent yellowish brown (10YR 5/4) and 10 percent light brownish gray (10YR 6/2) silty clay loam; strong medium prismatic structure; firm; common very fine roots; common distinct brown (10YR 4/3) clay films on faces of peds; many fine distinct yellowish brown (10YR 5/6) accumulations of iron with diffuse boundaries in ped interiors; slightly acid; gradual smooth boundary.

Bt3—22 to 31 inches; yellowish brown (10YR 5/4) and light brownish gray (10YR 6/2) silty clay loam; moderate medium prismatic structure; firm; few very fine roots; few distinct dark grayish brown (10YR 4/2) clay films lining root channels and pores; many fine distinct yellowish brown (10YR 5/6) accumulations of iron with diffuse boundaries and few fine manganese concretions with sharp

boundaries in ped interiors; slightly effervescent; neutral; gradual smooth boundary.

C1—31 to 56 inches; yellowish brown (10YR 5/4) and light brownish gray (10YR 6/2) silt loam; massive; friable; few very fine roots; few distinct dark grayish brown (10YR 4/2) clay films lining root channels and pores; common fine distinct yellowish brown (10YR 5/6) accumulations of iron with diffuse boundaries and few fine manganese concretions with sharp boundaries throughout; slightly effervescent; slightly alkaline; gradual smooth boundary.

C2—56 to 60 inches; light brownish gray (10YR 6/2) silt loam; massive; friable; common fine distinct light yellowish brown (10YR 6/4) accumulations of iron with diffuse boundaries and few fine manganese concretions with sharp boundaries throughout; slightly effervescent; moderately alkaline.

Range in Characteristics

Depth to base of diagnostic horizon: 22 to 40 inches

Depth to carbonates: 22 to 40 inches

Ap or A horizon(s):

Hue—10YR

Value—4 or 5

Chroma—2 to 4

Texture—silty clay loam or silt loam

Bt or BC horizon(s):

Hue—10YR or 7.5YR

Value—4 or 5

Chroma—3 to 6

Texture—silty clay loam or silt loam

C horizon(s):

Hue—10YR or 2.5Y

Value—4 to 6

Chroma—2 to 6

Texture—silt loam

630C2—Navlys silt loam, 5 to 10 percent slopes, eroded

Setting

Landform: Ground moraines (fig. 4)

Position on landform: Shoulders and backslopes

Map Unit Composition

Navlys and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a darker surface layer
- Soils that have carbonates at a depth of more than 40 inches
- Soils that have more clay in the surface layer
- Soils that have more sand in the lower part of the subsoil and in the underlying material
- Soils that have carbonates within a depth of 20 inches

Dissimilar soils:

- The somewhat poorly drained Keomah soils on summits

Properties and Qualities of the Navlys Soil

Parent material: Loess

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.1 inches to a depth of 60 inches

Content of organic matter in the surface layer: 1.0 to 2.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: 4 to 6 feet below the surface

Flooding: None

Accelerated erosion: The surface layer of this soil has less clay and more organic matter than the surface layer in the Typical Pedon.

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Medium

Susceptibility to water erosion: High

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Navlys—3e

Prime farmland status: Navlys—not prime farmland

Hydric soil status: Navlys—not hydric

630D2—Navlys silt loam, 10 to 18 percent slopes, eroded

Setting

Landform: Ground moraines

Position on landform: Backslopes

Map Unit Composition

Navlys and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a darker surface layer
- Soils that have carbonates at a depth of more than 40 inches
- Soils that have more clay in the surface layer
- Soils that have more sand in the lower part of the subsoil and in the underlying material
- Soils that have carbonates within a depth of 20 inches

Dissimilar soils:

- The well drained Rozetta soils on shoulders and summits

Properties and Qualities of the Navlys Soil

Parent material: Loess

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.1 inches to a depth of 60 inches

Content of organic matter in the surface layer: 1.0 to 2.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: 4 to 6 feet below the surface

Flooding: None

Accelerated erosion: The surface layer of this soil has less clay and more organic matter than the surface layer in the Typical Pedon.

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Medium

Susceptibility to water erosion: High

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Navlys—3e

Prime farmland status: Navlys—not prime farmland

Hydric soil status: Navlys—not hydric

630D3—Navlys silty clay loam, 10 to 18 percent slopes, severely eroded

Setting

Landform: Ground moraines

Position on landform: Backslopes

Map Unit Composition

Navlys and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have carbonates at a depth of more than 40 inches
- Soils that have less clay in the surface layer
- Soils that have more sand in the lower part of the subsoil and in the underlying material
- Soils that have carbonates within a depth of 20 inches

Dissimilar soils:

- The well drained Rozetta soils on shoulders and summits

Properties and Qualities of the Navlys Soil

Parent material: Loess

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.1 inches to a depth of 60 inches

Content of organic matter in the surface layer: 0.5 to 1.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: 4 to 6 feet below the surface

Flooding: None

Accelerated erosion: The surface layer is mostly subsoil material.

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Medium

Susceptibility to water erosion: High

Susceptibility to wind erosion: Very low

Interpretive Groups

Land capability classification: Navlys—4e

Prime farmland status: Navlys—not prime farmland

Hydric soil status: Navlys—not hydric

801C—Orthents, silty, rolling

Setting

Landform: Ground moraines

Position on landform: Summits and backslopes in cut and fill and borrow areas where the soils have been disturbed; including the proving ground area for heavy construction equipment near Mechanicsburg

Map Unit Composition

Orthents: 85 percent

Dissimilar soils: 15 percent

Minor Components

Dissimilar soils:

- The moderately well drained Buckhart soils in undisturbed areas
- The somewhat poorly drained Ipava soils in undisturbed areas
- The well drained Hickory and Rozetta soils in undisturbed areas

Properties and Qualities of the Orthents

Parent material: Earthy fill consisting of silty material derived from former soil layers and silty underlying material

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:
Moderately slow

Permeability below a depth of 60 inches: Moderate or moderately slow

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.0 inches to a depth of 60 inches

Content of organic matter in the surface layer: 0.2 to 1.0 percent

Shrink-swell potential: Moderate

Flooding: None

Potential for frost action: High

Corrosivity: High for steel and moderate for concrete

Surface runoff class: Medium

Susceptibility to water erosion: High

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Orthents—3e

Prime farmland status: Orthents—not prime farmland

Hydric soil status: Orthents—not hydric

Oscos Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Typic Argiudolls

Typical Pedon (OSD)

Oscos silt loam, 2 to 5 percent slopes, at an elevation of 858 feet; Carroll County, Illinois; 316 feet north and 88 feet west of the southeast corner of sec. 23, T. 24 N., R. 6 E.; USGS Lanark quadrangle; lat. 42 degrees 03 minutes 13.4 seconds N. and long. 89 degrees 45 minutes 48.2 seconds W., NAD 83:

Ap—0 to 10 inches; very dark brown (10YR 2/2) silt loam, very dark grayish brown (10YR 3/2) dry; moderate fine granular structure; friable; common fine roots; slightly acid; abrupt smooth boundary.

A—10 to 14 inches; very dark grayish brown (10YR 3/2) silt loam, dark grayish brown (10YR 4/2) dry; moderate medium and coarse granular structure; friable; common fine roots; strongly acid; clear smooth boundary.

BA—14 to 20 inches; dark yellowish brown (10YR 3/4) and dark brown (10YR 3/3) silt loam, brown (10YR 5/3) dry; weak fine subangular blocky structure; friable; common fine roots; few distinct light brownish gray (10YR 6/2 dry) silt coatings on faces of peds; strongly acid; clear smooth boundary.

Bt1—20 to 26 inches; brown (10YR 4/3) silty clay loam; moderate fine subangular blocky structure; friable; few fine roots; few distinct gray (10YR 6/1 dry) silt coatings and common faint dark brown (10YR 3/3) organo-clay films on faces of peds; strongly acid; clear smooth boundary.

Bt2—26 to 37 inches; dark yellowish brown (10YR 4/4) silty clay loam; moderate medium subangular blocky structure; firm; few fine roots; common distinct light brownish gray (10YR 6/2 dry) silt coatings and many faint dark yellowish brown (10YR 4/4) clay films on faces of peds; common fine faint brown (10YR 5/3) and common medium prominent strong brown (7.5YR 5/8) accumulations of iron and manganese; many prominent very dark gray (N 3/0) and dark brown (7.5YR 3/2) manganese concretions; strongly acid; clear smooth boundary.

Bt3—37 to 45 inches; light yellowish brown (10YR 6/4) silty clay loam; moderate coarse subangular blocky structure; friable; few fine roots; many faint dark yellowish brown (10YR 4/4) clay films on faces of peds; common fine distinct light brownish gray (10YR 6/2) iron depletions and few medium prominent strong brown (7.5YR 5/8) accumulations of iron and manganese; strongly acid; gradual smooth boundary.

BC—45 to 55 inches; yellowish brown (10YR 5/4) and brown (10YR 4/3) silty clay loam; weak coarse angular blocky structure; friable; few fine distinct light brownish gray (10YR 6/2) iron depletions; strongly acid; gradual smooth boundary.

C—55 to 60 inches; yellowish brown (10YR 5/4) and brown (10YR 4/3) silt loam; massive; friable; many fine distinct yellowish brown (10YR 5/6) accumulations of iron and manganese and

common medium distinct grayish brown (10YR 5/2) iron depletions; moderately acid.

Range in Characteristics

Thickness of the mollic epipedon: 10 to 18 inches

Depth to base of diagnostic horizon: 40 to more than 66 inches

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 or 2

Texture—silt loam

Bt horizon(s):

Hue—10YR

Value—4 to 6

Chroma—3 or 4

Texture—silty clay loam or silt loam

C horizon(s):

Hue—10YR

Value—4 or 5

Chroma—3 to 6

Texture—silt loam or silty clay loam

Taxadjunct Feature

Oscosilt loam, 5 to 10 percent slopes, eroded, has a dark surface soil that is thinner than is definitive for the series. This difference, however, does not significantly affect the use and management of the soil. The soil is classified as a fine-silty, mixed, superactive, mesic Mollic Hapludalf.

86B—Oscosilt loam, 2 to 5 percent slopes

Setting

Landform: Ground moraines (fig. 5)

Position on landform: Summits and shoulders

Map Unit Composition

Oscos and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a thinner dark surface soil
- Soils that have carbonates within a depth of 40 inches
- Soils that have more sand in the lower part of the subsoil and in the underlying material

Dissimilar soils:

- The somewhat poorly drained Ipava soils on summits

Properties and Qualities of the Osco Soil*Parent material:* Loess*Drainage class:* Well drained*Slowest permeability within a depth of 40 inches:*
Moderate*Permeability below a depth of 60 inches:* Moderate*Depth to restrictive feature:* More than 80 inches*Available water capacity:* About 11.8 inches to a depth of 60 inches*Content of organic matter in the surface layer:* 3.0 to 4.0 percent*Shrink-swell potential:* Moderate*Apparent seasonal high water table:* 4 to 6 feet below the surface*Flooding:* None*Potential for frost action:* High*Corrosivity:* Moderate for steel and concrete*Surface runoff class:* Low*Susceptibility to water erosion:* Moderate*Susceptibility to wind erosion:* Low**Interpretive Groups***Land capability classification:* Osco—2e*Prime farmland status:* Osco—prime farmland in all areas*Hydric soil status:* Osco—not hydric**86C2—Osco silt loam, 5 to 10 percent slopes, eroded****Setting***Landform:* Ground moraines*Position on landform:* Shoulders and backslopes**Map Unit Composition**

Osco and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components*Similar soils:*

- Soils that have carbonates within a depth of 40 inches
- Soils that have more clay in the surface layer
- Soils that have a thicker dark surface soil
- Soils that have more sand in the lower part of the subsoil and in the underlying material

Dissimilar soils:

- The somewhat poorly drained Ipava soils on summits

Properties and Qualities of the Osco Soil*Parent material:* Loess*Drainage class:* Well drained*Slowest permeability within a depth of 40 inches:*
Moderate*Permeability below a depth of 60 inches:* Moderate*Depth to restrictive feature:* More than 80 inches*Available water capacity:* About 11.7 inches to a depth of 60 inches*Content of organic matter in the surface layer:* 2.0 to 3.0 percent*Shrink-swell potential:* Moderate*Apparent seasonal high water table:* 4 to 6 feet below the surface*Flooding:* None*Accelerated erosion:* The surface layer has been thinned by erosion.*Potential for frost action:* High*Corrosivity:* Moderate for steel and concrete*Surface runoff class:* Medium*Susceptibility to water erosion:* Moderate*Susceptibility to wind erosion:* Low**Interpretive Groups***Land capability classification:* Osco—3e*Prime farmland status:* Osco—not prime farmland*Hydric soil status:* Osco—not hydric**864—Pits, quarries****Setting**

This map unit consists of open excavations from which limestone and/or shale has been removed or is currently being removed.

Map Unit Composition

Pits, quarries: 90 percent

Dissimilar components: 10 percent

Minor Components*Dissimilar components:*

- The well drained Marseilles soils in undisturbed areas
- Areas where the soil has been mixed with rock fragments and other debris during mining operations
- Stockpiles of crushed rock
- Areas of machinery
- Areas of water less than 3 acres in size

Interpretive Groups*Land capability classification:* Pits—none assigned*Prime farmland status:* Pits—not prime farmland

Hydric soil status: Pits—unranked

862—Pits, sand

Setting

This map unit consists of open excavations from which sand has been removed or is currently being removed.

Map Unit Composition

Pits, sand: 90 percent

Dissimilar components: 10 percent

Minor Components

Dissimilar components:

- The well drained Alvin, Camden, and Plano soils in undisturbed areas
- Areas of water less than 3 acres in size
- Areas where the soil has been disturbed by grading or by sand-processing operations
- Stockpiles of sand
- Areas of machinery

Interpretive Groups

Land capability classification: Pits—none assigned

Prime farmland status: Pits—not prime farmland

Hydric soil status: Pits—unranked

Plano Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Typic Arguidolls

Typical Pedon (OSD)

Plano silt loam, 0 to 2 percent slopes, at an elevation of 715 feet; Stark County, Illinois; 1,200 feet south and 1,920 feet east of the northwest corner of sec. 13, T. 12 N., R. 7 E.; USGS Castleton topographic quadrangle; lat. 41 degrees 01 minute 45 seconds N. and long. 89 degrees 39 minutes 00 seconds W., NAD 27:

Ap—0 to 9 inches; very dark brown (10YR 2/2) silt loam, grayish brown (10YR 5/2) dry; moderate fine granular structure; friable; few very fine roots; slightly acid; clear smooth boundary.

A—9 to 14 inches; dark brown (10YR 3/3) silt loam, brown (10YR 5/3) dry; moderate fine granular structure; friable; many very fine roots; slightly acid; clear smooth boundary.

Bt1—14 to 19 inches; dark yellowish brown (10YR 4/4) silty clay loam; moderate fine subangular blocky structure; friable; common very fine roots; many

distinct dark brown (10YR 3/3) organo-clay films on faces of peds; slightly acid; clear smooth boundary.

Bt2—19 to 31 inches; dark yellowish brown (10YR 4/4) silty clay loam; moderate medium subangular blocky structure; friable; common very fine roots; many distinct brown (10YR 4/3) clay films on faces of peds; slightly acid; clear smooth boundary.

Bt3—31 to 43 inches; dark yellowish brown (10YR 4/4) silty clay loam; moderate medium prismatic structure parting to moderate medium subangular blocky; friable; few very fine roots; many distinct brown (10YR 4/3) clay films on faces of peds; common distinct very pale brown (10YR 7/3 dry) clay depletions on faces of peds; few fine faint yellowish brown (10YR 5/4) accumulations of iron in the matrix; slightly acid; clear smooth boundary.

Bt4—43 to 49 inches; dark yellowish brown (10YR 4/4) silt loam; moderate medium prismatic structure; friable; few very fine roots; many distinct brown (10YR 4/3) clay films on faces of peds; few distinct very pale brown (10YR 7/3 dry) clay depletions on faces of peds; slightly acid; clear smooth boundary.

2Bt5—49 to 53 inches; dark yellowish brown (10YR 4/4) clay loam; weak medium prismatic structure; friable; few fine roots; many distinct brown (10YR 4/3) clay films on faces of peds; neutral; clear smooth boundary.

2BC—53 to 60 inches; brown (7.5YR 4/4) sandy loam; weak medium subangular blocky structure; very friable; many distinct dark yellowish brown (10YR 3/4) clay films bridging sand grains; about 5 percent gravel; neutral; gradual smooth boundary.

2C—60 to 72 inches; stratified yellowish brown (10YR 5/6) and brown (7.5YR 4/4) sandy loam, loam, and loamy sand; massive; friable; about 12 percent gravel; neutral.

Range in Characteristics

Thickness of the loess: 40 to 60 inches

Thickness of the mollic epipedon: 10 to 20 inches

Depth to base of diagnostic horizon: 44 to 70 inches

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 to 3

Texture—silt loam

AB or BA horizon(s), where present:

Hue—10YR

Value—3 or 4

Chroma—2 to 4

Texture—silt loam or silty clay loam

Bt horizon (upper and middle parts):

Hue—10YR

Value—4 or 5

Chroma—3 or 4

Texture—silt loam or silty clay loam

Bt horizon (lower part):

Hue—7.5YR or 10YR

Value—3 to 5

Chroma—2 to 4

Texture—silt loam or silty clay loam

2Bt or 2BC horizon(s):

Hue—7.5YR or 10YR

Value—3 to 5

Chroma—2 to 6

Texture—silt loam, loam, sandy loam, clay loam, or sandy clay loam

2C horizon(s):

Hue—7.5YR, 10YR, or 2.5Y

Value—3 to 5

Chroma—3 to 6

Texture—loam, loamy sand, sandy loam, or silt loam

199A—Plano silt loam, 0 to 2 percent slopes

Setting

Landform: Stream terraces

Position on landform: Summits

Map Unit Composition

Plano and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have more sand in the upper part of the subsoil
- Soils that have a thinner dark surface soil
- Soils that have less sand in the lower part of the subsoil and in the underlying material

Dissimilar soils:

- The poorly drained Vesser soils on flood plains
- The somewhat poorly drained Elburn and Kendall soils on summits and footslopes

Properties and Qualities of the Plano Soil

Parent material: Loess over outwash

Drainage class: Well drained

Slowest permeability within a depth of 40 inches: Moderate

Permeability below a depth of 60 inches: Moderate rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.3 inches to a depth of 60 inches

Content of organic matter in the surface layer: 3.0 to 5.0 percent

Shrink-swell potential: Moderate

Flooding: None

Potential for frost action: High

Corrosivity: Moderate for steel and low for concrete

Surface runoff class: Low

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Plano—1

Prime farmland status: Plano—prime farmland in all areas

Hydric soil status: Plano—not hydric

199B—Plano silt loam, 2 to 5 percent slopes

Setting

Landform: Stream terraces (fig. 3)

Position on landform: Summits

Map Unit Composition

Plano and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have more sand in the upper part of the subsoil
- Soils that have a thinner dark surface soil
- Soils that have less sand in the lower part of the subsoil and in the underlying material

Dissimilar soils:

- The somewhat poorly drained Elburn and Kendall soils on summits and footslopes

Properties and Qualities of the Plano Soil

Parent material: Loess over outwash

Drainage class: Well drained

Slowest permeability within a depth of 40 inches: Moderate

Permeability below a depth of 60 inches: Moderately rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.1 inches to a depth of 60 inches

Content of organic matter in the surface layer: 3.0 to 5.0 percent

Shrink-swell potential: Moderate

Flooding: None

Potential for frost action: High

Corrosivity: Moderate for steel and low for concrete

Surface runoff class: Low

Susceptibility to water erosion: Moderate

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Plano—2e

Prime farmland status: Plano—prime farmland in all areas

Hydric soil status: Plano—not hydric

Proctor Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Typic Argiudolls

Typical Pedon (OSD)

Proctor silt loam, 0 to 2 percent slopes, at an elevation of 705 feet; Peoria County, Illinois; 204 feet north and 2,460 feet west of the southeast corner of sec. 3, T. 11 N., R. 6 E.; USGS Princeville topographic quadrangle; lat. 40 degrees 57 minutes 37 seconds N. and long. 89 degrees 47 minutes 59 seconds W., NAD 27:

Ap—0 to 8 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; weak fine granular structure; friable; common very fine roots; moderately acid; clear smooth boundary.

A—8 to 11 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; moderate fine granular structure; friable; common very fine roots; neutral; clear smooth boundary.

Bt1—11 to 16 inches; dark yellowish brown (10YR 4/4) silty clay loam; moderate very fine subangular blocky structure; friable; common very fine roots; common distinct very dark grayish brown (10YR 3/2) organo-clay films on faces of peds; moderately acid; clear smooth boundary.

Bt2—16 to 23 inches; dark yellowish brown (10YR 4/4) silty clay loam; moderate very fine and fine

subangular blocky structure; friable; common very fine roots; many distinct brown (10YR 4/3) clay films on faces of peds; moderately acid; clear smooth boundary.

Bt3—23 to 28 inches; yellowish brown (10YR 5/4) silty clay loam; moderate fine subangular blocky structure; friable; common very fine roots; many distinct brown (10YR 4/3) clay films on faces of peds; moderately acid; clear smooth boundary.

2Bt4—28 to 33 inches; yellowish brown (10YR 5/4) loam; moderate medium subangular blocky structure; friable; few very fine roots; common faint dark yellowish brown (10YR 4/4) clay films on faces of peds; moderately acid; clear smooth boundary.

2Bt5—33 to 46 inches; strong brown (7.5YR 5/6), stratified loam and sandy loam; weak coarse subangular blocky structure; very friable; few very fine roots; common faint brown (7.5YR 4/4) clay films on faces of peds; slightly acid; gradual smooth boundary.

2C—46 to 60 inches; strong brown (7.5YR 5/6), stratified sandy loam and loamy sand; massive; very friable; slightly acid.

Range in Characteristics

Thickness of the loess: 20 to 40 inches

Thickness of the mollic epipedon: 10 to 20 inches

Depth to base of diagnostic horizon: 40 to 65 inches

Ap, A, or AB horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 to 3

Texture—silt loam or silty clay loam

Bt or BA horizon(s):

Hue—7.5YR or 10YR

Value—3 to 6

Chroma—3 to 6

Texture—silty clay loam or silt loam

2Bt or 2BC horizon(s):

Hue—7.5YR, 10YR, or 2.5Y

Value—4 to 6

Chroma—3 to 6

Texture—silty clay loam, silt loam, clay loam, sandy clay loam, loam, or sandy loam

2C horizon(s):

Hue—7.5YR, 10YR, or 2.5Y

Value—4 to 6

Chroma—3 to 6

Texture—sandy loam, loam, silt loam, loamy sand, or sand

7148A—Proctor silt loam, 0 to 2 percent slopes, rarely flooded

Setting

Landform: Flood-plain steps

Map Unit Composition

Proctor and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have less sand in the lower part of the subsoil
- Soils that have more sand in the upper part of the subsoil
- Soils that have a thinner dark surface soil

Dissimilar soils:

- The well drained Camden soils on summits and risers
- The somewhat poorly drained Kendall soils on summits
- The poorly drained Sawmill soils on flood plains

Properties and Qualities of the Proctor Soil

Parent material: Loess or other silty material

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:

Moderate

Permeability below a depth of 60 inches: Moderately

rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 10.6 inches to a depth of 60 inches

Content of organic matter in the surface layer: 3.0 to 4.0 percent

Shrink-swell potential: Moderate

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Low

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Proctor—1

Prime farmland status: Proctor—prime farmland in all areas

Hydric soil status: Proctor—not hydric

Radford Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Fluvaquentic Hapludolls

Typical Pedon (OSD)

Radford silt loam, 0 to 2 percent slopes, frequently flooded (fig. 8), at an elevation of 567 feet; Cass County, Illinois; 2,700 feet east and 1,320 feet south of the northeast corner of sec. 2, T. 17 N., R. 9 W.; USGS Ashland topographic quadrangle; lat. 39 degrees 56 minutes 26 seconds and long. 90 degrees 04 minutes 43 seconds W.; NAD 27:

Ap—0 to 7 inches; very dark grayish brown (10YR 3/2) silt loam, brown (10YR 5/3 dry); weak medium subangular blocky structure parting to moderate fine and medium granular; friable; few very fine roots; neutral; clear smooth boundary.

A—7 to 12 inches; very dark grayish brown (10YR 3/2) silt loam, brown (10YR 5/3) dry; weak medium subangular blocky structure; friable; few very fine roots; neutral; clear smooth boundary.

C—12 to 33 inches; dark grayish brown (10YR 4/2) and very dark grayish brown (10YR 3/2) silt loam with common thin grayish brown (10YR 5/2) and brown (10YR 5/3) lenses; massive; friable; few very fine roots; common faint very dark grayish brown (10YR 3/2) organic coatings in worm channels; few fine rounded black (7.5YR 2.5/1) weakly cemented manganese concretions with diffuse boundaries throughout; neutral; clear smooth boundary.

Ab1—33 to 42 inches; very dark gray (10YR 3/1) silt loam; weak fine subangular blocky structure parting to moderate medium granular; friable; few very fine roots; few fine rounded black (7.5YR 2.5/1) weakly cemented manganese concretions with diffuse boundaries lining root channels and pores; few fine prominent strong brown (7.5YR 5/8) accumulations of iron in the matrix; slightly alkaline; gradual smooth boundary.

Ab2—42 to 72 inches; very dark gray (10YR 3/1) silt loam; moderate fine subangular blocky structure; friable; few very fine roots; few distinct gray (10YR 6/1 dry) clay depletions on faces of peds; few fine rounded black (7.5YR 2.5/1) weakly cemented manganese concretions with diffuse boundaries lining root channels and pores; few fine prominent strong brown (7.5YR 5/8) accumulations of iron in the matrix; slightly alkaline; clear smooth boundary.

Bgb—72 to 80 inches; grayish brown (10YR 5/2) silt loam; moderate medium prismatic structure parting to weak medium subangular blocky;

friable; few very fine roots; common distinct very dark gray (10YR 3/1) organic coatings lining root channels and pores; slightly alkaline.



Figure 8.—Typical profile of Radford silt loam, 0 to 2 percent slopes, frequently flooded.

Range in Characteristics

Thickness of the mollic epipedon: 10 to 20 inches

Depth to the buried soil: 20 to 40 inches

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 or 2

Texture—silt loam

C horizon(s):

Hue—10YR

Value—2 to 6

Chroma—1 to 4

Texture—silt loam

Ab horizon(s):

Hue—10YR or neutral

Value—2 or 3

Chroma—0 or 1

Texture—silt loam, silty clay loam, clay loam, or loam

Bgb horizon(s), where present:

Hue—10YR, 2.5Y, 5Y, or neutral

Value—3 to 6

Chroma—0 to 2

Texture—silt loam, silty clay loam, clay loam, or loam

3074A—Radford silt loam, 0 to 2 percent slopes, frequently flooded

Setting

Landform: Flood plains

Map Unit Composition

Radford and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a lighter colored surface soil
- Soils that have a buried soil at a depth of more than 40 inches
- Soils that have a buried soil within a depth of 20 inches

Dissimilar soils:

- The poorly drained Sawmill soils on flood plains

Properties and Qualities of the Radford Soil

Parent material: Alluvium

Drainage class: Somewhat poorly drained

Slowest permeability within a depth of 40 inches:

Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.3 inches to a depth of 60 inches

Content of organic matter in the surface layer: 2.0 to 4.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: 1 to 2 feet below the surface

Potential for frost action: High

Corrosivity: High for steel and moderate for concrete

Surface runoff class: Low

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Radford—3w

Prime farmland status: Radford—prime farmland where protected from flooding or not frequently flooded during the growing season

Hydric soil status: Radford—not hydric

Ross Series

Taxonomic classification: Fine-loamy, mixed, superactive, mesic Cumulic Hapludolls

Typical Pedon

Ross silt loam, 0 to 2 percent slopes, frequently flooded, at an elevation of 590 feet; Tazewell County, Illinois; 1,490 feet west and 232 feet north of the southeast corner of sec. 28, T. 23 N., R. 3 W.; USGS Hopedale topographical quadrangle; lat. 40 degrees 24 minutes 36 seconds N. and long. 89 degrees 26 minutes 27 seconds W., NAD 27:

Ap—0 to 8 inches; very dark gray (10YR 3/1) silt loam, dark grayish brown (10YR 4/2) dry; weak fine granular structure; friable; common very fine roots; neutral; clear smooth boundary.

A—8 to 13 inches; very dark grayish brown (10YR 3/2) silt loam, dark grayish brown (10YR 4/2) dry; weak fine subangular blocky structure parting to weak fine granular; friable; common very fine and fine roots; common faint very dark gray (10YR 3/1) organic coatings on faces of peds; neutral; clear smooth boundary.

Bw1—13 to 27 inches; very dark grayish brown (10YR 3/2) loam, dark grayish brown (10YR 4/2) dry; weak fine subangular blocky structure; friable; few very fine roots; few faint very dark gray (10YR 3/1)

organic coatings on faces of peds; neutral; gradual smooth boundary.

Bw2—27 to 34 inches; dark brown (10YR 3/3) loam, brown (10YR 4/3) dry; weak fine and medium subangular blocky structure; friable; few very fine and coarse roots; common distinct very dark gray (10YR 3/1) and few faint very dark grayish brown (10YR 3/2) organic coatings on faces of peds; neutral; gradual smooth boundary.

Bw3—34 to 43 inches; brown (10YR 4/3) loam; weak medium subangular blocky structure; very friable; few very fine roots; many distinct very dark grayish brown (10YR 3/2) organic coatings on faces of peds; neutral; gradual smooth boundary.

C1—43 to 54 inches; brown (10YR 4/3) sandy loam; massive; very friable; few very fine and fine roots; neutral; gradual smooth boundary.

C2—54 to 60 inches; brown (10YR 4/3) sandy loam; massive; very friable; few fine faint grayish brown (10YR 5/2) iron depletions; 5 percent gravel; neutral.

Range in Characteristics

Depth to base of diagnostic horizon: 24 to 45 inches

Thickness of the mollic epipedon: 24 to 40 inches

Depth to carbonates: More than 45 inches

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 to 3

Texture—loam, silt loam, or silty clay loam

Bw horizon(s):

Hue—10YR

Value—2 to 5

Chroma—1 to 4

Texture—sandy loam, loam, silt loam, clay loam, or silty clay loam

C horizon(s):

Hue—10YR, 7.5YR, or 2.5Y

Value—4 to 6

Chroma—1 to 4

Texture—sandy loam, loam, silt loam, sandy clay loam, or the gravelly analogs of those textures; also, strata containing more sand below a depth of 40 inches in some pedons

3073A—Ross silt loam, 0 to 2 percent slopes, frequently flooded

Setting

Landform: Flood plains (fig. 3)

Map Unit Composition

Ross and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have less sand in the subsoil
- Soils that have more sand in the surface layer
- Soils that have less clay throughout
- Soils that have a thinner dark surface soil
- Soils that have a buried soil

Dissimilar soils:

- The somewhat poorly drained Lawson and Tice soils on flood plains
- The poorly drained Sawmill soils on flood plains

Properties and Qualities of the Ross Soil

Parent material: Alluvium

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:

Moderate

Permeability below a depth of 60 inches: Moderate or moderately rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.6 inches to a depth of 60 inches

Content of organic matter in the surface layer: 2.0 to 4.0 percent

Shrink-swell potential: Low

Apparent seasonal high water table: 4 to 6 feet below the surface

Potential for frost action: Moderate

Corrosivity: Low for steel and concrete

Surface runoff class: Low

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Ross—2w

Prime farmland status: Ross—prime farmland where protected from flooding or not frequently flooded during the growing season

Hydric soil status: Ross—not hydric

Rozetta Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Typic Hapludalfs

Typical Pedon (OSD)

Rozetta silt loam, 0 to 2 percent slopes, at an elevation of 890 feet; Stephenson County, Illinois; 150 feet south

and 500 feet east of the center of sec. 18, T. 27 N., R. 6 E.; USGS Pearl City quadrangle; lat. 42 degrees 20 minutes 00 seconds N. and long. 89 degrees 51 minutes 19 seconds W., NAD 27:

A—0 to 4 inches; very dark gray (10YR 3/1) silt loam, gray (10YR 6/1) dry; weak medium granular structure; friable; many fine roots throughout; moderately acid; clear wavy boundary.

E—4 to 11 inches; dark grayish brown (10YR 4/2) silt loam; weak medium platy structure; friable; many fine roots throughout; strongly acid; clear smooth boundary.

BE—11 to 14 inches; brown (10YR 4/3) silty clay loam; weak medium subangular blocky structure; firm; many fine roots between pedes; few faint brown (10YR 5/3 dry) clay depletions on faces of pedes; strongly acid; clear smooth boundary.

Bt1—14 to 21 inches; dark yellowish brown (10YR 4/4) silty clay loam; moderate fine and medium subangular blocky structure; firm; many fine roots between pedes; many faint brown (10YR 5/3) clay films on faces of pedes; strongly acid; clear smooth boundary.

Bt2—21 to 39 inches; brown (10YR 5/3) silty clay loam; moderate medium and coarse subangular blocky structure; firm; common fine roots; many faint dark yellowish brown (10YR 4/4) clay films on faces of pedes; common faint pale brown (10YR 6/3 dry) silt coatings on faces of pedes; few medium faint grayish brown (10YR 5/2) iron depletions in the matrix; common medium faint light yellowish brown (10YR 6/4) and brown (10YR 4/3) accumulations of iron in the matrix; strongly acid; clear smooth boundary.

Bt3—39 to 50 inches; yellowish brown (10YR 5/4) silty clay loam; weak coarse subangular blocky structure; firm; common fine roots; few faint brown (10YR 4/3) clay films on faces of pedes; common medium distinct grayish brown (10YR 5/2) iron depletions in the matrix; common medium faint pale brown (10YR 6/3) accumulations of iron in the matrix; moderately acid; clear smooth boundary.

C—50 to 60 inches; yellowish brown (10YR 5/4) silt loam; massive; friable; common medium distinct dark grayish brown (10YR 4/2) iron depletions in the matrix; slightly acid.

Range in Characteristics

Depth to base of diagnostic horizon: 42 to 72 inches

Ap or A horizon:
Hue—10YR

Value—3 to 5
 Chroma—1 to 3
 Texture—silt loam

E horizon, where present:

Hue—10YR
 Value—4 to 6
 Chroma—2 or 3
 Texture—silt loam

Bt horizon:

Hue—10YR or 7.5YR
 Value—4 to 6
 Chroma—3 to 6
 Texture—silty clay loam

C horizon:

Hue—10YR
 Value—4 to 6
 Chroma—2 to 6
 Texture—silt loam or silty clay loam

279B—Rozetta silt loam, 2 to 5 percent slopes

Setting

Landform: Ground moraines (fig. 4)

Position on landform: Shoulders and summits

Map Unit Composition

Rozetta and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a thicker, darker surface layer
- Soils that have a seasonal high water table at a depth of more than 6 feet
- Soils that have more sand in the lower part of the subsoil and in the underlying material
- Soils that have carbonates within a depth of 40 inches

Dissimilar soils:

- The somewhat poorly drained Keomah soils on summits
- The well drained Hickory soils on backslopes

Properties and Qualities of the Rozetta Soil

Parent material: Loess

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:
 Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.3 inches to a depth of 60 inches

Content of organic matter in the surface layer: 1.0 to 3.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: 4 to 6 feet below the surface

Flooding: None

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Low

Susceptibility to water erosion: Moderate

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Rozetta—2e

Prime farmland status: Rozetta—prime farmland in all areas

Hydric soil status: Rozetta—not hydric

Sable Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Typic Endoaquolls

Typical Pedon (OSD)

Sable silty clay loam, 0 to 2 percent slopes (fig. 9), at an elevation of 732 feet; Warren County, Illinois; 1,281 feet south and 97 feet west of the northeast corner of sec. 14, T. 9 N., R. 3 W.; USGS Kirkwood East topographic quadrangle; lat. 40 degrees 46 minutes 22.4 seconds N. and long. 90 degrees 41 minutes 33.7 seconds W., NAD 27:

Ap—0 to 8 inches; black (10YR 2/1) silty clay loam, dark gray (10YR 4/1) dry; moderate fine and medium granular structure; firm; moderately acid; abrupt smooth boundary.

A—8 to 19 inches; black (10YR 2/1) silty clay loam, dark gray (10YR 4/1) dry; moderate very fine angular blocky structure; firm; few fine rounded dark reddish brown (5YR 3/2) concretions of iron and manganese throughout; slightly acid; clear smooth boundary.

AB—19 to 23 inches; very dark gray (10YR 3/1) silty clay loam, grayish brown (10YR 5/2) dry; moderate fine angular blocky structure; firm; few faint very dark grayish brown (10YR 3/2) organic coatings on faces of peds; few fine rounded dark reddish brown (5YR 3/2) concretions of iron and manganese throughout; slightly acid; clear smooth boundary.

Bg—23 to 29 inches; dark gray (10YR 4/1) silty clay



Figure 9.—Typical profile of Sable silty clay loam, 0 to 2 percent slopes.

loam; moderate fine and medium subangular blocky structure; firm; common faint very dark gray (10YR 3/1) organic coatings on faces of peds; common fine and medium rounded dark reddish brown (5YR 3/2) concretions of iron and manganese throughout; common medium distinct brown (10YR 5/3) accumulations of iron in the matrix; few medium faint dark grayish brown (10YR 4/2) iron depletions in the matrix; neutral; clear smooth boundary.

Btg1—29 to 38 inches; grayish brown (2.5Y 5/2) silty clay loam; moderate medium and coarse subangular blocky structure; firm; few distinct dark gray (10YR 4/1) clay films on faces of peds; many fine and medium rounded dark reddish brown (5YR 3/2) concretions of iron and manganese throughout; many medium prominent yellowish brown (10YR 5/6) accumulations of iron in the matrix; neutral; clear wavy boundary.

Btg2—38 to 47 inches; gray (N 5/0) silt loam; weak

medium prismatic structure parting to weak medium and coarse angular blocky; firm; few distinct grayish brown (10YR 5/2) clay films on faces of peds; common fine rounded dark reddish brown (5YR 3/2) concretions of iron and manganese throughout; many medium prominent yellowish brown (10YR 5/6) accumulations of iron in the matrix; slightly alkaline; gradual smooth boundary.

Cg—47 to 60 inches; gray (N 5/0) silt loam; massive; friable; many medium prominent yellowish brown (10YR 5/6) accumulations of iron in the matrix; slightly effervescent; slightly alkaline.

Range in Characteristics

Thickness of the mollic epipedon: 12 to 24 inches

Depth to free carbonates: More than 40 inches

Depth to base of diagnostic horizon: 40 to 60 inches

Ap, A, AB, or BA horizon(s):

Hue—10YR, 5Y, or neutral

Value—2 or 3

Chroma—0 or 1

Texture—silty clay loam or silt loam

Btg, Bg, BC, or BCg horizon(s):

Hue—10YR, 2.5Y, 5Y, or neutral

Value—3 to 6

Chroma—0 to 2

Texture—silty clay loam or silt loam

Cg horizon(s):

Hue—10YR, 2.5Y, 5Y, or neutral

Value—3 to 6

Chroma—0 to 2

Texture—silt loam or silty clay loam

68A—Sable silty clay loam, 0 to 2 percent slopes

Setting

Landform: Ground moraines (fig. 2)

Position on landform: Toeslopes

Map Unit Composition

Sable and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils in which the dark surface soil is more than 24 inches thick
- Soils that have more clay in the subsoil
- Soils that have more sand in the lower part of the subsoil and in the underlying material

- Soils that have carbonates within a depth of 40 inches

Dissimilar soils:

- The moderately well drained Buckhart soils on summits and backslopes
- The well drained Osco soils on summits and shoulders

Properties and Qualities of the Sable Soil

Parent material: Loess

Drainage class: Poorly drained

Slowest permeability within a depth of 40 inches:

Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.9 inches to a depth of 60 inches

Content of organic matter in the surface layer: 5.0 to 6.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: At the surface to 1 foot below the surface

Ponding: At the surface to 0.5 foot above the surface

Flooding: None

Potential for frost action: High

Corrosivity: High for steel and low for concrete

Surface runoff class: Negligible

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Very low

Interpretive Groups

Land capability classification: Sable—2w

Prime farmland status: Sable—prime farmland where drained

Hydric soil status: Sable—hydric

Sawmill Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Cumulic Endoaquolls

Typical Pedon (OSD)

Sawmill silty clay loam, 0 to 2 percent slopes, frequently flooded, at an elevation of 535 feet; Sangamon County, Illinois; 300 feet south and 750 feet east of the northwest corner of sec. 20, T. 15 N., R. 4 W.; USGS New City topographic quadrangle; lat. 39 degrees 44 minutes 34 seconds N. and long. 89 degrees 34 minutes 15 seconds W., NAD 27:

Ap—0 to 10 inches; very dark gray (10YR 3/1) and very dark grayish brown (10YR 3/2) silty clay loam, gray (10YR 5/1) dry; weak fine subangular

blocky structure; firm; few fine roots; few subrounded pebbles 1 to 3 millimeters in diameter; slightly acid; clear smooth boundary.

A1—10 to 17 inches; black (10YR 2/1) and very dark grayish brown (10YR 3/2) silty clay loam, dark gray (10YR 4/1) dry; moderate fine subangular blocky structure; firm; few fine roots; few subrounded pebbles 1 to 3 millimeters in diameter; few fine rounded black (7.5YR 2.5/1) weakly cemented concretions of iron and manganese with diffuse boundaries lining root channels and pores; few fine prominent yellowish brown (10YR 5/6) accumulations of iron in the matrix; neutral; clear smooth boundary.

A2—17 to 25 inches; black (10YR 2/1) silty clay loam, dark gray (10YR 4/1) dry; moderate fine and medium angular blocky structure; firm; few fine roots; few fine rounded black (7.5YR 2.5/1) weakly cemented concretions of iron and manganese with diffuse boundaries lining root channels and pores; few fine prominent yellowish brown (10YR 5/6) accumulations of iron in the matrix; neutral; clear smooth boundary.

AB—25 to 32 inches; very dark gray (10YR 3/1) silty clay loam, gray (10YR 5/1) dry; weak medium prismatic structure parting to moderate fine subangular blocky; firm; few fine roots; few fine rounded black (7.5YR 2.5/1) weakly cemented concretions of iron and manganese with diffuse boundaries lining root channels and pores; few fine prominent yellowish brown (10YR 5/6) accumulations of iron in the matrix; neutral; clear smooth boundary.

Bg—32 to 40 inches; dark gray (10YR 4/1) silty clay loam; weak medium prismatic structure parting to moderate fine and medium angular blocky; firm; common faint very dark gray (10YR 3/1) organic coatings on faces of peds; few fine roots; few fine rounded black (7.5YR 2.5/1) weakly cemented concretions of iron and manganese with diffuse boundaries lining root channels and pores; few fine prominent strong brown (7.5YR 5/6) accumulations of iron in the matrix; slightly alkaline; clear smooth boundary.

Btg1—40 to 49 inches; grayish brown (10YR 5/2) silty clay loam; moderate medium prismatic structure parting to weak medium angular blocky; firm; common distinct dark gray (10YR 4/1) clay films on faces of peds; few fine rounded black (7.5YR 2.5/1) weakly cemented concretions of iron and manganese with diffuse boundaries lining root channels and pores; few fine prominent strong brown (7.5YR 5/6) and common fine distinct yellowish brown (10YR 5/4) accumulations of iron

in the matrix; slightly alkaline; clear smooth boundary.

Btg2—49 to 58 inches; grayish brown (2.5Y 5/2) silty clay loam; moderate medium prismatic structure; firm; common distinct gray (10YR 5/1) clay films on faces of ped; few fine rounded black (7.5YR 2.5/1) weakly cemented concretions of iron and manganese with diffuse boundaries lining pores; few fine prominent yellowish brown (10YR 5/6) accumulations of iron in the matrix; slightly alkaline; clear smooth boundary.

Cg—58 to 65 inches; grayish brown (2.5Y 5/2) silty clay loam; massive; firm; very dark gray (10YR 3/1) channel linings and fillings; many medium prominent yellowish brown (10YR 5/6) accumulations of iron lining pores; slightly alkaline.

Range in Characteristics

Thickness of the mollic epipedon: 24 to 36 inches

Depth to base of diagnostic horizon: 36 to 60 inches

Ap, A, or AB horizon(s):

Hue—10YR, 2.5Y, 5Y, or neutral

Value—2 or 3

Chroma—0 to 2

Texture—silty clay loam or silt loam

Bg or Btg horizon(s):

Hue—10YR, 2.5Y, or 5Y

Value—3 to 6

Chroma—1 or 2

Texture—silty clay loam

Cg horizon(s):

Hue—10YR, 2.5Y, or 5Y

Value—3 to 6

Chroma—1 or 2

Texture—silty clay loam, clay loam, silt loam, or loam

3107A—Sawmill silty clay loam, 0 to 2 percent slopes, frequently flooded

Setting

Landform: Flood plains (fig. 3)

Map Unit Composition

Sawmill and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a thinner dark surface soil
- Soils that have more clay in the subsoil
- Soils that have a thicker dark surface soil

Dissimilar soils:

- The well drained Huntsville and Ross soils on flood plains

Properties and Qualities of the Sawmill Soil

Parent material: Alluvium

Drainage class: Poorly drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.3 inches to a depth of 60 inches

Content of organic matter in the surface layer: 4.0 to 5.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: At the surface to 1 foot below the surface

Ponding: At the surface to 0.5 foot above the surface

Potential for frost action: High

Corrosivity: High for steel and low for concrete

Surface runoff class: Negligible

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Very low

Interpretive Groups

Land capability classification: Sawmill—3w

Prime farmland status: Sawmill—prime farmland where drained and either protected from flooding or not frequently flooded during the growing season

Hydric soil status: Sawmill—hydric

Shiloh Series

Taxonomic classification: Fine, smectitic, mesic
Cumulic Vertic Endoaquolls

Typical Pedon

Shiloh silty clay loam, 0 to 2 percent slopes, at an elevation of 595 feet; Christian County, Illinois; 2,600 feet east and 132 feet south of the northwest corner of sec. 34, T. 16 N., R. 1 W.; USGS Niantic topographical quadrangle; lat. 39 degrees 48 minutes 0 seconds N. and long. 89 degrees 11 minutes 18 seconds W., NAD 83:

- Ap—0 to 7 inches; black (10YR 2/1) silty clay loam, dark gray (10YR 4/1) dry; moderate fine granular structure; firm; few very fine roots; few fine rounded black (7.5YR 2/1) iron and manganese concretions; slightly acid; abrupt smooth boundary.
- A—7 to 15 inches; black (N 2/0) silty clay loam, very dark gray (10YR 3/1) dry; moderate very fine

subangular blocky structure; firm; few very fine roots; few fine rounded black (7.5YR 2/1) iron and manganese concretions; neutral; clear smooth boundary.

BA—15 to 27 inches; black (N 2/0) silty clay, very dark gray (10YR 3/1) dry; moderate very fine angular blocky structure; firm; few very fine roots; few fine rounded black (7.5YR 2/1) iron and manganese concretions; neutral; clear smooth boundary.

Bg1—27 to 32 inches; olive gray (5Y 5/2) silty clay; moderate fine subangular blocky structure; firm; few very fine roots; many prominent very dark gray (10YR 3/1) pressure faces on peds; few fine prominent strong brown (7.5YR 5/6) accumulations of iron and manganese in the matrix; neutral; clear smooth boundary.

Bg2—32 to 39 inches; olive gray (5Y 5/2) silty clay loam; moderate medium subangular blocky structure; friable; few very fine roots; many distinct dark grayish brown (10YR 4/2) pressure faces on peds; few prominent very dark gray (10YR 3/1) organic coatings lining pores; few fine rounded black (7.5YR 2/1) iron and manganese concretions; many fine prominent yellowish brown (10YR 5/6) accumulations of iron and manganese throughout; neutral; clear smooth boundary.

Bg3—39 to 52 inches; olive gray (5Y 5/2) silty clay loam; moderate medium subangular blocky structure; friable; few fine roots; few prominent very dark gray (10YR 3/1) organic coatings lining pores; few fine rounded black (10YR 2/1) iron and manganese concretions; many medium prominent yellowish brown (10YR 5/6) accumulations of iron and manganese throughout; neutral; clear smooth boundary.

BCg—52 to 60 inches; olive gray (5Y 5/2) silt loam; weak medium subangular blocky structure; friable; few very fine roots; few prominent very dark gray (10YR 3/1) organic coatings lining pores; many medium prominent yellowish brown (10YR 5/6) accumulations of iron and manganese throughout; neutral; clear smooth boundary.

Cg—60 to 80 inches; gray (10YR 6/1) silt loam; massive; friable; many medium prominent yellowish brown (10YR 5/6) accumulations of iron and manganese throughout; very slightly effervescent; slightly alkaline.

Range in Characteristics

Depth to base of diagnostic horizon: 40 to 70 inches

Thickness of the mollic epipedon: 24 to 48 inches

Depth to carbonates: More than 39 inches

Profile feature: An AB horizon in some pedons

Ap or A horizon(s):

Hue—10YR, 2.5Y, 5Y, or neutral

Value—2 or 3

Chroma—0 to 2

Texture—silty clay loam or silty clay

Bg horizon(s):

Hue—10YR, 2.5Y, 5Y, or neutral

Value—2 to 5

Chroma—0 to 2

Texture—silty clay or silty clay loam

Cg horizon(s):

Hue—10YR, 2.5Y, 5Y, or neutral

Value—2 to 6

Chroma—0 to 2

Texture—silty clay loam, silt loam, or silty clay

138A—Shiloh silty clay loam, 0 to 2 percent slopes

Setting

Landform: Depressions

Map Unit Composition

Shiloh and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils in which the dark surface soil is less than 24 inches thick
- Soils that have less clay in the subsoil
- Soils that have less clay in the surface layer

Dissimilar soils:

- The moderately well drained Buckhart soils on summits
- The well drained Osco soils on summits and shoulders

Properties and Qualities of the Shiloh Soil

Parent material: Loess

Drainage class: Poorly drained

Slowest permeability within a depth of 40 inches:

Moderately slow

Permeability below a depth of 60 inches: Moderately slow

Depth to restrictive feature: More than 80 inches

Available water capacity: About 10.4 inches to a depth of 60 inches

Content of organic matter in the surface layer: 4.0 to 6.0 percent

Shrink-swell potential: High

Apparent seasonal high water table: At the surface to 1 foot below the surface

Ponding: At the surface to 1 foot above the surface

Flooding: None

Potential for frost action: High

Corrosivity: High for steel and low for concrete

Surface runoff class: Negligible

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Very low

Interpretive Groups

Land capability classification: Shiloh—2w

Prime farmland status: Shiloh—prime farmland where drained

Hydric soil status: Shiloh—hydric

Spaulding Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Typic Calciaquolls

Typical Pedon (OSD)

Spaulding silty clay loam, 0 to 2 percent slopes, at an elevation of about 612 feet; Sangamon County, Illinois; about 2 miles east of Buffalo Hart; 2,410 feet east and 1,300 feet south of the northwest corner of sec. 22, T. 17 N., R. 3 W.; USGS Cornland topographic quadrangle; lat. 39 degrees 54 minutes 52 seconds N. and long. 89 degrees 24 minutes 54 seconds W., NAD 27:

Apk—0 to 9 inches; black (10YR 2/1) silty clay loam, very dark gray (10YR 3/1) dry; weak and moderate fine granular structure; friable; many fine roots throughout; few snail shells; violently effervescent; 15 percent calcium carbonate equivalent; moderately alkaline; abrupt smooth boundary.

Ak1—9 to 18 inches; black (10YR 2/1) silty clay loam, dark gray (10YR 4/1) dry; moderate very fine and fine subangular blocky structure; friable; many fine roots throughout; few snail shells; violently effervescent; 22 percent calcium carbonate equivalent; moderately alkaline; clear smooth boundary.

Ak2—18 to 22 inches; very dark gray (10YR 3/1) silty clay loam, gray (10YR 5/1) dry; moderate very fine and fine subangular blocky structure; firm; common fine roots throughout; few fine prominent light olive brown (2.5Y 5/6) accumulations of iron along micropores; few snail shells; violently effervescent; 22 percent calcium carbonate equivalent; moderately alkaline; clear smooth boundary.

Btkg1—22 to 26 inches; dark gray (2.5Y 4/1) silty clay loam; moderate very fine and fine subangular blocky structure; firm; common fine roots throughout; common distinct very dark gray (10YR 3/1) organo-clay films on faces of peds; common distinct black (10YR 2/1) organic coatings in root channels and/or pores; few fine prominent light olive brown (2.5Y 5/6) accumulations of iron along micropores; few fine carbonate nodules; strongly effervescent; 12 percent calcium carbonate equivalent; moderately alkaline; clear smooth boundary.

Btkg2—26 to 32 inches; olive gray (5Y 5/2) silty clay loam; moderate fine and medium subangular blocky structure; firm; common fine roots throughout; few distinct gray (5Y 5/1) clay films on faces of peds; common fine rounded prominent black (10YR 2/1) accumulations of manganese in the matrix; common medium prominent light olive brown (2.5Y 5/6) and yellowish brown (10YR 5/6) accumulations of iron in the matrix; common medium and coarse carbonate nodules; strongly effervescent; 12 percent calcium carbonate equivalent; moderately alkaline; clear smooth boundary.

Btkg3—32 to 38 inches; gray (5Y 6/1) silty clay loam; moderate fine and medium subangular blocky structure; firm; few distinct gray (5Y 5/1) clay films on faces of peds; very few distinct very dark gray (10YR 3/1) organic coatings in root channels and/or pores; many fine prominent light olive brown (2.5Y 5/6) and few fine prominent yellowish brown (10YR 5/6) accumulations of iron in the matrix; few fine carbonate nodules; strongly effervescent; 16 percent calcium carbonate equivalent; moderately alkaline; clear smooth boundary.

BCkg—38 to 44 inches; gray (5Y 6/1) silty clay loam; weak medium subangular blocky structure; firm; few distinct gray (5Y 5/1) clay films in root channels and/or pores; few distinct very dark gray (10YR 3/1) organic coatings in root channels and/or pores; many fine prominent light olive brown (2.5Y 5/6) and few fine prominent yellowish brown (10YR 5/6) accumulations of iron in the matrix; few fine carbonate nodules; strongly effervescent; 16 percent calcium carbonate equivalent; moderately alkaline; clear smooth boundary.

Cg—44 to 80 inches; gray (5Y 6/1) silt loam; massive; friable; many medium prominent strong brown (7.5YR 5/8) accumulations of iron in the matrix; strongly effervescent; 19 percent calcium carbonate equivalent; moderately alkaline.

Range in Characteristics

Thickness of the mollic epipedon: 10 to 24 inches
Depth to carbonates: 0 to 16 inches
Depth to base of diagnostic horizon: 22 to 60 inches
Profile feature: A BCg horizon in some pedons

Apk or Ak horizon(s):

Hue—10YR, 2.5Y, 5Y, or neutral
 Value—2 or 3
 Chroma—0 or 1
 Texture—silty clay loam

Bkg or Btkg horizon(s):

Hue—10YR, 2.5Y, 5Y, or neutral
 Value—3 to 6
 Chroma—0 to 2
 Texture—silty clay loam or silt loam

Cg horizon(s):

Hue—7.5YR, 10YR, 2.5Y, or 5Y
 Value—4 to 6
 Chroma—1 to 8
 Texture—silt loam

712A—Spaulding silty clay loam, 0 to 2 percent slopes

Setting

Landform: Depressions

Map Unit Composition

Spaulding and similar soils: 90 percent
 Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have less clay in the subsoil
- Soils that have carbonates at a depth of more than 16 inches

Dissimilar soils:

- The somewhat poorly drained Ipava soils on summits
- The poorly drained Sable soils in depressions

Properties and Qualities of the Spaulding Soil

Parent material: Calcareous loess

Drainage class: Poorly drained

Slowest permeability within a depth of 40 inches:
 Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 14.0 inches to a depth of 60 inches

Content of organic matter in the surface layer: 4.0 to 6.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: At the surface to 1 foot below the surface

Ponding: At the surface to 0.5 foot above the surface

Flooding: None

Potential for frost action: High

Corrosivity: High for steel and low for concrete

Surface runoff class: Negligible

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Spaulding—2w

Prime farmland status: Spaulding—prime farmland where drained

Hydric soil status: Spaulding—hydric

Tama Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Typic Argiudolls

Typical Pedon

Tama silt loam, 2 to 5 percent slopes, at an elevation of 640 feet; Sangamon County, Illinois; about 1,600 feet south and 2,480 feet east of the northwest corner of sec. 34, T. 17 N., R. 3 W.; USGS Cornland topographical quadrangle; lat. 39 degrees 53 minutes 6 seconds N. and 89 degrees 24 minutes 55 seconds W., NAD 83:

Ap—0 to 11 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; moderate fine granular structure; friable; few fine roots throughout; neutral; clear smooth boundary.

BA—11 to 19 inches; brown (10YR 4/3) silt loam; moderate medium subangular blocky structure parting to moderate fine subangular blocky; friable; few fine roots throughout; many distinct very dark grayish brown (10YR 3/2) organic coatings on faces of peds; slightly acid; clear smooth boundary.

Bt1—19 to 30 inches; brown (10YR 4/3) silty clay loam; moderate fine prismatic structure parting to moderate fine subangular blocky; friable; few fine roots throughout; very few distinct very dark grayish brown (10YR 3/2) organic coatings along pores; many distinct dark brown (10YR 3/3) organo-clay films on faces of peds; slightly acid; gradual smooth boundary.

Bt2—30 to 39 inches; dark yellowish brown (10YR 4/4) silty clay loam; moderate medium prismatic structure parting to moderate medium subangular blocky; friable; common fine roots throughout; many distinct brown (10YR 4/3) clay films on faces of peds; slightly acid; gradual smooth boundary.

Bt3—39 to 58 inches; yellowish brown (10YR 5/4) silty clay loam; weak medium prismatic structure; friable; few fine roots throughout; common distinct brown (10YR 4/3) clay films on faces of peds and few distinct brown (10YR 5/3) silt coatings on faces of peds; moderately acid; clear smooth boundary.

BC—58 to 65 inches; yellowish brown (10YR 5/4) silt loam; weak coarse prismatic structure; friable; few fine roots throughout; common distinct brown (10YR 4/3) clay films and few distinct brown (10YR 5/3) silt coatings along pores; slightly acid; clear smooth boundary.

C1—65 to 77 inches; 55 percent yellowish brown (10YR 5/4) and 43 percent brown (10YR 5/3) silt loam; few fine distinct yellowish brown (10YR 5/6) mottles along pores; massive; very friable; very few distinct brown (10YR 4/3) clay films along pores; slightly acid; gradual smooth boundary.

C2—77 to 92 inches; 55 percent yellowish brown (10YR 5/4) and 43 percent brown (10YR 5/3) silt loam; common fine distinct yellowish brown (10YR 5/6) mottles along pores; massive; very friable; few fine distinct rounded black (10YR 2/1) accumulations of iron and manganese throughout; slightly acid.

Range in Characteristics

Thickness of mollic epipedon: 10 to 20 inches

Depth to carbonates: More than 48 inches

Depth to base of diagnostic horizon: 36 to more than 60 inches

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 or 2

Texture—silt loam

Bt horizon(s):

Hue—10YR

Value—4 or 5

Chroma—3 or 4

Texture—silty clay loam

C horizon(s):

Hue—10YR

Value—4 or 5

Chroma—3 to 6

Texture—silt loam

Taxadjunct Feature

Tama silt loam, 5 to 10 percent slopes, eroded, has a dark surface soil that is thinner than is definitive for the series. This difference, however, does not significantly affect the use and management of the soil. The soil is classified as a fine-silty, mixed, superactive, mesic Mollic Hapludalf.

36B—Tama silt loam, 2 to 5 percent slopes

Setting

Landform: Loessial hills

Position on landform: Summits and backslopes

Map Unit Composition

Tama and similar soils: 95 percent

Dissimilar soils: 5 percent

Minor Components

Similar soils:

- Soils that have a thinner dark surface soil
- Soils that have carbonates within a depth of 40 inches
- Soils that have a seasonal high water table at a depth of 4 to 6 feet
- Soils that have more sand in the lower part of the subsoil and in the underlying material

Dissimilar soils:

- The somewhat poorly drained Ipava soils on footslopes

Properties and Qualities of the Tama Soil

Parent material: Loess

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.1 inches to a depth of 60 inches

Content of organic matter in the surface layer: 3.0 to 4.0 percent

Shrink-swell potential: Moderate

Flooding: None

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Low

Susceptibility to water erosion: Moderate

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Tama—2e

Prime farmland status: Tama—prime farmland in all areas

Hydric soil status: Tama—not hydric

36C2—Tama silt loam, 5 to 10 percent slopes, eroded

Setting

Landform: Loessial hills

Position on landform: Shoulders and backslopes

Map Unit Composition

Tama and similar soils: 95 percent

Dissimilar soils: 5 percent

Minor Components

Similar soils:

- Soils that have carbonates within a depth of 40 inches
- Soils that have more clay in the surface layer
- Soils that have less clay in the subsoil
- Soils that have more sand in the lower part of the subsoil and in the underlying material

Dissimilar soils:

- The somewhat poorly drained Ipava soils on footslopes

Properties and Qualities of the Tama Soil

Parent material: Loess

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.7 inches to a depth of 60 inches

Content of organic matter in the surface layer: 2.0 to 3.0 percent

Shrink-swell potential: Moderate

Flooding: None

Accelerated erosion: The surface layer has been thinned by erosion.

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Medium

Susceptibility to water erosion: Moderate

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Tama—3e

Prime farmland status: Tama—not prime farmland

Hydric soil status: Tama—not hydric

Thebes Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Typic Hapludalfs

Typical Pedon (OSD)

Thebes silt loam, 5 to 10 percent slopes, at an elevation of 670 feet; Mercer County, Illinois; 1,060 feet west and 1,800 feet south of the northeast corner of sec. 3, T. 13 N., R. 3 W.; USGS Aledo East topographic quadrangle; lat. 41 degrees 09 minutes 02 seconds N. and long. 90 degrees 42 minutes 30 seconds W., NAD 27:

Ap—0 to 9 inches; dark grayish brown (10YR 4/2) silt loam, light brownish gray (10YR 6/2) dry; moderate medium granular structure; friable; slightly acid; clear smooth boundary.

Bt1—9 to 14 inches; yellowish brown (10YR 5/4) silty clay loam; weak fine and medium subangular blocky structure; friable; few distinct brown (10YR 5/3) clay films on faces of peds and in pores; strongly acid; clear wavy boundary.

Bt2—14 to 26 inches; dark yellowish brown (10YR 4/4) silty clay loam; moderate medium subangular blocky structure; friable; common distinct brown (10YR 4/3) clay films on faces of peds and in pores; moderately acid; clear wavy boundary.

Bt3—26 to 31 inches; dark yellowish brown (10YR 4/4) silty clay loam; moderate medium subangular blocky structure; friable; common distinct brown (10YR 4/3) clay films on faces of peds and in pores; few medium faint pale brown (10YR 6/3) and few medium distinct strong brown (7.5YR 4/6) accumulations of iron in the matrix; slightly acid; clear wavy boundary.

2Bt4—31 to 40 inches; dark yellowish brown (10YR 4/4) loam; moderate medium subangular blocky structure; friable; few distinct brown (10YR 4/3) clay films on faces of peds and in pores; common coarse faint pale brown (10YR 6/3) and common coarse distinct strong brown (7.5YR 4/6) accumulations of iron in the matrix; slightly acid; clear wavy boundary.

2BC—40 to 50 inches; 55 percent yellowish brown (10YR 5/4) and 43 percent brown (7.5YR 4/4), stratified sandy loam and loamy sand; weak medium subangular blocky structure; friable; few medium distinct pale brown (10YR 6/3) accumulations of iron in the matrix; moderately acid; clear wavy boundary.

2C—50 to 80 inches; dark yellowish brown (10YR 4/4), stratified loamy sand and sand; single grain; loose; common medium and coarse faint brown (7.5YR 4/4) accumulations of iron in the matrix; slightly acid.

Range in Characteristics

Thickness of the loess or other silty material: 20 to 40 inches

Depth to base of diagnostic horizon: 25 to 55 inches

Ap or A horizon(s):

Hue—10YR

Value—4 or 5

Chroma—2 to 4

Texture—silt loam or silty clay loam

E horizon(s), where present:

Hue—10YR

Value—4 or 5

Chroma—3 to 6

Texture—silt loam

Bt horizon(s):

Hue—10YR or 7.5YR

Value—4 or 5

Chroma—3 to 6

Texture—silty clay loam or silt loam

2Bt horizon(s):

Hue—10YR or 7.5YR

Value—4 or 5

Chroma—4 to 6

Texture—loam, sandy loam, fine sandy loam, sandy clay loam, or clay loam

2BC horizon(s), where present:

Hue—10YR or 7.5YR

Value—4 or 5

Chroma—4 to 6

Texture—sandy loam, loamy sand, or sand

2C horizon(s):

Hue—10YR or 7.5YR

Value—4 to 6

Chroma—3 to 6

Texture—loamy sand, fine sand, loamy fine sand, or sand; strata of sandy loam, fine sandy loam, or silt loam in the lower part of the profile in some pedons

212C2—Thebes silt loam, 5 to 10 percent slopes, eroded

Setting

Landform: Knolls and ground moraines

Position on landform: Summits and backslopes

Map Unit Composition

Thebes and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have more sand in the surface layer
- Soils that have less sand in lower part of the subsoil
- Soils that have more clay in the surface layer
- Soils that have more sand in the upper part of the subsoil

Dissimilar soils:

- The well drained Fayette soils on shoulders and backslopes

Properties and Qualities of the Thebes Soil

Parent material: Loess over sandy eolian material

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:

Moderate

Permeability below a depth of 60 inches: Rapid

Depth to restrictive feature: More than 80 inches

Available water capacity: About 9.5 inches to a depth of 60 inches

Content of organic matter in the surface layer: 1.0 to 2.0 percent

Shrink-swell potential: Moderate

Flooding: None

Accelerated erosion: The surface layer has been thinned by erosion.

Potential for frost action: High

Corrosivity: Moderate for steel and concrete

Surface runoff class: Medium

Susceptibility to water erosion: High

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Thebes—3e

Prime farmland status: Thebes—not prime farmland

Hydric soil status: Thebes—not hydric

Tice Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Fluvaquentic Hapludolls

Typical Pedon (OSD)

Tice silty clay loam, 0 to 2 percent slopes, occasionally flooded; at an elevation of about 465 feet; Adams County, Illinois; 1,670 feet north and 990 feet west of the southeast corner of sec. 22, T. 2 S., R. 9

W.; USGS Quincy West topographic quadrangle, lat. 39 degrees 52 minutes 56 seconds N. and long. 91 degrees 25 minutes 7 seconds W., NAD 27:

Ap—0 to 9 inches; very dark grayish brown (10YR 3/2) silty clay loam, grayish brown (10YR 5/2) dry; weak fine subangular blocky structure parting to weak medium granular; firm; common very fine roots throughout; neutral; abrupt smooth boundary.

A—9 to 14 inches; very dark grayish brown (10YR 3/2) silty clay loam, grayish brown (10YR 5/2) dry; moderate fine subangular blocky structure; firm; few very fine roots throughout; few fine faint brown (10YR 4/3) accumulations of iron in the matrix; neutral; clear smooth boundary.

BA—14 to 19 inches; dark grayish brown (10YR 4/2) silty clay loam; weak fine prismatic structure parting to moderate medium subangular blocky; firm; few very fine roots throughout; common distinct very dark grayish brown (10YR 3/2) organic coatings on faces of peds; common fine faint brown (7.5YR 4/3) accumulations of iron in the matrix; few fine faint grayish brown (10YR 5/2) iron depletions in the matrix; neutral; clear smooth boundary.

Bw—19 to 35 inches; brown (10YR 4/3) silty clay loam; weak medium prismatic structure parting to moderate medium subangular blocky; firm; few very fine roots throughout; common distinct very dark grayish brown (10YR 3/2) organo-clay films on faces of peds; many medium prominent strong brown (7.5YR 4/6) accumulations of iron in the matrix; few fine faint grayish brown (10YR 5/2) iron depletions in the matrix; moderately acid; clear smooth boundary.

Bg1—35 to 44 inches; dark grayish brown (10YR 4/2) silty clay loam; weak medium prismatic structure parting to moderate medium subangular blocky; firm; few very fine roots throughout; common distinct very dark gray (10YR 3/1) organic coatings on faces of peds; many medium prominent strong brown (7.5YR 4/6) accumulations of iron in the matrix; moderately acid; gradual smooth boundary.

Bg2—44 to 61 inches; dark grayish brown (2.5Y 4/2) silty clay loam; weak medium prismatic structure; firm; common distinct very dark gray (10YR 3/1) organic coatings on faces of peds; common medium prominent strong brown (7.5YR 4/6) accumulations of iron in the matrix; slightly acid; clear smooth boundary.

Bg3—61 to 80 inches; dark grayish brown (2.5Y 4/2) silty clay loam; weak medium prismatic structure;

firm; common distinct very dark gray (10YR 3/1) organic coatings on faces of peds; common medium prominent strong brown (7.5YR 4/6) accumulations of iron in the matrix; slightly acid.

Range in Characteristics

Thickness of the mollic epipedon: 10 to 24 inches
Depth to base of diagnostic horizon: 30 to more than 80 inches

Profile feature: An AB horizon in some pedons

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 or 2

Texture—silty clay loam or silt loam

Bw or Bg horizon(s):

Hue—10YR or 2.5Y

Value—4 or 5

Chroma—2 to 4

Texture—silty clay loam or silt loam

BC or BCg horizon(s), where present:

Hue—10YR, 2.5Y, or 5Y

Value—4 or 5

Chroma—1 to 4

Texture—silty clay loam or silt loam; strata of loam, clay loam, or sandy loam in some pedons

Cg or C horizon(s), where present:

Hue—10YR, 2.5Y, or 5Y

Value—4 to 6

Chroma—1 to 3

Texture—silty clay loam, clay loam, loam, sandy loam, or silt loam

3284A—Tice silty clay loam, 0 to 2 percent slopes, frequently flooded

Setting

Landform: Flood plains (fig. 3)

Map Unit Composition

Tice and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a thicker dark surface soil
- Soils that have more sand in the upper part of the subsoil
- Soils that have less clay in the surface layer
- Soils that have a buried soil

Dissimilar soils:

- The poorly drained Sawmill soils on flood plains
- The well drained Ross soils on flood plains

Properties and Qualities of the Tice Soil

Parent material: Alluvium

Drainage class: Somewhat poorly drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 12.8 inches to a depth of 60 inches

Content of organic matter in the surface layer: 2.0 to 4.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: 1 to 2 feet below the surface

Potential for frost action: High

Corrosivity: High for steel and low for concrete

Surface runoff class: Low

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Very low

Interpretive Groups

Land capability classification: Tice—3w

Prime farmland status: Tice—prime farmland where protected from flooding or not frequently flooded during the growing season

Hydric soil status: Tice—not hydric

533—Urban land**Setting**

This map unit consists of areas covered by parking lots, streets, buildings, and other structures.

Map Unit Composition

Urban land: 85 percent

Dissimilar components: 15 percent

Minor Components

Dissimilar components:

- The well drained, silty Orthents in open areas

Interpretive Groups

Land capability classification: Urban land—none assigned

Prime farmland status: Urban land—not prime farmland

Hydric soil status: Urban land—unranked

Vesser Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Argiaquic Argialbolls

Typical Pedon

Vesser silt loam, 0 to 2 percent slopes, occasionally flooded, at an elevation of 480 feet; Adams County, Illinois; 360 feet west and 220 feet south of the northeast corner of sec. 4, T. 1 N., R. 9 W.; USGS Long Island topographic quadrangle; lat. 40 degrees 6 minutes 37 seconds N. and long. 91 degrees 26 minutes 15 seconds W., NAD 27:

Ap—0 to 8 inches; very dark gray (10YR 3/1) silt loam, gray (10YR 5/1) dry; weak fine granular structure; friable; neutral; abrupt smooth boundary.

A—8 to 14 inches; very dark gray (10YR 3/1) silt loam, gray (10YR 5/1) dry; weak fine subangular blocky structure parting to weak medium granular; friable; common fine prominent dark brown (7.5YR 3/4) accumulations of iron throughout; neutral; gradual smooth boundary.

Eg1—14 to 20 inches; dark gray (10YR 4/1) silt loam, gray (10YR 6/1) dry; weak medium platy structure parting to weak very fine subangular blocky; friable; very few distinct very dark gray (10YR 3/1) organic coatings on faces of peds; common medium faint gray (10YR 5/1) clay depletions between peds and common fine prominent dark brown (7.5YR 3/4) accumulations of iron throughout; slightly acid; clear smooth boundary.

Eg2—20 to 26 inches; dark gray (10YR 4/1) silt loam, gray (10YR 6/1) dry; weak thick platy structure parting to weak very fine subangular blocky; friable; very few distinct very dark gray (10YR 3/1) organic coatings on faces of peds; common medium faint gray (10YR 6/1) clay depletions between peds and common fine prominent brown (7.5YR 4/4) accumulations of iron throughout; slightly acid; gradual smooth boundary.

Btg1—26 to 34 inches; gray (10YR 5/1) silty clay loam; weak medium prismatic structure; friable; very few distinct very dark gray (10YR 3/1) organo-clay films on faces of peds and few distinct gray (10YR 6/1) silt coatings in root channels and/or pores; common medium prominent dark brown (7.5YR 3/4) accumulations of iron throughout; moderately acid; gradual smooth boundary.

Btg2—34 to 48 inches; gray (10YR 5/1) silty clay loam; weak medium prismatic structure; firm; very few distinct very dark gray (10YR 3/1) organo-clay films on faces of peds and very few distinct light

brownish gray (10YR 6/2) silt coatings in root channels and/or pores; common medium prominent dark brown (7.5YR 3/4) accumulations of iron throughout; moderately acid; gradual smooth boundary.

Btg3—48 to 58 inches; gray (10YR 5/1) silty clay loam; weak medium prismatic structure; firm; few distinct light brownish gray (10YR 6/2) silt coatings in root channels and/or pores and very few distinct very dark gray (10YR 3/1) organo-clay films on faces of peds; common medium prominent dark brown (7.5YR 3/4) accumulations of iron throughout; slightly acid; clear smooth boundary.

BCg—58 to 80 inches; gray (10YR 5/1) silty clay loam; weak coarse prismatic structure; firm; very few distinct dark gray (10YR 4/1) clay films on faces of peds and very few distinct light brownish gray (10YR 6/2) silt coatings in root channels and/or pores; common medium prominent dark brown (7.5YR 3/4) accumulations of iron throughout; slightly acid.

Range in Characteristics

Thickness of the mollic epipedon: 10 to 20 inches

Depth to base of diagnostic horizon: More than 60 inches

Ap or A horizon(s):

Hue—10YR

Value—2 or 3

Chroma—1 or 2

Texture—silt loam

E or Eg horizon(s):

Hue—10YR

Value—3 to 5

Chroma—1 or 2

Texture—silt loam

Btg horizon(s):

Hue—10YR or 2.5Y

Value—3 to 5

Chroma—1 or 2

Texture—silty clay loam

8396A—Vesser silt loam, 0 to 2 percent slopes, occasionally flooded

Setting

Landform: Flood plains

Map Unit Composition

Vesser and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have more clay in the subsoil
- Soils that have a thinner dark surface layer
- Soils that have a lighter colored surface layer
- Soils that have more sand in the lower part of the subsoil

Dissimilar soils:

- The well drained Huntsville soils on flood plains
- The well drained Proctor soils on flood-plain steps

Properties and Qualities of the Vesser Soil

Parent material: Alluvium

Drainage class: Poorly drained

Slowest permeability within a depth of 40 inches:

Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.9 inches to a depth of 60 inches

Content of organic matter in the surface layer: 2.0 to 3.0 percent

Shrink-swell potential: Moderate

Apparent seasonal high water table: At the surface to 1 foot below the surface

Potential for frost action: High

Corrosivity: High for steel and moderate for concrete

Surface runoff class: Negligible

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Vesser—2w

Prime farmland status: Vesser—prime farmland where drained

Hydric soil status: Vesser—hydric

Viriden Series

Taxonomic classification: Fine, smectitic, mesic Vertic Argiaquolls

Typical Pedon (OSD)

Viriden silty clay loam, 0 to 2 percent slopes, at an elevation of 699 feet; Adams County, Illinois; 140 feet west and 54 feet north of the southeast corner of sec. 3, T. 2 N., R. 6 W.; USGS Bowen topographic quadrangle; lat. 40 degrees 10 minutes 52 seconds N. and long. 91 degrees 4 minutes 5 seconds W., NAD 27:

Ap—0 to 8 inches; black (10YR 2/1) silty clay loam, dark gray (10YR 4/1) dry; moderate medium

granular structure; firm; slightly alkaline; abrupt smooth boundary.

A—8 to 16 inches; black (10YR 2/1) silty clay loam, dark gray (10YR 4/1) dry; moderate fine granular structure; firm; moderately acid; clear smooth boundary.

Btg1—16 to 23 inches; very dark gray (10YR 3/1) silty clay, grayish brown (10YR 5/2) dry; strong fine angular blocky structure; firm; few faint black (10YR 2/1) organo-clay films on faces of peds; few fine faint black (10YR 2/1) concretions of iron and manganese throughout; slightly acid; clear smooth boundary.

Btg2—23 to 34 inches; gray (5Y 5/1) silty clay loam; weak coarse prismatic structure parting to moderate medium angular blocky; firm; few distinct dark gray (10YR 4/1) clay films on faces of peds; many medium prominent brownish yellow (10YR 6/6) accumulations of iron and few fine prominent black (10YR 2/1) accumulations of iron and manganese throughout; slightly acid; clear smooth boundary.

Btg3—34 to 42 inches; gray (5Y 5/1) silty clay loam; weak and moderate coarse prismatic structure parting to moderate coarse angular blocky; firm; few distinct dark gray (5Y 4/1) clay films on faces of peds; common medium prominent light olive brown (2.5Y 5/6) accumulations of iron and few fine prominent black (10YR 2/1) accumulations of iron and manganese throughout; neutral; clear smooth boundary.

Btg4—42 to 49 inches; gray (5Y 5/1) silty clay loam; moderate coarse prismatic structure parting to weak coarse angular blocky; firm; very few distinct dark gray (N 4/0) clay films on faces of peds; many medium prominent olive brown (2.5Y 4/4) accumulations of iron throughout; neutral; gradual smooth boundary.

Cg—49 to 60 inches; gray (5Y 5/1) silty clay loam; massive; firm; common medium prominent olive brown (2.5Y 4/4) accumulations of iron throughout; neutral.

Range in Characteristics

Thickness of the mollic epipedon: 10 to 24 inches

Depth to carbonates, where present: More than 50 inches

Depth to base of diagnostic horizon: 40 to 60 inches

Ap or A horizon(s):

Hue—10YR, 2.5Y, or 5Y

Value—2 or 3

Chroma—1 or 2

Texture—silty clay loam or silt loam

Btg horizon(s):

Hue—10YR, 2.5Y, 5Y, or neutral

Value—2 to 6

Chroma—0 to 4

Texture—silty clay loam, silty clay, or silt loam

Cg horizon(s):

Hue—10YR, 2.5Y, 5Y, or neutral

Value—4 to 6

Chroma—0 to 4

Texture—silty clay loam or silt loam

50A—Virden silty clay loam, 0 to 2 percent slopes

Setting

Landform: Ground moraines

Position on landform: Toeslopes

Map Unit Composition

Virden and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils in which the dark surface soil is more than 24 inches thick
- Soils that have less clay in the subsoil
- Soils that have less clay in the surface layer
- Soils that have carbonates within a depth of 50 inches

Dissimilar soils:

- The moderately well drained Buckhart and Harrison soils on summits and backslopes

Properties and Qualities of the Virden Soil

Parent material: Loess

Drainage class: Poorly drained

Slowest permeability within a depth of 40 inches:

Moderately slow

Permeability below a depth of 60 inches: Moderately slow

Depth to restrictive feature: More than 80 inches

Available water capacity: About 11.1 inches to a depth of 60 inches

Content of organic matter in the surface layer: 3.0 to 6.0 percent

Shrink-swell potential: High

Apparent seasonal high water table: At the surface to 1 foot below the surface

Ponding: At the surface to 0.5 foot above the surface

Flooding: None

Potential for frost action: High
Corrosivity: High for steel and moderate for concrete
Surface runoff class: Negligible
Susceptibility to water erosion: Low
Susceptibility to wind erosion: Very low

Interpretive Groups

Land capability classification: Virden—2w
Prime farmland status: Virden—prime farmland where drained
Hydric soil status: Virden—hydric

Worthen Series

Taxonomic classification: Fine-silty, mixed, superactive, mesic Cumulic Hapludolls

Typical Pedon (OSD)

Worthen silt loam, 2 to 5 percent slopes, rarely flooded, at an elevation of 465 feet; Scott County, Illinois; 160 feet south and 640 feet west of the northeast corner of sec. 26, T. 13 N., R. 13 W.; USGS Bedford topographic quadrangle; lat. 39 degrees 33 minutes 00 seconds N. and long. 90 degrees 30 minutes 33 seconds W., NAD 27:

Ap—0 to 9 inches; very dark grayish brown (10YR 3/2) silt loam, grayish brown (10YR 5/2) dry; weak fine granular structure; friable; common very fine and fine roots; neutral; abrupt smooth boundary.

A—9 to 20 inches; dark brown (10YR 3/3) silt loam, brown (10YR 5/3) dry; weak medium granular structure; friable; few very fine and fine roots; common distinct very dark grayish brown (10YR 3/2) organic coatings on faces of peds; slightly acid; clear smooth boundary.

AB—20 to 29 inches; dark brown (10YR 3/3) silt loam, brown (10YR 5/3) dry; weak fine subangular blocky structure; friable; few very fine and fine roots; common distinct very dark grayish brown (10YR 3/2) organic coatings on faces of peds; neutral; clear smooth boundary.

Bw1—29 to 41 inches; brown (10YR 4/3) silt loam; weak fine subangular blocky structure; friable; few very fine and fine roots; common distinct dark brown (10YR 3/3) organic coatings on faces of peds, few distinct very dark grayish brown (10YR 3/2) organic coatings in root channels and/or pores, and few distinct very pale brown (10YR 7/3) silt coatings on faces of peds; neutral; clear smooth boundary.

Bw2—41 to 64 inches; dark yellowish brown (10YR 4/4) silt loam; weak fine subangular blocky

structure; friable; few very fine and fine roots; few distinct dark brown (10YR 3/3) organic coatings in root channels and/or pores and few distinct very pale brown (10YR 7/3) silt coatings on faces of peds; neutral; gradual smooth boundary.

C—64 to 80 inches; yellowish brown (10YR 5/4) silt loam; weak medium subangular blocky structure; friable; neutral.

Range in Characteristics

Thickness of the mollic epipedon: 24 to 36 inches

Depth to carbonates, where present: More than 50 inches

Depth to base of diagnostic horizon: 30 to 80 inches

Ap, A, or AB horizon:

Hue—10YR

Value—2 or 3

Chroma—1 to 3

Texture—silt loam

Bw horizon:

Hue—10YR or 7.5YR

Value—3 to 5

Chroma—2 to 6

Texture—silt loam

C horizon:

Hue—10YR or 7.5YR

Value—4 or 5

Chroma—3 to 6

Texture—silt loam

7037A—Worthen silt loam, 0 to 2 percent slopes, rarely flooded

Setting

Landform: Alluvial fans

Map Unit Composition

Worthen and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a thinner dark surface soil
- Soils that have more clay in the subsoil
- Soils that have more sand in the lower part of the subsoil and in the underlying material

Dissimilar soils:

- The somewhat poorly drained Kendall soils on flood-plain steps
- The poorly drained Vesser soils on flood plains

Properties and Qualities of the Worthen Soil

Parent material: Silty local alluvium

Drainage class: Well drained

Slowest permeability within a depth of 40 inches:
Moderate

Permeability below a depth of 60 inches: Moderate

Depth to restrictive feature: More than 80 inches

Available water capacity: About 13.1 inches to a depth of 60 inches

Content of organic matter in the surface layer: 2.0 to 4.0 percent

Shrink-swell potential: Low

Potential for frost action: High

Corrosivity: Low for steel and concrete

Surface runoff class: Low

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Low

Interpretive Groups

Land capability classification: Worthen—1

Prime farmland status: Worthen—prime farmland in all areas

Hydric soil status: Worthen—not hydric

Zook Series

Taxonomic classification: Fine, smectitic, mesic
Cumulic Vertic Endoaquolls

Typical Pedon

Zook silty clay loam, 0 to 2 percent slopes, frequently flooded; at an elevation of 600 feet; Warren County, Illinois; 2,640 feet west and 1,200 feet south of the northeast corner of sec. 22, T. 12 N., R. 3 W.; USGS Little York topographic quadrangle; lat. 41 degrees 01 minute 14 seconds N. and long. 90 degrees 43 minutes 03 seconds W., NAD 27:

Ap—0 to 8 inches; black (10YR 2/1) silty clay loam, dark gray (10YR 4/1) dry; moderate fine granular structure; friable; many fine roots; slightly acid; clear smooth boundary.

A—8 to 22 inches; very dark gray (N 3/0) silty clay, gray (N 5/0) dry; moderate fine subangular blocky structure; firm; common medium and many fine roots; slightly acid; clear smooth boundary.

Bg1—22 to 38 inches; very dark gray (10YR 3/1) silty clay, gray (10YR 5/1) dry; weak fine prismatic structure parting to moderate fine subangular and angular blocky; firm; common fine and medium and few coarse roots; few fine prominent strong brown (7.5YR 5/6) accumulations of iron; neutral; clear wavy boundary.

Bg2—38 to 55 inches; dark gray (5Y 4/1) silty clay;

weak medium prismatic structure parting to moderate fine subangular and angular blocky; firm; common fine and medium and few coarse roots; common faint very dark gray (5Y 3/1) organic coatings in root channels and krotovinas; common medium faint olive gray (5Y 5/2) iron depletions and common medium prominent strong brown (7.5YR 5/6) accumulations of iron; neutral; clear wavy boundary.

BCg—55 to 60 inches; olive gray (5Y 5/2) silty clay loam; weak fine and medium subangular blocky structure; firm; few medium and coarse and common fine roots; common faint gray (5Y 5/1) organic coatings in root channels and on faces of peds; common medium prominent strong brown (7.5YR 5/6) accumulations of iron; neutral.

Range in Characteristics

Thickness of the mollic epipedon: 36 to 60 inches

Ap or A horizon(s):

Hue—10YR or neutral

Value—2 or 3

Chroma—0 or 1

Texture—silty clay loam or silty clay

Bg or Cg horizon(s), where present:

Hue—10YR, 2.5Y, or 5Y

Value—2 to 5

Chroma—1 or 2

Texture—silty clay, silty clay loam, or silt loam

3405A—Zook silty clay loam, 0 to 2 percent slopes, frequently flooded

Setting

Landform: Flood plains

Map Unit Composition

Zook and similar soils: 90 percent

Dissimilar soils: 10 percent

Minor Components

Similar soils:

- Soils that have a thinner dark surface soil
- Soils that have less clay in the subsoil
- Soils that have more sand in the underlying material
- Soils that have less clay in the surface layer

Dissimilar soils:

- The well drained Huntsville soils on flood plains

Properties and Qualities of the Zook Soil

Parent material: Alluvium

Drainage class: Poorly drained

Slowest permeability within a depth of 40 inches: Slow

Permeability below a depth of 60 inches: Slow or moderately slow

Depth to restrictive feature: More than 80 inches

Available water capacity: About 8.2 inches to a depth of 60 inches

Content of organic matter in the surface layer: 4.0 to 5.0 percent

Shrink-swell potential: High

Apparent seasonal high water table: At the surface to 1 foot below the surface

Potential for frost action: High

Corrosivity: High for steel and moderate for concrete

Surface runoff class: Negligible

Susceptibility to water erosion: Low

Susceptibility to wind erosion: Very low

Interpretive Groups

Land capability classification: Zook—3w

Prime farmland status: Zook—prime farmland where drained and either protected from flooding or not frequently flooded during the growing season

Hydric soil status: Zook—hydric

Use and Management of the Soils

This soil survey is an inventory and evaluation of the soils in the survey area. It can be used to adjust land uses to the limitations and potentials of natural resources and the environment. Also, it can help to prevent soil-related failures in land uses.

In preparing a soil survey, soil scientists, conservationists, engineers, and others collect extensive field data about the nature and behavioral characteristics of the soils. They collect data on erosion, droughtiness, flooding, and other factors that affect various soil uses and management. Field experience and collected data on soil properties and performance are used as a basis in predicting soil behavior.

Information in this section can be used to plan the use and management of soils for crops and pasture; as forestland; as sites for buildings, sanitary facilities, highways and other transportation systems, and parks and other recreational facilities; for agricultural waste management; and as wildlife habitat. It can be used to identify the potentials and limitations of each soil for specific land uses and to help prevent construction failures caused by unfavorable soil properties.

Planners and others using soil survey information can evaluate the effect of specific land uses on productivity and on the environment in all or part of the survey area. The survey can help planners to maintain or create a land use pattern in harmony with the natural soil.

Contractors can use this survey to locate sources of sand and gravel, roadfill, and topsoil. They can use it to identify areas where bedrock, wetness, or very firm soil layers can cause difficulty in excavation.

Health officials, highway officials, engineers, and others may also find this survey useful. The survey can help them plan the safe disposal of wastes and locate sites for pavements, sidewalks, campgrounds, playgrounds, lawns, and trees and shrubs.

Interpretive Ratings

The interpretive tables in this survey rate the soils in the survey area for various uses. Many of the tables identify the limitations that affect specified uses and

indicate the severity of those limitations. The ratings in these tables are both verbal and numerical.

Rating Class Terms

Rating classes are expressed in the tables in terms that indicate the extent to which the soils are limited by all of the soil features that affect a specified use or in terms that indicate the suitability of the soils for the use. Thus, the tables may show limitation classes or suitability classes. Terms for the limitation classes are *not limited*, *somewhat limited*, and *very limited*. The suitability ratings are expressed as *well suited*, *moderately suited*, *poorly suited*, and *unsuited* or as *good*, *fair*, and *poor*.

Numerical Ratings

Numerical ratings in the tables indicate the relative severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use and the point at which the soil feature is not a limitation. The limitations appear in order from the most limiting to the least limiting. Thus, if more than one limitation is identified, the most severe limitation is listed first and the least severe one is listed last.

Crops and Pasture

General management needed for crops and pasture is suggested in this section. The estimated yields of the main crops and pasture plants are listed, the system of land capability classification used by the Natural Resources Conservation Service is explained, and prime farmland is described.

Planners of management systems for individual fields or farms should consider the detailed information given in the description of each soil under the heading "Detailed Soil Map Units." Specific information can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

A total of 436,440 acres in Sangamon County is cropland (USDA, 1997). The major row crops are corn and soybeans. The major small grain crop is wheat.

The soils in Sangamon County have good potential for continued crop production, especially if the latest crop-production technology is applied. This soil survey can be used as a guide in applying this technology.

Management Considerations on Cropland

The management concerns affecting the use of the soils in Sangamon County for crops and pasture are shown in the table 6. The main concerns in managing cropland are crusting, flooding, poor tilth, water erosion, and wetness. Excess lime, excessive permeability, high pH, and ponding are additional management concerns.

Crusting occurs when flowing water or raindrops break down soil structural units, moving clay downward and leaving a concentration of sand grains and silt particles on the surface. Crusts can reduce the rate of water infiltration, increase the runoff rate, and restrict seedling emergence and oxygen diffusion to seedlings.

Crusting can be minimized by increasing soil aggregate stability through the addition of organic matter to the surface and by maintaining a cover of plants or crop residue, which reduces the impact of raindrops.

Flooding occurs in unprotected areas along the major rivers and their tributaries. Levees or diversions reduce the extent of the crop damage caused by floodwater. Surface drainage ditches help to remove floodwater where suitable outlets are available. Management of drainage systems in conformance with regulations influencing wetlands may require special permits and extra planning. Selecting crop varieties that are adapted to shorter growing seasons and wetter conditions reduces the extent of flood damage.

Poor tilth can occur in soils when part of the subsoil is incorporated into the plow layer because of erosion. Incorporation of subsoil material decreases the amount of organic matter and increases the content of clay in the surface soil. Intensive rainfall often causes surface crusting. Poor tilth also occurs in poorly drained soils with a high content of clay, regardless of the content of organic matter, and in soils that have been excessively tilled.

Poor tilth decreases the rate of water infiltration and increases the runoff rate and the susceptibility to erosion on the more sloping soils. Soils with poor tilth generally have a surface layer that is sticky when wet

and hard and cloddy when dry. They can be tilled only within a narrow range of moisture content. As a result, seedbed preparation is difficult.

Returning crop residue to the soil, regularly adding other organic material, minimizing tillage, and applying conservation tillage systems during periods of near optimal soil moisture conditions improve tilth.

Water erosion can occur when the surface soil is not protected against the impact of raindrops, which can reduce the stability of soil aggregates. This reduced stability decreases the rate of water infiltration and increases the surface runoff rate. Soils with long or steep slopes are more susceptible to water erosion than other soils.

Erosion, primarily sheet and rill erosion, removes the surface soil, which commonly has more biological activity and organic matter than any other part of the soil. Soil productivity decreases as the content of organic matter and level of natural fertility are lowered. Poor tilth and crusting occur as the subsoil, which is generally higher in content of clay than the surface soil, is incorporated through tillage into the plow layer.

Excessive runoff decreases the quality of surface water through sedimentation and contamination by pesticides.

Erosion can be controlled by a conservation tillage system that leaves crop residue on the surface after planting or by a cropping system that includes grasses and legumes in the cropping sequence. Contour farming and/or terraces in combination with a conservation tillage system can help to control erosion on soils with long, uniform slopes.

Wetness occurs in soils when the seasonal high water table is at or near the surface. Subsurface tile drains can lower the seasonal high water table if suitable outlets are available. In soils with a high content of clay and restricted permeability, a subsurface drainage system may not be practical. In these soils surface ditches can reduce the wetness. Management of drainage in conformance with regulations influencing wetlands may require special permits and extra planning.

Excess lime occurs in soils that contain a high content of calcium carbonate at or near the surface. The lime affects the availability of many plant nutrients and influences the effectiveness of herbicides. Frequent applications of a small amount of fertilizer are needed to correct nutrient imbalances. Crops may respond well to additions of phosphate fertilizer on these soils. Applications of herbicide should be adjusted as the level of alkalinity increases. Incorporating green manure crops, manure, or crop residue into the soil, applying a system of

conservation tillage, and using conservation cropping systems also help to overcome this limitation.

Excessive permeability can occur in soils that have a high content of sand and many of the larger diameter pores. The capacity of the soils to retain moisture for plant use is limited. Deep leaching of nutrients and pesticides is possible. It increases the risk of ground-water pollution.

Irrigation can supply the moisture needed for crops. Frequent applications of a small amount of fertilizer are needed. One application of a large amount can result in excessive loss of plant nutrients through leaching.

High pH, or a pH of more than 7.9, affects the availability of many plant nutrients and influences the effectiveness of herbicides. Frequent applications of a small amount of fertilizer are needed to correct nutrient imbalances. Crops may respond well to additions of phosphate fertilizer on the soils with a high pH. Applications of herbicide should be adjusted as the level of alkalinity increases. Incorporating green manure crops, manure, or crop residue into the soil, applying a system of conservation tillage, and using conservation cropping systems help to overcome this limitation.

Ponding occurs on soils when the seasonal high water table is above the surface. Land grading helps to control ponding. Surface ditches and surface inlet tile also can help to remove the excess water if suitable outlets are available. Management of drainage in conformance with regulations influencing wetlands may require special permits and extra planning.

Explanation of Criteria

Crusting.—In the surface layer, the average content of organic matter is 2.5 percent or less and the content of clay is between 20 and 35 percent.

Excess lime.—The calcium carbonate equivalent is 15 percent or more within 16 inches of the surface.

Excessive permeability.—The lower limit of the permeability rate is 6.0 or more inches per hour within the soil profile.

Flooding.—The soil is occasionally flooded or frequently flooded.

High pH.—The pH is more than 7.9 within a depth of 40 inches.

Ponding.—The seasonal high water table is above the surface.

Poor tilth.—The content of clay in the surface layer is 27 percent or more.

Water erosion.—The Kw factor multiplied by the slope is more than 0.8, and the slope is 3 percent or more.

Wetness.—The water table is within a depth of 1.5 feet at some time during the growing season in normal years.

Management Considerations on Pasture

The management concerns affecting the use of the soils in the county for pasture are shown in the table 6. The main management concerns are frost heave, low pH, and water erosion. Additional management concerns are an equipment limitation, flooding, high pH, low fertility, ponding, and wetness.

Frost heave occurs when ice lenses or bands that drive an ice wedge between two layers develop near the surface layer of a soil. The ice wedges heave the overlying soil layer upward, snapping the roots. Soils with a low content of sand have small pores that hold water and enable ice lenses to form. Selecting adapted forage and hay varieties helps to reduce the effects of frost heave. Timely deferment of grazing helps to maintain a protective cover that insulates the soil, thereby reducing the effects of frost heave.

Low pH, or a pH of 5.5 or less, can decrease the solubility and availability of plant nutrients. Selecting adapted forage and hay varieties and applying lime according to the results of soil tests help to overcome this limitation.

Water erosion can occur in overgrazed areas or during periods of pasture establishment and renovation when the surface soil is not protected against the impact of raindrops, which can cause poor tilth. Deterioration of tilth decreases the rate of water infiltration and increases the surface runoff rate. Soils with long or steep slopes are more susceptible to water erosion than other soils.

Erosion can be controlled by deferred grazing, which helps to prevent overgrazing and thus also helps to prevent surface compaction and excessive runoff and erosion. Tilling on the contour, using a no-till system of seeding when a seedbed is prepared or the pasture is renovated, and selecting adapted forage and hay varieties also help to control erosion.

An *equipment limitation* occurs on soils with slopes of more than 18 percent. This limitation can cause rapid wear of equipment. It can also hinder fertilization, harvest, pasture renovation, and seedbed preparation. It cannot be easily overcome.

Flooding occurs in unprotected areas along the major rivers and their tributaries. Surface drainage ditches help to remove floodwater where suitable outlets are available. Management of drainage in conformance with regulations may require special permits and extra planning. Selecting forage and hay

varieties adapted to shorter growing seasons and wetter conditions reduces the extent of flood damage. Restricted use during wet periods helps to keep the pasture in good condition.

High pH.—A pH of more than 7.9, affects the availability of many plant nutrients. Frequent applications of a small amount of fertilizer are needed to correct nutrient imbalances. Selecting adapted forage and hay varieties helps to overcome this limitation.

Low fertility occurs in soils with a low content of organic matter and a low cation-exchange capacity. The capacity of the soil to retain nutrients for plant use is limited. Frequent applications of small amounts of fertilizer help to prevent excessive loss of plant nutrients through leaching. When used as part of a seeding mixture, legumes can provide nitrogen to the grass varieties. Timely deferment of grazing helps to maintain the surface cover and the content of organic matter, a source of nutrients in the soil.

Ponding occurs on soils when the seasonal high water table is above the surface. Land grading helps to control ponding. Surface ditches and surface inlet tile also can help to remove the excess water if suitable outlets are available. Management of drainage in conformance with regulations may require special permits and extra planning. Selecting forage and hay varieties adapted to wet conditions improves forage production. Restricted use during wet periods helps to keep the pasture in good condition.

Wetness occurs in soils when the seasonal high water table is at or near the surface. Subsurface tile drains can help to lower the seasonal high water table if suitable outlets are available. Management of drainage in conformance with regulations may require special permits and extra planning. Selecting forage and hay varieties adapted to wet conditions improves forage production. Restricted use during wet periods helps to keep the pasture in good condition.

Explanation of Criteria

Equipment limitation.—The slope is more than 18 percent.

Flooding.—The soil is occasionally flooded or frequently flooded.

Frost heave.—The potential for frost action is moderate or high, and the soil is poorly drained or very poorly drained.

High pH.—The pH is more than 7.9 within a depth of 40 inches.

Low fertility.—The average content of organic matter in the surface layer is less than 1 percent, or the cation-exchange capacity, expressed in terms of milliequivalents per 100 grams of soil, is 7 or less.

Low pH.—The pH is 5.5 or less within a depth of 40 inches.

Ponding.—The seasonal high water table is above the surface.

Water erosion.—The Kw factor multiplied by the slope is more than 1.0, and the slope is 3 percent or more.

Wetness.—The seasonal high water table is within a depth of 1.5 feet.

Yields per Acre

The average yields per acre that can be expected of the principal crops under a high level of management are shown in table 7. In any given year, yields may be higher or lower than those indicated in the table because of variations in rainfall and other climatic factors. The land capability classification of map units in the survey area also is shown in the table.

The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby counties and results of field trials and demonstrations also are considered (Fehrenbacher et al., 1978).

The management needed to obtain the indicated yields of the various crops depends on the kind of soil and the crop. Management can include drainage, erosion control, and protection from flooding; the proper planting and seeding rates; suitable high-yielding crop varieties; appropriate and timely tillage; control of weeds, plant diseases, and harmful insects; favorable soil reaction and optimum levels of nitrogen, phosphorus, potassium, and trace elements for each crop; effective use of crop residue, barnyard manure, and green manure crops; and harvesting that ensures the smallest possible loss.

The estimated yields reflect the productive capacity of each soil for each of the principal crops. Yields are likely to increase as new production technology is developed. The productivity of a given soil compared with that of other soils, however, is not likely to change.

Crops other than those shown in table 7 are grown in the survey area, but estimated yields are not listed because the acreage of such crops is small. The local office of the Natural Resources Conservation Service or of the Cooperative Extension Service can provide information about the management and productivity of the soils for those crops.

Pasture and Hay Yields

Under good management, proper grazing is essential for the production of high-quality forage,

stand survival, and erosion control. Proper grazing helps the plants to maintain sufficient and generally vigorous top growth during the growing season. Brush control is essential in many areas, and weed control generally is needed. Rotation grazing and renovation also are important management practices.

Yield estimates are often given in animal unit months (AUMs), or the amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

The local office of the Natural Resources Conservation Service or of the Cooperative Extension Service can provide information about forage yields other than those shown in table 7.

Land Capability Classification

Land capability classification shows, in a general way, the suitability of soils for most kinds of field crops, including corn, small grain, and hay. Crops that require special management are excluded. The soils are grouped according to their limitations for field crops, the risk of damage if they are used for crops, and the way they respond to management. The criteria used in grouping the soils do not include major and generally expensive landforming that would change slope, depth, or other characteristics of the soils, nor do they include possible but unlikely major reclamation projects. Capability classification is not a substitute for interpretations designed to show suitability and limitations of groups of soils for woodland or for engineering purposes.

In the capability system, soils are generally grouped at three levels—capability class, subclass, and unit (USDA, 1961). Only class and subclass are used in this survey.

Capability classes, the broadest groups, are designated by the numbers 1 through 8. The numbers indicate progressively greater limitations and narrower choices for practical use. The classes are defined as follows:

Class 1 soils have slight limitations that restrict their use.

Class 2 soils have moderate limitations that restrict the choice of plants or that require moderate conservation practices.

Class 3 soils have severe limitations that restrict the choice of plants or that require special conservation practices, or both.

Class 4 soils have very severe limitations that restrict the choice of plants or that require very careful management, or both.

Class 5 soils are subject to little or no erosion but have other limitations, impractical to remove, that

restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

Class 6 soils have severe limitations that make them generally unsuitable for cultivation and that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

Class 7 soils have very severe limitations that make them unsuitable for cultivation and that restrict their use mainly to grazing, forestland, or wildlife habitat.

Class 8 soils and miscellaneous areas have limitations that preclude commercial plant production and that restrict their use to recreational purposes, wildlife habitat, watershed, or esthetic purposes.

Capability subclasses are soil groups within one class. They are designated by adding a small letter, *e*, *w*, *s*, or *c*, to the class numeral, for example, 2e. The letter *e* shows that the main hazard is the risk of erosion unless close-growing plant cover is maintained; *w* shows that water in or on the soil interferes with plant growth or cultivation (in some soils the wetness can be partly corrected by artificial drainage); *s* shows that the soil is limited mainly because it is shallow, droughty, or stony; and *c*, used in only some parts of the United States, shows that the chief limitation is climate that is very cold or very dry.

In class 1 there are no subclasses because the soils of this class have few limitations. Class 5 contains only the subclasses indicated by *w*, *s*, or *c* because the soils in class 5 are subject to little or no erosion. They have other limitations that restrict their use to pasture, rangeland, forestland, wildlife habitat, or recreation.

The capability classification of map units in this survey area is given in the section “Detailed Soil Map Units” and in table 7.

Prime Farmland

Prime farmland is one of several kinds of important farmland defined by the U.S. Department of Agriculture. It is of major importance in meeting the Nation’s short- and long-range needs for food and fiber. Because the supply of high-quality farmland is limited, the U.S. Department of Agriculture recognizes that responsible levels of government, as well as individuals, should encourage and facilitate the wise use of our Nation’s prime farmland.

Prime farmland, as defined by the U.S. Department of Agriculture, is land that has the best combination of physical and chemical characteristics for producing food, feed, forage, fiber, and oilseed crops and is available for these uses. It could be cultivated land, pastureland, forestland, or other land, but it is not urban or built-up land or water areas. The soil qualities,

growing season, and moisture supply are those needed for the soil to economically produce sustained high yields of crops when proper management, including water management, and acceptable farming methods are applied. In general, prime farmland has an adequate and dependable supply of moisture from precipitation or irrigation, a favorable temperature and growing season, acceptable acidity or alkalinity, an acceptable salt and sodium content, and few or no rocks. It is permeable to water and air. It is not excessively erodible or saturated with water for long periods, and it either is not frequently flooded during the growing season or is protected from flooding. Slope ranges mainly from 0 to 6 percent. More detailed information about the criteria for prime farmland is available at the local office of the Natural Resources Conservation Service.

A recent trend in land use in some parts of the survey area has been the loss of some prime farmland to industrial and urban uses. The loss of prime farmland to other uses puts pressure on marginal lands, which generally are more erodible, droughty, and less productive and cannot be easily cultivated.

About 465,325 acres, or nearly 83 percent of the county, meets the requirements for prime farmland.

The map units in the county that are considered prime farmland are listed in table 8. This list does not constitute a recommendation for a particular land use. On some soils included in the list, measures that overcome a hazard or limitation, such as flooding and wetness, are needed. Onsite evaluation is needed to determine whether or not the hazard or limitation has been overcome by corrective measures. The extent of each listed map unit is shown in table 5. The location is shown on the detailed soil maps. The soil qualities that affect use and management are described under the heading "Detailed Soil Map Units."

Hydric Soils

In this section, hydric soils are defined and described and the hydric soils in the survey area are listed.

The three essential characteristics of wetlands are hydrophytic vegetation, hydric soils, and wetland hydrology (Cowardin et al., 1979; U.S. Army Corps of Engineers, 1987; National Research Council, 1995; Tiner, 1985). Criteria for each of the characteristics must be met for areas to be identified as wetlands. Undrained hydric soils that have natural vegetation should support a dominant population of ecological wetland plant species. Hydric soils that have been

converted to other uses should be capable of being restored to wetlands.

Hydric soils are defined by the National Technical Committee for Hydric Soils (NTCHS) as soils that formed under conditions of saturation, flooding, or ponding long enough during the growing season to develop anaerobic conditions in the upper part (Federal Register, 1994). These soils are either saturated or inundated long enough during the growing season to support the growth and reproduction of hydrophytic vegetation.

The NTCHS definition identifies general soil properties that are associated with wetness. In order to determine whether a specific soil is a hydric soil or nonhydric soil, however, more specific information, such as information about the depth and duration of the water table, is needed. Thus, criteria that identify those estimated soil properties unique to hydric soils have been established (Federal Register, 1995). These criteria are used to identify a phase of a soil series that normally is associated with wetlands. The criteria used are selected estimated soil properties that are described in "Soil Taxonomy" (Soil Survey Staff, 1999) and in the "Soil Survey Manual" (Soil Survey Division Staff, 1993).

If soils are wet enough for a long enough period to be considered hydric, they should exhibit certain properties that can be easily observed in the field. These visible properties are indicators of hydric soils. The indicators used to make onsite determinations of hydric soils in this survey area are specified in "Field Indicators of Hydric Soils in the United States" (Hurt et al., 1998).

Hydric soils are identified by examining and describing the soil to a depth of about 20 inches. This depth may be greater if determination of an appropriate indicator so requires. It is always recommended that soils be excavated and described to the depth necessary for an understanding of the redoximorphic processes. Then, using the completed soil descriptions, soil scientists can compare the soil features required by each indicator and specify which indicators have been matched with the conditions observed in the soil. The soil can be identified as a hydric soil if at least one of the approved indicators is present.

The following map units meet the definition of hydric soils and, in addition, have at least one of the hydric soil indicators. This list can help in planning land uses; however, onsite investigation is recommended to determine the hydric soils on a specific site (National Research Council, 1995; Hurt et al., 1998).

- 45A Denny silt loam, 0 to 2 percent slopes
- 50A Virden silty clay loam, 0 to 2 percent slopes
- 68A Sable silty clay loam, 0 to 2 percent slopes
- 112A Cowden silt loam, 0 to 2 percent slopes
- 138A Shiloh silty clay loam, 0 to 2 percent slopes
- 244A Hartsburg silty clay loam, 0 to 2 percent slopes
- 249A Edinburg silty clay loam, 0 to 2 percent slopes
- 712A Spaulding silty clay loam, 0 to 2 percent slopes
- 3107A Sawmill silty clay loam, 0 to 2 percent slopes, frequently flooded
- 3405A Zook silty clay loam, 0 to 2 percent slopes, frequently flooded
- 8396A Vesser silt loam, 0 to 2 percent slopes, occasionally flooded

Map units that are made up mainly of hydric soils may have small areas, or inclusions, of nonhydric soils in the higher positions on the landform, and map units made up mainly of nonhydric soils may have inclusions of hydric soils in the lower positions on the landform. Table 9 indicates the hydric and nonhydric soils identified in the names of the detailed map units in the county. The table also identifies the included soils that are considered hydric. Onsite investigation is recommended to determine whether hydric soils occur and the location of the included hydric soils.

Windbreaks and Environmental Plantings

Windbreaks protect livestock, buildings, yards, fruit trees, gardens, and cropland from wind and snow; help to keep snow on fields; and provide food and cover for wildlife. Several rows of low- and high-growing broadleaf and coniferous trees and shrubs provide the most protection.

Field windbreaks are narrow plantings made at right angles to the prevailing wind and at specific intervals across the field. The interval depends on the erodibility of the soil.

Environmental plantings help to beautify and screen houses and other buildings and to abate noise. The plants, mostly evergreen shrubs and trees, are closely spaced. To ensure plant survival, a healthy planting stock of suitable species should be planted properly on a well prepared site and maintained in good condition.

Windbreaks are often planted on land that did not originally support trees. Knowledge of how trees perform on such land can be gained only by observing and recording the performance of trees that have been planted and have survived. Many popular windbreak

species are not indigenous to the areas in which they are planted.

Each tree or shrub species has certain climatic and physiographic limits. Within these parameters, a tree or shrub may grow well or grow poorly, depending on the characteristics of the soil. Each tree or shrub has definable potential heights in a given physiographic area and under a given climate. Accurate definitions of the potential heights are necessary when a windbreak is planned and designed.

Table 10 shows the height that locally grown trees and shrubs are expected to reach in 20 years on various soils. The estimates in table 10 are based on measurements and observation of established plantings that have been given adequate care. They can be used as a guide in planning windbreaks and screens. Additional information on planning windbreaks and screens and planting and caring for trees and shrubs can be obtained from the local office of the Natural Resources Conservation Service or of the Cooperative Extension Service or from a commercial nursery.

Forestland

John Churan, district forester, Illinois Department of Natural Resources, helped prepare this section.

Approximately 5 percent (29,100 acres) of Sangamon County is forested. Of this total, only about 2,000 acres is under management by a forester (Iverson et al., 1989).

The forestland is mainly in sloping areas, on the narrower bottomland, and in strips along the Sangamon River and its tributaries. Originally, these areas were heavily if not entirely forested, but the flatter areas were converted to agriculture long ago. Adjoining the upper slopes were flat forest areas that transitioned into prairie, gradually in some places (savannah sites) and abruptly in others. These areas have also, for the most part, been converted to agriculture, although forested areas remain in a few places. Overall, Sangamon County is part of the "Grand Prairie" (Swegman, 1979), and of some 129,400 original forest acres, about 20 percent remains today.

By far, the bulk of forest in the county is in areas of the sloping Hickory soils. Some forested areas extend onto the adjoining Alvin, Elco, Fayette, Keomah, Rozetta, Thebes, and other soils. Upland tree species are sensitive to differences in soils. Some species are adapted to certain sites but are only marginally adapted or are not adapted at all to other sites. A broad listing of species would include red chinquapin oak, white chinquapin oak, black chinquapin oak, bur

oak, hickory, walnut, elm, and hackberry. A serious disease affects white ash, which was once prominent. Hard maple is making strong incursions onto many of the shadier, moist slopes typically occupied by red oak. American elm is ubiquitous in the understory but occurs rarely as a larger tree.

Remnant forest sites are on the flood plains. These sites can be seeded readily, though reed canarygrass can be preemptive. Radford soils are the most common forest-associated soils on these sites, but Lawson, Tice, Sawmill, and certain other soils also are well represented. Soft maple is ubiquitous, but common associates include cottonwood, green ash, and sycamore. Some sites in which the soils have better internal drainage support walnut, hackberry, bitternut hickory, and assorted other bottomland species. Box elder, which is fairly widespread, is generally considered a “weed species.”

The many values of forestland include economic benefits (timber production), wildlife habitat, erosion control or prevention, ground-water infiltration or retention, water quality (including filtering and cleansing of agricultural pollutants), esthetic values, and recreational opportunities. All of these features can be enhanced through proper management.

Soil properties that affect the growth of trees include reaction (pH), fertility, drainage, texture, structure, and soil depth. The soil also serves as a reservoir for moisture, provides an anchor for tree roots, and supplies essential plant nutrients. Soils that do not have extremes of these properties and that have an effective rooting depth of more than 40 inches allow the best growth for wood production.

Site characteristics that affect tree growth include aspect (the direction in which the slopes face) and degree of slope. These site characteristics influence the amount of available sunlight, drainage, soil temperature, soil moisture, and relative humidity.

Typically, north and east aspects and the lower slope positions, which are cooler and have better moisture conditions than other sites, are the best upland sites for tree growth. The most productive sites on bottomland are generally the deep, moderately well drained soils that are subject to occasional flooding.

Management activities can influence forestland productivity and should be aimed at eliminating factors that cause tree stress. Generally, these activities involve thinning overstocked young stands; harvesting old, mature trees; and eliminating wildfire and grazing. Wildfire and grazing have very negative impacts on forest growth and quality. Grazing destroys the leaf layer on the surface, compacts the soils, and eliminates or damages tree seedlings. Forestland sites

that are not grazed have the highest potential for optimum timber production. Currently, relatively little forestland in the survey area is subject to grazing.

Where regeneration of forest stands is desired, the best approach is to concentrate on the mixture (or “composition”) of the species that originally occupied the site. Reforestation can be attempted by various means, including planting seedlings or tree seeds, encouraging natural regeneration, or a combination of these. Because of varying field conditions, planting trees in areas of bottomland along the Sangamon River can be difficult, especially in terms of timing, scheduling, and other logistics. Assistance can be obtained from a professional forester.

Similarly, a forester’s assistance should be sought before a stand is manipulated for the purpose of improving or optimizing composition, growth rates, and tree quality.

The tables in this section (tables 11 through 16) can help forest owners or managers plan the use of soils for wood crops. They show the potential productivity of the soils for wood crops and rate the soils according to the limitations that affect various aspects of forest management.

Forest Productivity

In table 11, the *potential productivity* of merchantable or *common trees* on a soil is expressed as a site index and as a volume number. The *site index* is the average height, in feet, that dominant and codominant trees of a given species attain in a specified number of years. The site index applies to fully stocked, even-aged, unmanaged stands. Commonly grown trees are those that forest managers generally favor in intermediate or improvement cuttings. They are selected on the basis of growth rate, quality, value, and marketability. More detailed information regarding site index is available in the “National Forestry Manual,” which is available in local offices of the Natural Resources Conservation Service or on the Internet.

The *volume of wood fiber*, a number, is the yield likely to be produced by the most important tree species. This number, expressed as cubic feet per acre per year and calculated at the age of culmination of the mean annual increment (CMAI), indicates the amount of fiber produced in a fully stocked, even-aged, unmanaged stand.

Suggested trees to plant are those that are preferred for planting, seeding, or natural regeneration and those that remain in the stand after thinning or partial harvest.

Forest Management

In tables 12 through 16, interpretive ratings are given for various aspects of forest management. The ratings are both verbal and numerical.

Some rating class terms indicate the degree to which the soils are suited to a specified forest management practice. *Well suited* indicates that the soil has features that are favorable for the specified practice and has no limitations. Good performance can be expected, and little or no maintenance is needed. *Moderately suited* indicates that the soil has features that are moderately favorable for the specified practice. One or more soil properties are less than desirable, and fair performance can be expected. Some maintenance is needed. *Poorly suited* indicates that the soil has one or more properties that are unfavorable for the specified practice. Overcoming the unfavorable properties requires special design, extra maintenance, and costly alteration. *Unsuited* indicates that the expected performance of the soil is unacceptable for the specified practice or that extreme measures are needed to overcome the undesirable soil properties.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.01 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the specified forest management practice (1.00) and the point at which the soil feature is not a limitation (0.00).

Rating class terms for fire damage and seedling mortality are expressed as *low*, *moderate*, and *high*. Where these terms are used, the numerical ratings indicate gradations between the point at which the potential for fire damage or seedling mortality is highest (1.00) and the point at which the potential is lowest (0.00).

The paragraphs that follow indicate the soil properties considered in rating the soils for forest management practices. More detailed information about the criteria used in the ratings is available in the "National Forestry Manual," which is available in local offices of the Natural Resources Conservation Service or on the Internet.

For *limitations affecting construction of haul roads and log landings*, the ratings are based on slope, flooding, permafrost, plasticity index, the hazard of soil slippage, content of sand, the Unified classification, rock fragments on or below the surface, depth to a restrictive layer that is indurated, depth to a water table, and ponding. The limitations are described as *slight*, *moderate*, or *severe*. A rating of *slight* indicates

that no significant limitations affect construction activities, *moderate* indicates that one or more limitations can cause some difficulty in construction, and *severe* indicates that one or more limitations can make construction very difficult or very costly.

The ratings of *suitability for log landings* are based on slope, rock fragments on the surface, plasticity index, content of sand, the Unified classification, depth to a water table, ponding, flooding, and the hazard of soil slippage. The soils are described as well suited, moderately suited, or poorly suited to use as log landings.

Ratings in the column *soil rutting hazard* are based on depth to a water table, rock fragments on or below the surface, the Unified classification, depth to a restrictive layer, and slope. Ruts form as a result of the operation of forest equipment. The hazard is described as *slight*, *moderate*, or *severe*. A rating of *slight* indicates that the soil is subject to little or no rutting, *moderate* indicates that rutting is likely, and *severe* indicates that ruts form readily.

Ratings in the column *hazard of off-road or off-trail erosion* are based on slope and on soil erodibility factor K. The soil loss is caused by sheet or rill erosion in off-road or off-trail areas where 50 to 75 percent of the surface has been exposed by logging, grazing, mining, or other kinds of disturbance. The hazard is described as *slight*, *moderate*, *severe*, or *very severe*. A rating of *slight* indicates that erosion is unlikely under ordinary climatic conditions; *moderate* indicates that some erosion is likely and that erosion-control measures may be needed; *severe* indicates that erosion is very likely and that erosion-control measures, including revegetation of bare areas, are advised; and *very severe* indicates that significant erosion is expected, loss of soil productivity and off-site damage are likely, and erosion-control measures are costly and generally impractical.

Ratings in the column *hazard of erosion on roads and trails* are based on the soil erodibility factor K, slope, and content of rock fragments. The ratings apply to unsurfaced roads and trails. The hazard is described as *slight*, *moderate*, or *severe*. A rating of *slight* indicates that little or no erosion is likely; *moderate* indicates that some erosion is likely, that the roads or trails may require occasional maintenance; and that simple erosion-control measures are needed; and *severe* indicates that significant erosion is expected, that the roads or trails require frequent maintenance, and that costly erosion-control measures are needed.

Ratings in the column *suitability for roads (natural surface)* are based on slope, rock fragments on the

surface, plasticity index, content of sand, the Unified classification, depth to a water table, ponding, flooding, and the hazard of soil slippage. The ratings indicate the suitability for using the natural surface of the soil for roads. The soils are described as well suited, moderately suited, or poorly suited to this use.

Ratings in the columns *suitability for hand planting* and *suitability for mechanical planting* are based on slope, depth to a restrictive layer, content of sand, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, moderately suited, poorly suited, or unsuited to these methods of planting. It is assumed that necessary site preparation is completed before seedlings are planted.

Ratings in the column *suitability for use of harvesting equipment* are based on slope, rock fragments on the surface, plasticity index, content of sand, the Unified classification, depth to a water table, and ponding. The soils are described as well suited, moderately suited, or poorly suited to this use.

Ratings in the column *suitability for mechanical site preparation (surface)* are based on slope, depth to a restrictive layer, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsuited to this management activity. The part of the soil from the surface to a depth of about 1 foot is considered in the ratings.

Ratings in the column *suitability for mechanical site preparation (deep)* are based on slope, depth to a restrictive layer, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsuited to this management activity. The part of the soil from the surface to a depth of about 3 feet is considered in the ratings.

Ratings in the column *potential for damage to soil by fire* are based on texture of the surface layer, content of rock fragments and organic matter in the surface layer, thickness of the surface layer, and slope. The soils are described as having a low, moderate, or high potential for this kind of damage. The ratings indicate an evaluation of the potential impact of prescribed fires or wildfires that are intense enough to remove the duff layer and consume organic matter in the surface layer.

Ratings in the column *potential for seedling mortality* are based on flooding, ponding, depth to a water table, content of lime, reaction, salinity, available water capacity, soil moisture regime, soil temperature regime, aspect, and slope. The soils are described as

having a low, moderate, or high potential for seedling mortality.

Recreation

Dan Stephenson, district fisheries biologist, Illinois Department of Natural Resources, helped prepare this section.

Sangamon County has many historic sites that attract tourists (USDA, 1972). Of special importance are the Lincoln Tomb State Memorial and the Lincoln Home State Memorial in Springfield.

The potential for development of recreational facilities is fair throughout most of the county. Lake Springfield and the Sangamon River provide good opportunities for fishing, boating, and swimming, and the surrounding areas provide good opportunities for picnicking, hunting, camping, and hiking. Several hundred acres of Sangchris Lake are in the southeastern part of the county. This State-owned lake is suitable for fishing, boating, and other recreational activities.

More than 70 species of fish inhabit the waters of Sangamon County. Prime opportunities for fishing are available on rivers, streams, lakes, and farm ponds. The county has about 428 miles, or 1,611 acres, of streams. The Sangamon River flows for more than 50 miles in the county. Major fishing areas are the Sangamon River, the South Fork of the Sangamon River, and Lick Creek and Sugar Creek directly west of Springfield. These areas support largemouth bass, white bass, crappie, sunfish species, walleye, drum, carp, and catfish species (Illinois Department of Conservation, 1971).

The county has approximately 1,100 lakes and ponds. Of these, about 1,000 are private ponds and 80 are organizational lakes. These private waters encompass about 1,116 acres. There are approximately 30 public bodies of water totaling more than 4,400 acres (Illinois Department of Natural Resources, 2000b).

Lake Springfield is the most heavily fished impoundment located entirely in the county. Approximately 25,000 fishing trips are made to that lake each year (Illinois Department of Conservation, 1991). Lake Sangchris, partially located in Sangamon County, is one of the most heavily fished lakes in the State. About 65,000 angling trips are made to this lake each year (Illinois Department of Natural Resources, 2000c). Both Lake Springfield and Lake Sangchris have excellent populations of largemouth bass, channel catfish, and flathead catfish. The populations

of crappie are very good in Lake Sangchris but have fallen off in recent years in Lake Springfield. Lake Springfield has a very good white bass and walleye fishery. Lake Sangchris supports a striped bass fishery and currently holds the State record for that species (31 pounds 7 ounces). Both lakes have average or slightly above average populations of bluegill.

There are several other public lakes in the county, including an Illinois Department of Transportation lake and Lakes Williamsville, Loami, and Washington Park. More than 13,000 residents of Sangamon County enjoy angling opportunities each year and add more than \$8 million to the local economy (Illinois Department of Natural Resources, 2000a).

The soils of the survey area are rated in tables 17 and 18 according to limitations that affect their suitability for recreation. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the recreational uses. *Not limited* indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. *Somewhat limited* indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. *Very limited* indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.01 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

The ratings in the tables are based on restrictive soil features, such as wetness, slope, and texture of the surface layer. Susceptibility to flooding is considered. Not considered in the ratings, but important in evaluating a site, are the location and accessibility of the area, the size and shape of the area and its scenic quality, vegetation, access to water, potential water impoundment sites, and access to public sewer lines. The capacity of the soil to absorb septic tank effluent and the ability of the soil to support

vegetation also are important. Soils that are subject to flooding are limited for recreational uses by the duration and intensity of flooding and the season when flooding occurs. In planning recreational facilities, onsite assessment of the height, duration, intensity, and frequency of flooding is essential.

The information in tables 17 and 18 can be supplemented by other information in this survey, for example, interpretations for building site development, construction materials, sanitary facilities, and water management.

Camp areas require site preparation, such as shaping and leveling the tent and parking areas, stabilizing roads and intensively used areas, and installing sanitary facilities and utility lines. Camp areas are subject to heavy foot traffic and some vehicular traffic. The ratings are based on the soil properties that affect the ease of developing camp areas and the performance of the areas after development. Slope, stoniness, and depth to bedrock or a cemented pan are the main concerns affecting the development of camp areas. The soil properties that affect the performance of the areas after development are those that influence trafficability and promote the growth of vegetation, especially in heavily used areas. For good trafficability, the surface of camp areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Picnic areas are subject to heavy foot traffic. Most vehicular traffic is confined to access roads and parking areas. The ratings are based on the soil properties that affect the ease of developing picnic areas and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of picnic areas. For good trafficability, the surface of picnic areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Playgrounds require soils that are nearly level, are free of stones, and can withstand intensive foot traffic.

The ratings are based on the soil properties that affect the ease of developing playgrounds and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of playgrounds. For good trafficability, the surface of the playgrounds should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Paths and trails for hiking and horseback riding should require little or no slope modification through cutting and filling. The ratings are based on the soil properties that affect trafficability and erodibility. These properties are stoniness, depth to a water table, ponding, flooding, slope, and texture of the surface layer.

Off-road motorcycle trails require little or no site preparation. They are not covered with surfacing material or vegetation. Considerable compaction of the soil material is likely. The ratings are based on the soil properties that influence erodibility, trafficability, dustiness, and the ease of revegetation. These properties are stoniness, slope, depth to a water table, ponding, flooding, and texture of the surface layer.

Golf fairways are subject to heavy foot traffic and some light vehicular traffic. Cutting or filling may be required. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer. The suitability of the soil for traps, tees, roughs, and greens is not considered in the ratings.

Wildlife Habitat

Michael Chandler, district wildlife habitat biologist, Illinois Department of Natural Resources, helped prepare this section.

The soils in Sangamon County support habitat for a variety of wildlife, including pheasant, quail, mourning dove, turkey, white-tailed deer, squirrel, rabbit, songbirds, fox, raccoon, mink, and muskrat. Snipe,

heron, and other shore birds inhabit the bottomland areas. The streams and lakes support smallmouth bass, catfish, carp, and sunfish. Many farm ponds are stocked with largemouth bass and bluegill. These ponds provide habitat for migratory ducks in spring and fall as well as habitat for giant Canada geese.

Most areas in the county can be improved for use as wildlife habitat. The map units described in the section "Soil Series and Detailed Soil Map Units" can be grouped into two major wildlife areas. These areas are described in the following paragraphs.

Wildlife area 1.—Osco, Keomah, Assumption, Ipava, Sable, Sawmill, Radford, and Tice soils are the major soil types in this wildlife area. These soils are nearly level to strongly sloping and are poorly drained to well drained. Sawmill, Radford, and Tice soils are subject to flooding.

This wildlife area consists mainly of cropland, much of which is used for corn or soybeans year after year. This area provides habitat for ring-necked pheasant, raccoon, deer, meadowlark, grasshoppers, sparrow, fox, snakes, and other openland wildlife.

The habitat is generally of poor quality because of the lack of crop residue, herbaceous nesting, roosting cover, woody cover, travel lanes, and hedgerows. Wildlife would benefit from delayed mowing of grassy cover on roadsides and ditchbanks and along waterways until after the nesting season. Protection of woody cover and management of crop residue are also important.

Wildlife area 2.—Fayette, Rozetta, Elco, Alvin, Navlys, Hickory, and Middletown soils are the major soil types in this wildlife area. These soils are gently sloping to steep and are well drained or moderately well drained.

This wildlife area borders the major streams in the county, and it provides much more diversified habitat than wildlife area 1. It consists of cropland, pasture, and forestland. The major game species are ring-necked pheasant, white-tailed deer, mourning dove, bobwhite quail, turkey, fox, squirrel, and rabbit. The nongame species include those that inhabit brushy cover and forestland in addition to those listed in the description of wildlife area 1.

Pasture management, protection of forestland from livestock, crop residue management, and delayed mowing of grassy cover can benefit wildlife in this area.

Soils affect the kind and amount of vegetation that is available to wildlife as food and cover. They also affect the construction of water impoundments. The kind and abundance of wildlife depend largely on the amount and distribution of food, cover, and water. Wildlife habitat can be created or improved by planting

appropriate vegetation, by maintaining the existing plant cover, or by promoting the natural establishment of desirable plants.

In table 19, the soils in the survey area are rated according to their potential for providing habitat for various kinds of wildlife. This information can be used in planning parks, wildlife refuges, nature study areas, and other developments for wildlife; in selecting soils that are suitable for establishing, improving, or maintaining specific elements of wildlife habitat; and in determining the intensity of management needed for each element of the habitat.

The potential of the soil is rated good, fair, poor, or very poor. A rating of *good* indicates that the element or kind of habitat is easily established, improved, or maintained. Few or no limitations affect management, and satisfactory results can be expected. A rating of *fair* indicates that the element or kind of habitat can be established, improved, or maintained in most places. Moderately intensive management is required for satisfactory results. A rating of *poor* indicates that limitations are severe for the designated element or kind of habitat. Habitat can be created, improved, or maintained in most places, but management is difficult and must be intensive. A rating of *very poor* indicates that restrictions for the element or kind of habitat are very severe and that unsatisfactory results can be expected. Creating, improving, or maintaining habitat is impractical or impossible.

The elements of wildlife habitat are described in the following paragraphs.

Grain and seed crops are domestic grains and seed-producing herbaceous plants. Soil properties and features that affect the growth of grain and seed crops are depth of the root zone, texture of the surface layer, available water capacity, wetness, slope, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of grain and seed crops are corn, soybeans, wheat, and oats.

Grasses and legumes are domestic perennial grasses and herbaceous legumes. Soil properties and features that affect the growth of grasses and legumes are depth of the root zone, texture of the surface layer, available water capacity, wetness, surface stoniness, flooding, and slope. Soil temperature and soil moisture also are considerations. Examples of grasses and legumes are brome grass, timothy, orchardgrass, clover, and alfalfa.

Wild herbaceous plants are native or naturally established grasses and forbs, including weeds. Soil properties and features that affect the growth of these plants are depth of the root zone, texture of the

surface layer, available water capacity, wetness, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of wild herbaceous plants are bluestem, indiangrass, goldenrod, beggarweed, ragweed, and foxtail.

Hardwood trees and woody understory produce nuts or other fruit, buds, catkins, twigs, bark, and foliage. Soil properties and features that affect the growth of hardwood trees and shrubs are depth of the root zone, available water capacity, and wetness. Examples of these plants are oak, cherry, cottonwood, apple, hawthorn, hickory, blackberry, elderberry, maple, green ash, and willow. Examples of fruit-producing shrubs that are suitable for planting on soils rated *good* are American plum, hazelnut, dogwood, and arrowwood.

Coniferous plants furnish browse and seeds. Soil properties and features that affect the growth of coniferous trees, shrubs, and ground cover are depth of the root zone, available water capacity, and wetness. Examples of coniferous plants are pine, spruce, cedar, fir, and juniper.

Wetland plants are annual and perennial wild herbaceous plants that grow on moist or wet sites. Submerged or floating aquatic plants are excluded. Soil properties and features affecting wetland plants are texture of the surface layer, wetness, reaction, salinity, slope, and surface stoniness. Examples of wetland plants are smartweed, wild millet, cordgrass, rushes, sedges, and reeds.

Shallow water areas have an average depth of less than 5 feet. Some are naturally wet areas. Others are created by dams, levees, or other water-control structures. Soil properties and features affecting shallow water areas are depth to bedrock, wetness, surface stoniness, slope, and permeability. Examples of shallow water areas are waterfowl feeding areas, wildlife watering developments, marshes, and beaver ponds and other wildlife ponds.

The habitat for various kinds of wildlife is described in the following paragraphs.

Habitat for openland wildlife consists of cropland, pasture, meadows, and areas that are overgrown with grasses, herbs, shrubs, and vines. These areas produce grain and seed crops, grasses and legumes, and wild herbaceous plants. Wildlife attracted to these areas include ring-necked pheasant, bobwhite quail, meadowlark, field sparrow, cottontail rabbit, and red fox.

Habitat for woodland wildlife consists of areas of deciduous and/or coniferous plants and associated grasses, legumes, and wild herbaceous plants. Wildlife attracted to these areas include wild turkey, thrushes,

woodpeckers, owls, tree squirrels, raccoon, woodcock, and white-tailed deer.

Habitat for wetland wildlife consists of open, marshy or swampy shallow water areas. Some of the wildlife attracted to such areas are ducks, geese, herons, shore birds, muskrat, mink, and beaver.

Engineering

This section provides information for planning land uses related to urban development and to water management. Soils are rated for various uses, and the most limiting features are identified. Ratings are given for building site development, sanitary facilities, construction materials, and water management. The ratings are based on observed performance of the soils and on the data in the tables described under the heading "Soil Properties."

Information in this section is intended for land use planning, for evaluating land use alternatives, and for planning site investigations prior to design and construction. The information, however, has limitations. For example, estimates and other data generally apply only to that part of the soil between the surface and a depth of 5 to 7 feet. Because of the map scale, small areas of different soils may be included within the mapped areas of a specific soil.

The information is not site specific and does not eliminate the need for onsite investigation of the soils or for testing and analysis by personnel experienced in the design and construction of engineering works.

Government ordinances and regulations that restrict certain land uses or impose specific design criteria were not considered in preparing the information in this section. Local ordinances and regulations should be considered in planning, in site selection, and in design.

Soil properties, site features, and observed performance were considered in determining the ratings in this section. During the fieldwork for this soil survey, determinations were made about particle-size distribution, liquid limit, plasticity index, soil reaction, depth to bedrock, hardness of bedrock within 5 to 7 feet of the surface, soil wetness, depth to a water table, ponding, slope, likelihood of flooding, natural soil structure aggregation, and soil density. Data were collected about kinds of clay minerals, mineralogy of the sand and silt fractions, and the kinds of adsorbed cations. Estimates were made for erodibility, permeability, corrosivity, shrink-swell potential, available water capacity, and other behavioral characteristics affecting engineering uses.

This information can be used to evaluate the potential of areas for residential, commercial,

industrial, and recreational uses; make preliminary estimates of construction conditions; evaluate alternative routes for roads, streets, highways, pipelines, and underground cables; evaluate alternative sites for sanitary landfills, septic tank absorption fields, and sewage lagoons; plan detailed onsite investigations of soils and geology; locate potential sources of gravel, sand, earthfill, and topsoil; plan drainage systems, irrigation systems, ponds, terraces, and other structures for soil and water conservation; and predict performance of proposed small structures and pavements by comparing the performance of existing similar structures on the same or similar soils.

The information in the tables, along with the soil maps, the soil descriptions, and other data provided in this survey, can be used to make additional interpretations.

Some of the terms used in this soil survey have a special meaning in soil science and are defined in the Glossary.

Building Site Development

Soil properties influence the development of building sites, including the selection of the site, the design of the structure, construction, performance after construction, and maintenance. Tables 20 and 21 show the degree and kind of soil limitations that affect dwellings with and without basements, small commercial buildings, local roads and streets, shallow excavations, and lawns and landscaping.

The ratings in the tables are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect building site development. *Not limited* indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. *Somewhat limited* indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. *Very limited* indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.01 to 1.00. They indicate gradations between the point at which a soil

feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Dwellings are single-family houses of three stories or less. For dwellings without basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. For dwellings with basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of about 7 feet. The ratings for dwellings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility. Compressibility is inferred from the Unified classification. The properties that affect the ease and amount of excavation include depth to a water table, ponding, flooding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Small commercial buildings are structures that are less than three stories high and do not have basements. The foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. The ratings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility (which is inferred from the Unified classification). The properties that affect the ease and amount of excavation include flooding, depth to a water table, ponding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Local roads and streets have an all-weather surface and carry automobile and light truck traffic all year. They have a subgrade of cut or fill soil material; a base of gravel, crushed rock, or soil material stabilized by lime or cement; and a surface of flexible material (asphalt), rigid material (concrete), or gravel with a binder. The ratings are based on the soil properties that affect the ease of excavation and grading and the traffic-supporting capacity. The properties that affect the ease of excavation and grading are depth to

bedrock or a cemented pan, hardness of bedrock or a cemented pan, depth to a water table, ponding, flooding, the amount of large stones, and slope. The properties that affect the traffic-supporting capacity are soil strength (as inferred from the AASHTO group index number), subsidence, linear extensibility (shrink-swell potential), the potential for frost action, depth to a water table, and ponding.

Shallow excavations are trenches or holes dug to a maximum depth of 5 or 6 feet for graves, utility lines, open ditches, or other purposes. The ratings are based on the soil properties that influence the ease of digging and the resistance to sloughing. Depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, the amount of large stones, and dense layers influence the ease of digging, filling, and compacting. Depth to the seasonal high water table, flooding, and ponding may restrict the period when excavations can be made. Slope influences the ease of using machinery. Soil texture, depth to the water table, and linear extensibility (shrink-swell potential) influence the resistance to sloughing.

Lawns and landscaping require soils on which turf and ornamental trees and shrubs can be established and maintained. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer.

Sanitary Facilities

Tables 22 and 23 show the degree and kind of soil limitations that affect septic tank absorption fields, sewage lagoons, sanitary landfills, and daily cover for landfill. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect these uses. *Not limited* indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. *Somewhat limited* indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. *Very limited* indicates that the soil has one

or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.01 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Septic tank absorption fields are areas in which effluent from a septic tank is distributed into the soil through subsurface tiles or perforated pipe. Only that part of the soil between depths of 24 and 60 inches is evaluated. The ratings are based on the soil properties that affect absorption of the effluent, construction and maintenance of the system, and public health. Permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, and flooding affect absorption of the effluent. Stones and boulders, ice, and bedrock or a cemented pan interfere with installation. Subsidence interferes with installation and maintenance. Excessive slope may cause lateral seepage and surfacing of the effluent in downslope areas.

Some soils are underlain by loose sand and gravel or fractured bedrock at a depth of less than 4 feet below the distribution lines. In these soils the absorption field may not adequately filter the effluent, particularly when the system is new. As a result, the ground water may become contaminated.

Sewage lagoons are shallow ponds constructed to hold sewage while aerobic bacteria decompose the solid and liquid wastes. Lagoons should have a nearly level floor surrounded by cut slopes or embankments of compacted soil. Nearly impervious soil material for the lagoon floor and sides is required to minimize seepage and contamination of ground water. Considered in the ratings are slope, permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, flooding, large stones, and content of organic matter.

Soil permeability is a critical property affecting the suitability for sewage lagoons. Most porous soils eventually become sealed when they are used as sites for sewage lagoons. Until sealing occurs, however, the hazard of pollution is severe. Soils that have a permeability rate of more than 2 inches per hour are too porous for the proper functioning of sewage lagoons. In these soils, seepage of the effluent can result in contamination of the ground water. Ground-water contamination is also a hazard if fractured

bedrock is within a depth of 40 inches, if the water table is high enough to raise the level of sewage in the lagoon, or if floodwater overtops the lagoon.

A high content of organic matter is detrimental to proper functioning of the lagoon because it inhibits aerobic activity. Slope, bedrock, and cemented pans can cause construction problems, and large stones can hinder compaction of the lagoon floor. If the lagoon is to be uniformly deep throughout, the slope must be gentle enough and the soil material must be thick enough over bedrock or a cemented pan to make land smoothing practical.

A trench sanitary landfill is an area where solid waste is placed in successive layers in an excavated trench. The waste is spread, compacted, and covered daily with a thin layer of soil excavated at the site. When the trench is full, a final cover of soil material at least 2 feet thick is placed over the landfill. The ratings in the table are based on the soil properties that affect the risk of pollution, the ease of excavation, trafficability, and revegetation. These properties include permeability, depth to bedrock or a cemented pan, depth to a water table, ponding, slope, flooding, texture, stones and boulders, highly organic layers, soil reaction, and content of salts and sodium. Unless otherwise stated, the ratings apply only to that part of the soil within a depth of about 6 feet. For deeper trenches, onsite investigation may be needed.

Hard, nonrippable bedrock, creviced bedrock, or highly permeable strata in or directly below the proposed trench bottom can affect the ease of excavation and the hazard of ground-water pollution. Slope affects construction of the trenches and the movement of surface water around the landfill. It also affects the construction and performance of roads in areas of the landfill.

Soil texture and consistence affect the ease with which the trench is dug and the ease with which the soil can be used as daily or final cover. They determine the workability of the soil when dry and when wet. Soils that are plastic and sticky when wet are difficult to excavate, grade, or compact and are difficult to place as a uniformly thick cover over a layer of refuse.

The soil material used as the final cover for a trench landfill should be suitable for plants. It should not have excess sodium or salts and should not be too acid. The surface layer generally has the best workability, the highest content of organic matter, and the best potential for plants. Material from the surface layer should be stockpiled for use as the final cover.

In an *area sanitary landfill*, solid waste is placed in successive layers on the surface of the soil. The waste is spread, compacted, and covered daily with a thin

layer of soil from a source away from the site. A final cover of soil material at least 2 feet thick is placed over the completed landfill. The ratings in the table are based on the soil properties that affect trafficability and the risk of pollution. These properties include flooding, permeability, depth to a water table, ponding, slope, and depth to bedrock or a cemented pan.

Flooding is a serious problem because it can result in pollution in areas downstream from the landfill. If permeability is too rapid or if fractured bedrock, a fractured cemented pan, or the water table is close to the surface, the leachate can contaminate the water supply. Slope is a consideration because of the extra grading required to maintain roads in the steeper areas of the landfill. Also, leachate may flow along the surface of the soils in the steeper areas and cause difficult seepage problems.

Daily cover for landfill is the soil material that is used to cover compacted solid waste in an area sanitary landfill. The soil material is obtained offsite, transported to the landfill, and spread over the waste. The ratings in the table also apply to the final cover for a landfill. They are based on the soil properties that affect workability, the ease of digging, and the ease of moving and spreading the material over the refuse daily during wet and dry periods. These properties include soil texture, depth to a water table, ponding, rock fragments, slope, depth to bedrock or a cemented pan, reaction, and content of salts, sodium, or lime.

Loamy or silty soils that are free of large stones and excess gravel are the best cover for a landfill. Clayey soils may be sticky and difficult to spread; sandy soils are subject to wind erosion.

Slope affects the ease of excavation and of moving the cover material. Also, it can influence runoff, erosion, and reclamation of the borrow area.

After soil material has been removed, the soil material remaining in the borrow area must be thick enough over bedrock, a cemented pan, or the water table to permit revegetation. The soil material used as the final cover for a landfill should be suitable for plants. It should not have excess sodium, salts, or lime and should not be too acid.

Construction Materials

Tables 24 and 25 give information about the soils as potential sources of gravel, sand, topsoil, and roadfill. Normal compaction, minor processing, and other standard construction practices are assumed.

Sand and *gravel* are natural aggregates suitable for commercial use with a minimum of processing. They are used in many kinds of construction. Specifications

for each use vary widely. In table 24, only the likelihood of finding material in suitable quantity is evaluated. The suitability of the material for specific purposes is not evaluated, nor are factors that affect excavation of the material. The properties used to evaluate the soil as a source of sand or gravel are gradation of grain sizes (as indicated by the Unified classification of the soil), the thickness of suitable material, and the content of rock fragments. If the bottom layer of the soil contains sand or gravel, the soil is considered a likely source regardless of thickness. The assumption is that the sand or gravel layer below the depth of observation exceeds the minimum thickness.

The soils are rated *good*, *fair*, or *poor* as potential sources of sand and gravel. A rating of *good* or *fair* means that the source material is likely to be in or below the soil. The bottom layer and the thickest layer of the soils are assigned numerical ratings. These ratings indicate the likelihood that the layer is a source of sand or gravel. The number 0.00 indicates that the layer is a poor source. The number 1.00 indicates that the layer is a good source. A number between 0.00 and 1.00 indicates the degree to which the layer is a likely source.

The soils are rated *good*, *fair*, or *poor* as potential sources of topsoil and roadfill. The features that limit the soils as sources of these materials are specified in the tables. The numerical ratings given after the specified features indicate the degree to which the features limit the soils as sources of topsoil or roadfill. The lower the number, the greater the limitation.

Topsoil is used to cover an area so that vegetation can be established and maintained. The upper 40 inches of a soil is evaluated for use as topsoil. Also evaluated is the reclamation potential of the borrow area. The ratings are based on the soil properties that affect plant growth; the ease of excavating, loading, and spreading the material; and reclamation of the borrow area. Toxic substances, soil reaction, and the properties that are inferred from soil texture, such as available water capacity and fertility, affect plant growth. The ease of excavating, loading, and spreading is affected by rock fragments, slope, depth to a water table, soil texture, and thickness of suitable material. Reclamation of the borrow area is affected by slope, depth to a water table, rock fragments, depth to bedrock or a cemented pan, and toxic material.

The surface layer of most soils is generally preferred for topsoil because of its organic matter content. Organic matter greatly increases the absorption and retention of moisture and nutrients for plant growth.

Roadfill is soil material that is excavated in one place and used in road embankments in another place. In this table, the soils are rated as a source of roadfill for low embankments, generally less than 6 feet high and less exacting in design than higher embankments.

The ratings are for the whole soil, from the surface to a depth of about 5 feet. It is assumed that soil layers will be mixed when the soil material is excavated and spread.

The ratings are based on the amount of suitable material and on soil properties that affect the ease of excavation and the performance of the material after it is in place. The thickness of the suitable material is a major consideration. The ease of excavation is affected by large stones, depth to a water table, and slope. How well the soil performs in place after it has been compacted and drained is determined by its strength (as inferred from the AASHTO classification of the soil) and linear extensibility (shrink-swell potential).

Water Management

Tables 26 through 28 give information on the soil properties and site features that affect water management. The degree and kind of soil limitations are given for pond reservoir areas; embankments, dikes, and levees; aquifer-fed excavated ponds; grassed waterways and surface drains; terraces and diversions; drainage; and sprinkler irrigation. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect these uses. *Not limited* indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. *Somewhat limited* indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. *Very limited* indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.01 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use

(1.00) and the point at which the soil feature is not a limitation (0.00).

Pond reservoir areas hold water behind a dam or embankment. Soils best suited to this use have low seepage potential in the upper 60 inches. The seepage potential is determined by the permeability of the soil and the depth to fractured bedrock or other permeable material. Excessive slope can affect the storage capacity of the reservoir area.

Embankments, dikes, and levees are raised structures of soil material, generally less than 20 feet high, constructed to impound water or to protect land against overflow. Embankments that have zoned construction (core and shell) are not considered. In this table, the soils are rated as a source of material for embankment fill. The ratings apply to the soil material below the surface layer to a depth of about 5 feet. It is assumed that soil layers will be uniformly mixed and compacted during construction.

The ratings do not indicate the ability of the natural soil to support an embankment. Soil properties to a depth even greater than the height of the embankment can affect performance and safety of the embankment. Generally, deeper onsite investigation is needed to determine these properties.

Soil material in embankments must be resistant to seepage, piping, and erosion and have favorable compaction characteristics. Unfavorable features include less than 5 feet of suitable material and a high content of stones or boulders, organic matter, or salts or sodium. A high water table affects the amount of usable material. It also affects trafficability.

Aquifer-fed excavated ponds are pits or dugouts that extend to a ground-water aquifer or to a depth below a permanent water table. Excluded are ponds that are fed only by surface runoff and embankment ponds that impound water 3 feet or more above the original surface. Excavated ponds are affected by depth to a permanent water table, permeability of the aquifer, and quality of the water as inferred from the salinity of the soil. Depth to bedrock and the content of large stones affect the ease of excavation.

Grassed waterways and surface drains are natural or constructed channels, generally broad and shallow, that conduct surface water to outlets at a nonerosive velocity. Large stones, wetness, slope, and depth to bedrock or a cemented pan affect the construction of grassed waterways. A hazard of wind erosion, low available water capacity, restricted rooting depth, toxic substances such as salts and sodium, and restricted permeability adversely affect the growth and maintenance of the grass after construction.

Terraces and diversions are embankments or a combination of channels and ridges constructed across a slope to control erosion and conserve moisture by intercepting runoff. Slope, wetness, large stones, and depth to bedrock or a cemented pan affect the construction of terraces and diversions. A restricted rooting depth, a severe hazard of wind erosion or water erosion, an excessively coarse texture, and restricted permeability adversely affect maintenance.

Drainage is the removal of excess surface and subsurface water from the soil. How easily and effectively the soil is drained depends on the depth to bedrock, a cemented pan, or other layers that affect the rate of water movement; permeability; depth to a high water table or depth of standing water if the soil is subject to ponding; slope; susceptibility to flooding; subsidence of organic layers; and the potential for frost action. Excavating and grading and the stability of ditchbanks are affected by depth to bedrock or a cemented pan, large stones, slope, and the hazard of cutbanks caving. The productivity of the soil after drainage is adversely affected by extreme acidity or by toxic substances in the root zone, such as salts, sodium, and sulfur. Availability of drainage outlets is not considered in the ratings.

Sprinkler irrigation is the controlled application of water to supplement rainfall and support plant growth. The design and management of an irrigation system are affected by depth to the water table, the need for drainage, flooding, available water capacity, intake rate, permeability, erosion hazard, and slope. The construction of a system is affected by large stones

and depth to bedrock or a cemented pan. The performance of a system is affected by the depth of the root zone, the amount of salts or sodium, and soil reaction.

Waste Management

Soil properties are important when organic waste is applied as fertilizer and wastewater is applied in irrigated areas. They also are important when the soil is used as a medium for the treatment and disposal of the organic waste and wastewater. Unfavorable soil properties can result in environmental damage.

The use of organic waste and wastewater as production resources results in energy and resource conservation and minimizes the problems associated with waste disposal. If disposal is the goal, applying a maximum amount of the organic waste or the wastewater to a minimal area holds costs to a minimum and environmental damage is the main hazard. If reuse is the goal, a minimum amount should be applied to a maximum area and environmental damage is unlikely.

Interpretations developed for waste management may include ratings for manure, food-processing waste, municipal sewage sludge, use of wastewater for irrigation, and treatment of wastewater by slow rate, overland flow, and rapid infiltration processes.

Specific information regarding waste management is available at the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

Soil Properties

Data relating to soil properties are collected during the course of the soil survey.

Soil properties are ascertained by field examination of the soils and by laboratory index testing of some benchmark soils. Established standard procedures are followed. During the survey, many shallow borings are made and examined to identify and classify the soils and to delineate them on the soil maps. Samples are taken from some typical profiles and tested in the laboratory to determine particle-size distribution, plasticity, and compaction characteristics.

Estimates of soil properties are based on field examinations, on laboratory tests of samples from the survey area, and on laboratory tests of samples of similar soils in nearby areas. Tests verify field observations, verify properties that cannot be estimated accurately by field observation, and help to characterize key soils.

The estimates of soil properties are shown in tables. They include engineering index properties, physical and chemical properties, and pertinent soil and water features.

Engineering Index Properties

Table 29 gives the engineering classifications and the range of index properties for the layers of each soil in the survey area.

Depth to the upper and lower boundaries of each layer is indicated.

Texture is given in the standard terms used by the U.S. Department of Agriculture. These terms are defined according to percentages of sand, silt, and clay in the fraction of the soil that is less than 2 millimeters in diameter (fig. 10). "Loam," for example, is soil that is 7 to 27 percent clay, 28 to 50 percent silt, and less than 52 percent sand. If the content of particles coarser than sand is 15 percent or more, an appropriate modifier is added, for example, "gravelly." Textural terms are defined in the Glossary.

Classification of the soils is determined according to the Unified soil classification system (ASTM, 2001) and the system adopted by the American Association




Figure 10.—Percentages of clay, silt, and sand in the basic USDA soil texture classes.

of State Highway and Transportation Officials (AASHTO, 2000).

The Unified system classifies soils according to properties that affect their use as construction material. Soils are classified according to particle-size distribution of the fraction less than 3 inches in diameter and according to plasticity index, liquid limit, and organic matter content. Sandy and gravelly soils are identified as GW, GP, GM, GC, SW, SP, SM, and SC; silty and clayey soils as ML, CL, OL, MH, CH, and OH; and highly organic soils as PT. Soils exhibiting engineering properties of two groups can have a dual classification, for example, CL-ML.

The AASHTO system classifies soils according to those properties that affect roadway construction and maintenance. In this system, the fraction of a mineral soil that is less than 3 inches in diameter is classified in one of seven groups from A-1 through A-7 on the basis of particle-size distribution, liquid limit, and

plasticity index. Soils in group A-1 are coarse grained and low in content of fines (silt and clay). At the other extreme, soils in group A-7 are fine grained. Highly organic soils are classified in group A-8 on the basis of visual inspection.

If laboratory data are available, the A-1, A-2, and A-7 groups are further classified as A-1-a, A-1-b, A-2-4, A-2-5, A-2-6, A-2-7, A-7-5, or A-7-6. As an additional refinement, the suitability of a soil as subgrade material can be indicated by a group index number. Group index numbers range from 0 for the best subgrade material to 20 or higher for the poorest.

Rock fragments larger than 10 inches in diameter and 3 to 10 inches in diameter are indicated as a percentage of the total soil on a dry-weight basis. The percentages are estimates determined mainly by converting volume percentage in the field to weight percentage.

Percentage (of soil particles) passing designated sieves is the percentage of the soil fraction less than 3 inches in diameter based on an oven-dry weight. The sieves, numbers 4, 10, 40, and 200 (USA Standard Series), have openings of 4.76, 2.00, 0.420, and 0.074 millimeters, respectively. Estimates are based on laboratory tests of soils sampled in the survey area and in nearby areas and on estimates made in the field.

Liquid limit and plasticity index (Atterberg limits) indicate the plasticity characteristics of a soil. The estimates are based on test data from the survey area or from nearby areas and on field examination.

The estimates of particle-size distribution, liquid limit, and plasticity index are generally rounded to the nearest 5 percent. Thus, if the ranges of gradation and Atterberg limits extend a marginal amount (1 or 2 percentage points) across classification boundaries, the classification in the marginal zone is generally omitted in the table.

Physical Properties

Table 30 shows estimates of some physical characteristics and features that affect soil behavior. These estimates are given for the layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils.

Depth to the upper and lower boundaries of each layer is indicated.

Particle size is the effective diameter of a soil particle as measured by sedimentation, sieving, or micrometric methods. Particle sizes are expressed as classes with specific effective diameter class limits.

The broad classes are sand, silt, and clay, ranging from the larger to the smaller.

Sand as a soil separate consists of mineral soil particles that are 0.05 millimeter to 2 millimeters in diameter. In table 30, the estimated sand content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Silt as a soil separate consists of mineral soil particles that are 0.002 to 0.05 millimeter in diameter. In table 30, the estimated silt content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Clay as a soil separate consists of mineral soil particles that are less than 0.002 millimeter in diameter. In table 30, the estimated clay content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

The content of sand, silt, and clay affects the physical behavior of a soil. Particle size is important for engineering and agronomic interpretations, for determination of soil hydrologic qualities, and for soil classification.

The amount and kind of clay affect the fertility and physical condition of the soil and the ability of the soil to adsorb cations and to retain moisture. They influence shrink-swell potential, permeability, plasticity, the ease of soil dispersion, and other soil properties. The amount and kind of clay in a soil also affect tillage and earthmoving operations.

Moist bulk density is the weight of soil (oven-dry) per unit volume. Volume is measured when the soil is at field moisture capacity, that is, the moisture content at $1/3$ - or $1/10$ -bar (33kPa or 10kPa) moisture tension. Weight is determined after the soil is dried at 105 degrees C. In the table, the estimated moist bulk density of each soil horizon is expressed in grams per cubic centimeter of soil material that is less than 2 millimeters in diameter. Bulk density data are used to compute shrink-swell potential, available water capacity, total pore space, and other soil properties. The moist bulk density of a soil indicates the pore space available for water and roots. Depending on soil texture, a bulk density of more than 1.4 can restrict water storage and root penetration. Moist bulk density is influenced by texture, kind of clay, content of organic matter, and soil structure.

Permeability (K_{sat}) refers to the ability of a soil to transmit water or air. The term "permeability," as used in soil surveys, indicates saturated hydraulic conductivity (K_{sat}). The estimates in the table indicate the rate of water movement, in inches per hour, when

the soil is saturated. They are based on soil characteristics observed in the field, particularly structure, porosity, and texture. Permeability is considered in the design of soil drainage systems and septic tank absorption fields.

Available water capacity refers to the quantity of water that the soil is capable of storing for use by plants. The capacity for water storage is given in inches of water per inch of soil for each soil layer. The capacity varies, depending on soil properties that affect retention of water. The most important properties are the content of organic matter, soil texture, bulk density, and soil structure. Available water capacity is an important factor in the choice of plants or crops to be grown and in the design and management of irrigation systems. Available water capacity is not an estimate of the quantity of water actually available to plants at any given time.

Linear extensibility refers to the change in length of an unconfined clod as moisture content is decreased from a moist to a dry state. It is an expression of the volume change between the water content of the clod at $1/3$ - or $1/10$ -bar tension (33kPa or 10kPa tension) and oven dryness. The volume change is reported in the table as percent change for the whole soil. Volume change is influenced by the amount and type of clay minerals in the soil.

Linear extensibility is used to determine the shrink-swell potential of soils. The shrink-swell potential is low if the soil has a linear extensibility of less than 3 percent; moderate if 3 to 6 percent; high if 6 to 9 percent; and very high if more than 9 percent. If the linear extensibility is more than 3, shrinking and swelling can cause damage to buildings, roads, and other structures and to plant roots. Special design commonly is needed.

Organic matter is the plant and animal residue in the soil at various stages of decomposition. In table 30, the estimated content of organic matter is expressed as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

The content of organic matter in a soil can be maintained by returning crop residue to the soil. Organic matter has a positive effect on available water capacity, water infiltration, soil organism activity, and tilth. It is a source of nitrogen and other nutrients for crops and soil organisms.

Erosion factors are shown in table 30 as the K factor (K_w and K_f) and the T factor. Erosion factor K indicates the susceptibility of a soil to sheet and rill erosion by water. Factor K is one of several factors used in the Universal Soil Loss Equation (USLE) and the Revised Universal Soil Loss Equation (RUSLE) to predict the average annual rate of soil loss by sheet

and rill erosion in tons per acre per year. The estimates are based primarily on percentage of silt, sand, and organic matter and on soil structure and permeability. Values of K range from 0.02 to 0.69. Other factors being equal, the higher the value, the more susceptible the soil is to sheet and rill erosion by water.

Erosion factor K_w indicates the erodibility of the whole soil. The estimates are modified by the presence of rock fragments.

Erosion factor K_f indicates the erodibility of the fine-earth fraction, or the material less than 2 millimeters in size.

Erosion factor T is an estimate of the maximum average annual rate of soil erosion by wind or water that can occur without affecting crop productivity over a sustained period. The rate is in tons per acre per year.

Wind erodibility groups are made up of soils that have similar properties affecting their susceptibility to wind erosion in cultivated areas. The soils assigned to group 1 are the most susceptible to wind erosion, and those assigned to group 8 are the least susceptible. The groups are described in the "National Soil Survey Handbook" (USDA, NRCS).

Wind erodibility index is a numerical value indicating the susceptibility of soil to wind erosion, or the tons per acre per year that can be expected to be lost to wind erosion. There is a close correlation between wind erosion and the texture of the surface layer, the size and durability of surface clods, rock fragments, organic matter, and a calcareous reaction. Soil moisture and frozen soil layers also influence wind erosion.

Chemical Properties

Table 31 shows estimates of some chemical characteristics and features that affect soil behavior. These estimates are given for the layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils.

Depth to the upper and lower boundaries of each layer is indicated.

Soil reaction is a measure of acidity or alkalinity. The pH of each soil horizon is based on many field tests. For many soils, values have been verified by laboratory analyses. Soil reaction is important in selecting crops and other plants, in evaluating soil amendments for fertility and stabilization, and in determining the risk of corrosion.

Cation-exchange capacity is the total amount of extractable bases that can be held by the soil,

expressed in terms of milliequivalents per 100 grams of soil at neutrality (pH 7.0) or at some other stated pH value. Soils having a low cation-exchange capacity hold fewer cations and may require more frequent applications of fertilizer than soils having a high cation-exchange capacity. The ability to retain cations reduces the hazard of ground-water pollution.

Calcium carbonate equivalent is the percent of carbonates, by weight, in the fraction of the soil less than 2 millimeters in size. The availability of plant nutrients is influenced by the amount of carbonates in the soil. Incorporating nitrogen fertilizer into calcareous soils helps to prevent nitrite accumulation and ammonium-N volatilization.

Water Features

Table 32 gives estimates of various water features. The estimates are used in land use planning that involves engineering considerations.

Hydrologic soil groups are based on estimates of runoff potential. Soils are assigned to one of four groups according to the rate of water infiltration when the soils are not protected by vegetation, are thoroughly wet, and receive precipitation from long-duration storms.

The four hydrologic soil groups are:

Group A. Soils having a high infiltration rate (low runoff potential) when thoroughly wet. These consist mainly of deep, well drained to excessively drained sands or gravelly sands. These soils have a high rate of water transmission.

Group B. Soils having a moderate infiltration rate when thoroughly wet. These consist chiefly of moderately deep or deep, moderately well drained or well drained soils that have moderately fine texture to moderately coarse texture. These soils have a moderate rate of water transmission.

Group C. Soils having a slow infiltration rate when thoroughly wet. These consist chiefly of soils having a layer that impedes the downward movement of water or soils of moderately fine texture or fine texture. These soils have a slow rate of water transmission.

Group D. Soils having a very slow infiltration rate (high runoff potential) when thoroughly wet. These consist chiefly of clays that have a high shrink-swell potential, soils that have a high water table, soils that have a claypan or clay layer at or near the surface, and soils that are shallow over nearly impervious material. These soils have a very slow rate of water transmission.

If a soil is assigned to a dual hydrologic group (A/D, B/D, or C/D), the first letter is for drained areas and the second is for undrained areas.

The *months* in the table indicate the portion of the year in which the feature is most likely to be a concern.

Water table refers to a saturated zone in the soil. Table 32 indicates, by month, depth to the top (*upper limit*) and base (*lower limit*) of the saturated zone in most years. Estimates of the upper and lower limits are based mainly on observations of the water table at selected sites and on evidence of a saturated zone, namely grayish colors or mottles (redoximorphic features) in the soil. A saturated zone that lasts for less than a month is not considered a water table.

Ponding is standing water in a closed depression. Unless a drainage system is installed, the water is removed only by percolation, transpiration, or evaporation. Table 32 indicates *surface water depth* and the *duration* and *frequency* of ponding. Duration is expressed as *very brief* if less than 2 days, *brief* if 2 to 7 days, *long* if 7 to 30 days, and *very long* if more than 30 days. Frequency is expressed as none, rare, occasional, and frequent. *None* means that ponding is not probable; *rare* that it is unlikely but possible under unusual weather conditions (the chance of ponding is nearly 0 percent to 5 percent in any year); *occasional* that it occurs, on the average, once or less in 2 years (the chance of ponding is 5 to 50 percent in any year); and *frequent* that it occurs, on the average, more than once in 2 years (the chance of ponding is more than 50 percent in any year).

Flooding is the temporary inundation of an area caused by overflowing streams, by runoff from adjacent slopes, or by tides. Water standing for short periods after rainfall or snowmelt is not considered flooding, and water standing in swamps and marshes is considered ponding rather than flooding.

Duration and *frequency* are estimated. Duration is expressed as *extremely brief* if 0.1 hour to 4 hours, *very brief* if 4 hours to 2 days, *brief* if 2 to 7 days, *long* if 7 to 30 days, and *very long* if more than 30 days. Frequency is expressed as none, very rare, rare, occasional, frequent, and very frequent. *None* means that flooding is not probable; *very rare* that it is very unlikely but possible under extremely unusual weather conditions (the chance of flooding is less than 1 percent in any year); *rare* that it is unlikely but possible under unusual weather conditions (the chance of flooding is 1 to 5 percent in any year); *occasional* that it occurs infrequently under normal weather conditions (the chance of flooding is 5 to 50 percent in any year); *frequent* that it is likely to occur often under normal weather conditions (the chance of flooding is more than 50 percent in any year but is less than 50 percent in all months in any year); and *very frequent* that it is likely to occur very often under normal weather

conditions (the chance of flooding is more than 50 percent in all months of any year).

The information is based on evidence in the soil profile, namely thin strata of gravel, sand, silt, or clay deposited by floodwater; irregular decrease in organic matter content with increasing depth; and little or no horizon development.

Also considered are local information about the extent and levels of flooding and the relation of each soil on the landscape to historic floods. Information on the extent of flooding based on soil data is less specific than that provided by detailed engineering surveys that delineate flood-prone areas at specific flood frequency levels.

Soil Features

Table 33 gives estimates of various soil features. The estimates are used in land use planning that involves engineering considerations.

A *restrictive layer* is a nearly continuous layer that has one or more physical, chemical, or thermal properties that significantly impede the movement of water and air through the soil or that restrict roots or otherwise provide an unfavorable root environment. Examples are bedrock, cemented layers, dense layers, and frozen layers. The table indicates the hardness and thickness of the restrictive layer, both of which significantly affect the ease of excavation. *Depth to top* is the vertical distance from the soil surface to the upper boundary of the restrictive layer.

Potential for frost action is the likelihood of upward or lateral expansion of the soil caused by the formation of segregated ice lenses (frost heave) and the subsequent collapse of the soil and loss of strength on thawing. Frost action occurs when moisture moves

into the freezing zone of the soil. Temperature, texture, density, permeability, content of organic matter, and depth to the water table are the most important factors considered in evaluating the potential for frost action. It is assumed that the soil is not insulated by vegetation or snow and is not artificially drained. Silty and highly structured, clayey soils that have a high water table in winter are the most susceptible to frost action. Well drained, very gravelly, or very sandy soils are the least susceptible. Frost heave and low soil strength during thawing cause damage to pavements and other rigid structures.

Risk of corrosion pertains to potential soil-induced electrochemical or chemical action that corrodes or weakens uncoated steel or concrete. The rate of corrosion of uncoated steel is related to such factors as soil moisture, particle-size distribution, acidity, and electrical conductivity of the soil. The rate of corrosion of concrete is based mainly on the sulfate and sodium content, texture, moisture content, and acidity of the soil. Special site examination and design may be needed if the combination of factors results in a severe hazard of corrosion. The steel or concrete in installations that intersect soil boundaries or soil layers is more susceptible to corrosion than the steel or concrete in installations that are entirely within one kind of soil or within one soil layer.

For uncoated steel, the risk of corrosion, expressed as *low*, *moderate*, or *high*, is based on soil drainage class, total acidity, electrical resistivity near field capacity, and electrical conductivity of the saturation extract.

For concrete, the risk of corrosion also is expressed as *low*, *moderate*, or *high*. It is based on soil texture, acidity, and amount of sulfates in the saturation extract.

References

- American Association of State Highway and Transportation Officials (AASHTO). 2000. Standard specifications for transportation materials and methods of sampling and testing. 20th edition, 2 volumes.
- American Society for Testing and Materials (ASTM). 2001. Standard classification of soils for engineering purposes. ASTM Standard D 2487-00.
- Bergstrom, Robert E., Kemal Piskin, and Leon R. Follmer. 1976. Geology for planning in the Springfield-Decatur region, Illinois. Illinois State Geological Survey Circular 497.
- Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.
- Federal Register. July 13, 1994. Changes in hydric soils of the United States.
- Federal Register. February 24, 1995. Hydric soils of the United States.
- Fehrenbacher, J.B., R.A. Pope, I.J. Jansen, J.D. Alexander, and B.W. Ray. 1978. Soil productivity in Illinois. University of Illinois, College of Agriculture, Cooperative Extension Service Circular 1156.
- Hudson, Berman D. 1992. The soil survey as paradigm-based science. Soil Science Society of America Journal 56(3): 836-41.
- Hurt, G.W., P.M. Whited, and R.F. Pringle, editors. Version 4.0, 1998. Field indicators of hydric soils in the United States.
- Illinois Agricultural Statistics Service. 2002. Illinois county estimates, corn, soybeans, and wheat 2000-2001. Supplement to Bulletin 01-1. United States Department of Agriculture and Illinois Department of Agriculture. (Available on-line at <http://www.agstats.state.il.us/website/reports.htm>)
- Illinois Department of Conservation, Division of Fisheries. 1971. Sangamon County surface water resources.
- Illinois Department of Conservation, Division of Fisheries. 1991. Lake Springfield creel survey.
- Illinois Department of Natural Resources, Division of Fisheries. 2000a. Illinois fishing license sales.
- Illinois Department of Natural Resources, Division of Fisheries. 2000b. Inventory of Illinois surface water resources.

Illinois Department of Natural Resources, Division of Fisheries. 2000c. Lake Sanchris creel survey.

Iverson L.R., R.L. Oliver, D.P. Tucker, P.G. Risser, C.D. Burnett, and R.G. Rayburn. 1989. The forest resources of Illinois: An atlas and analysis of spatial and temporal trends. Illinois Natural History Survey Special Publication 11.

Jenny, Hans. 1941. Factors of soil formation.

National Research Council. 1995. Wetlands: Characteristics and boundaries.

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18.

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service. U.S. Department of Agriculture Handbook 436.

Steinkamp, James F. 1980. Soil survey of Sangamon County, Illinois. United States Department of Agriculture, Soil Conservation Service.

Swegman, John. 1979. The natural divisions of Illinois, land cover of Illinois. Illinois Department of Natural Resources, Forestry Division.

Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.

United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.

United States Census Bureau. 2002. 2000 Census of population and housing. Available online at <http://factfinder.census.gov/servlet/BasicFactsServlet> (March 2002).

United States Department of Agriculture. 1972. Study of the resources of central Illinois. Section VI—recreation enterprises.

United States Department of Agriculture, National Agricultural Statistics Service. 1997. 1997 Census of agriculture. (Reports available online at <http://www.nass.usda.gov/census/>)

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. (<http://soils.usda.gov/technical/handbook/>)

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210.

United States Department of Agriculture, Soil Conservation Service. 1981. Land resource regions and major land resource areas of the United States. U.S. Department of Agriculture Handbook 296. Map revised in 2004.

Glossary

ABC soil. A soil having an A, a B, and a C horizon.

Ablation till. Loose, permeable till deposited during the final downwasting of glacial ice. Lenses of crudely sorted sand and gravel are common.

AC soil. A soil having only an A and a C horizon. Commonly, such soil formed in recent alluvium or on steep, rocky slopes.

Aeration, soil. The exchange of air in soil with air from the atmosphere. The air in a well aerated soil is similar to that in the atmosphere; the air in a poorly aerated soil is considerably higher in carbon dioxide and lower in oxygen.

Aggregate, soil. Many fine particles held in a single mass or cluster. Natural soil aggregates, such as granules, blocks, or prisms, are called peds. Clods are aggregates produced by tillage or logging.

Alluvial fan. The fanlike deposit of a stream where it issues from a gorge upon a plain or of a tributary stream near or at its junction with its main stream.

Alluvium. Material, such as sand, silt, or clay, deposited on land by streams.

Alpha,alpha-dipyridyl. A dye that when dissolved in 1N ammonium acetate is used to detect the presence of reduced iron (Fe II) in the soil. A positive reaction indicates a type of redoximorphic feature.

Animal unit month (AUM). The amount of forage required by one mature cow of approximately 1,000 pounds weight, with or without a calf, for 1 month.

Aquic conditions. Current soil wetness characterized by saturation, reduction, and redoximorphic features.

Argillic horizon. A subsoil horizon characterized by an accumulation of illuvial clay.

Aspect. The direction in which a slope faces.

Available water capacity (available moisture capacity). The capacity of soils to hold water available for use by most plants. It is commonly defined as the difference between the amount of soil water at field moisture capacity and the amount at wilting point. It is commonly expressed as inches of water per inch of soil. The capacity, in

inches, in a 60-inch profile or to a limiting layer is expressed as:

Very low	0 to 3
Low	3 to 6
Moderate	6 to 9
High	9 to 12
Very high	more than 12

Backslope. The position that forms the steepest and generally linear, middle portion of a hillslope. In profile, backslopes are commonly bounded by a convex shoulder above and a concave footslope below.

Basal till. Compact glacial till deposited beneath the ice.

Base saturation. The degree to which material having cation-exchange properties is saturated with exchangeable bases (sum of Ca, Mg, Na, and K), expressed as a percentage of the total cation-exchange capacity.

Base slope. A geomorphic component of hills consisting of the concave to linear (perpendicular to the contour) slope that, regardless of the lateral shape, forms an apron or wedge at the bottom of a hillside dominated by colluvium and slope-wash sediments (for example, slope alluvium).

Bedding planes. Fine strata, less than 5 millimeters thick, in unconsolidated alluvial, eolian, lacustrine, or marine sediment.

Bedrock. The solid rock that underlies the soil and other unconsolidated material or that is exposed at the surface.

Bisequum. Two sequences of soil horizons, each of which consists of an illuvial horizon and the overlying eluvial horizons.

Bottomland. The normal flood plain of a stream, subject to flooding.

Boulders. Rock fragments larger than 2 feet (60 centimeters) in diameter.

Brush management. Use of mechanical, chemical, or biological methods to make conditions favorable for reseeding or to reduce or eliminate competition from woody vegetation and thus allow understory grasses and forbs to recover. Brush management

increases forage production and thus reduces the hazard of erosion. It can improve the habitat for some species of wildlife.

Calcareous soil. A soil containing enough calcium carbonate (commonly combined with magnesium carbonate) to effervesce visibly when treated with cold, dilute hydrochloric acid.

Capillary water. Water held as a film around soil particles and in tiny spaces between particles. Surface tension is the adhesive force that holds capillary water in the soil.

Catena. A sequence, or “chain,” of soils on a landscape that formed in similar kinds of parent material but have different characteristics as a result of differences in relief and drainage.

Cation. An ion carrying a positive charge of electricity. The common soil cations are calcium, potassium, magnesium, sodium, and hydrogen.

Cation-exchange capacity. The total amount of exchangeable cations that can be held by the soil, expressed in terms of milliequivalents per 100 grams of soil at neutrality (pH 7.0) or at some other stated pH value. The term, as applied to soils, is synonymous with base-exchange capacity but is more precise in meaning.

Chemical treatment. Control of unwanted vegetation through the use of chemicals.

Chiseling. Tillage with an implement having one or more soil-penetrating points that shatter or loosen hard, compacted layers to a depth below normal plow depth.

Clay. As a soil separate, the mineral soil particles less than 0.002 millimeter in diameter. As a soil textural class, soil material that is 40 percent or more clay, less than 45 percent sand, and less than 40 percent silt.

Clay depletions. Low-chroma zones having a low content of iron, manganese, and clay because of the chemical reduction of iron and manganese and the removal of iron, manganese, and clay. A type of redoximorphic depletion.

Clay film. A thin coating of oriented clay on the surface of a soil aggregate or lining pores or root channels. Synonyms: clay coating, clay skin.

Climax plant community. The stabilized plant community on a particular site. The plant cover reproduces itself and does not change so long as the environment remains the same.

Coarse textured soil. Sand or loamy sand.

Cobble (or cobblestone). A rounded or partly rounded fragment of rock 3 to 10 inches (7.6 to 25 centimeters) in diameter.

Cobbly soil material. Material that has 15 to 35 percent, by volume, rounded or partially rounded

rock fragments 3 to 10 inches (7.6 to 25 centimeters) in diameter. Very cobbly soil material has 35 to 60 percent of these rock fragments, and extremely cobbly soil material has more than 60 percent.

COLE (coefficient of linear extensibility). See Linear extensibility.

Colluvium. Soil material or rock fragments, or both, moved by creep, slide, or local wash and deposited at the base of steep slopes.

Complex slope. Irregular or variable slope. Planning or establishing terraces, diversions, and other water-control structures on a complex slope is difficult.

Complex, soil. A map unit of two or more kinds of soil or miscellaneous areas in such an intricate pattern or so small in area that it is not practical to map them separately at the selected scale of mapping. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas.

Concretions. Cemented bodies with crude internal symmetry organized around a point, a line, or a plane. They typically take the form of concentric layers visible to the naked eye. Calcium carbonate, iron oxide, and manganese oxide are common compounds making up concretions. If formed in place, concretions of iron oxide or manganese oxide are generally considered a type of redoximorphic concentration.

Conservation cropping system. Growing crops in combination with needed cultural and management practices. In a good conservation cropping system, the soil-improving crops and practices more than offset the effects of the soil-depleting crops and practices. Cropping systems are needed on all tilled soils. Soil-improving practices in a conservation cropping system include the use of rotations that contain grasses and legumes and the return of crop residue to the soil. Other practices include the use of green manure crops of grasses and legumes, proper tillage, adequate fertilization, and weed and pest control.

Conservation tillage. A tillage system that does not invert the soil and that leaves a protective amount of crop residue on the surface throughout the year.

Consistence, soil. Refers to the degree of cohesion and adhesion of soil material and its resistance to deformation when ruptured. Consistence includes resistance of soil material to rupture and to penetration; plasticity, toughness, and stickiness of puddled soil material; and the manner in which the soil material behaves when subject to

compression. Terms describing consistence are defined in the "Soil Survey Manual."

Control section. The part of the soil on which classification is based. The thickness varies among different kinds of soil, but for many it is that part of the soil profile between depths of 10 inches and 40 or 80 inches.

Corrosion. Soil-induced electrochemical or chemical action that dissolves or weakens concrete or uncoated steel.

Cover crop. A close-growing crop grown primarily to improve and protect the soil between periods of regular crop production, or a crop grown between trees and vines in orchards and vineyards.

Crop residue management. Returning crop residue to the soil, which helps to maintain soil structure, organic matter content, and fertility and helps to control erosion.

Cropping system. Growing crops according to a planned system of rotation and management practices.

Culmination of the mean annual increment (CMAI).

The average annual increase per acre in the volume of a stand. Computed by dividing the total volume of the stand by its age. As the stand increases in age, the mean annual increment continues to increase until mortality begins to reduce the rate of increase. The point where the stand reaches its maximum annual rate of growth is called the culmination of the mean annual increment.

Cutbanks cave (in tables). The walls of excavations tend to cave in or slough.

Deferred grazing. Postponing grazing or resting grazing land for a prescribed period.

Depression. A relatively sunken, low area surrounded by higher ground. Unlike an open depression, a closed depression has no natural outlet for surface water.

Depth, soil. Generally, the thickness of the soil over bedrock. Very deep soils are more than 60 inches deep over bedrock; deep soils, 40 to 60 inches; moderately deep, 20 to 40 inches; shallow, 10 to 20 inches; and very shallow, less than 10 inches.

Diversion (or diversion terrace). A ridge of earth, generally a terrace, built to protect downslope areas by diverting runoff from its natural course.

Drainage class (natural). Refers to the frequency and duration of wet periods under conditions similar to those under which the soil formed. Alterations of the water regime by human activities, either through drainage or irrigation, are not a consideration unless they have significantly changed the morphology of the soil. Seven

classes of natural soil drainage are recognized—*excessively drained, somewhat excessively drained, well drained, moderately well drained, somewhat poorly drained, poorly drained, and very poorly drained*. These classes are defined in the "Soil Survey Manual."

Drainage, surface. Runoff, or surface flow of water, from an area.

Drainageway. A relatively small, linear depression that at some time moves concentrated water and either has no defined channel or has a small defined channel.

Eluviation. The movement of material in true solution or colloidal suspension from one place to another within the soil. Soil horizons that have lost material through eluviation are eluvial; those that have received material are illuvial.

End moraine. A ridgelike accumulation produced at the outer margin of an actively flowing glacier at any given time.

Endosaturation. A type of saturation of the soil in which all horizons between the upper boundary of saturation and a depth of 2 meters are saturated.

Eolian deposit. Earthy parent material accumulated through wind action; commonly refers to sandy material in dunes or to loess in blankets on the surface.

Ephemeral stream. A stream, or reach of a stream, that flows only in direct response to precipitation. It receives no long-continued supply from melting snow or other source, and its channel is above the water table at all times.

Episaturation. A type of saturation indicating a perched water table in a soil in which saturated layers are underlain by one or more unsaturated layers within 2 meters of the surface.

Erosion. The wearing away of the land surface by water, wind, ice, or other geologic agents and by such processes as gravitational creep.

Erosion (geologic). Erosion caused by geologic processes acting over long geologic periods and resulting in the wearing away of mountains and the building up of such landscape features as flood plains and coastal plains. Synonym: natural erosion.

Erosion (accelerated). Erosion much more rapid than geologic erosion, mainly as a result of human or animal activities or of a catastrophe in nature, such as a fire, that exposes the surface.

Escarpment. A relatively continuous and steep slope or cliff breaking the general continuity of more gently sloping land surfaces and resulting from erosion or faulting. Synonym: scarp.

Fan terrace. A relict alluvial fan, no longer a site of active deposition, incised by younger and lower alluvial surfaces.

Fertility, soil. The quality that enables a soil to provide plant nutrients, in adequate amounts and in proper balance, for the growth of specified plants when light, moisture, temperature, tilth, and other growth factors are favorable.

Field moisture capacity. The moisture content of a soil, expressed as a percentage of the oven-dry weight, after the gravitational, or free, water has drained away; the field moisture content 2 or 3 days after a soaking rain; also called *normal field capacity*, *normal moisture capacity*, or *capillary capacity*.

Fine textured soil. Sandy clay, silty clay, or clay.

First bottom. The normal flood plain of a stream, subject to frequent or occasional flooding.

Flaggy soil material. Material that has, by volume, 15 to 35 percent flagstones. Very flaggy soil material has 35 to 60 percent flagstones, and extremely flaggy soil material has more than 60 percent flagstones.

Flagstone. A thin fragment of sandstone, limestone, slate, shale, or (rarely) schist 6 to 15 inches (15 to 38 centimeters) long.

Flood plain. A nearly level alluvial plain that borders a stream and is subject to flooding unless protected artificially.

Footslope. The position that forms the inner, gently inclined surface at the base of a hillslope. In profile, footslopes are commonly concave. A footslope is a transition zone between upslope sites of erosion and transport (shoulders and backslopes) and downslope sites of deposition (toeslopes).

Forb. Any herbaceous plant not a grass or a sedge.

Forest cover. All trees and other woody plants (underbrush) covering the ground in a forest.

Forest habitat type. An association of dominant tree and ground flora species in a climax community.

Forest type. A stand of trees similar in composition and development because of given physical and biological factors by which it may be differentiated from other stands.

Genesis, soil. The mode of origin of the soil. Refers especially to the processes or soil-forming factors responsible for the formation of the solum, or true soil, from the unconsolidated parent material.

Geomorphology. The science that treats the general configuration of the earth's surface; specifically, the study of the classification, description, nature, origin, and development of landforms and their relationships to underlying structures and the

history of geologic changes as recorded by these surface features. The term is especially applied to the genetic interpretation of landforms.

Glacial drift. Pulverized and other rock material transported by glacial ice and then deposited. Also, the sorted and unsorted material deposited by streams flowing from glaciers.

Glacial outwash. Gravel, sand, and silt, commonly stratified, deposited by glacial meltwater.

Glaciofluvial deposits. Material moved by glaciers and subsequently sorted and deposited by streams flowing from the melting ice. The deposits are stratified and occur as kames, eskers, deltas, and outwash plains.

Gleyed soil. Soil that formed under poor drainage, resulting in the reduction of iron and other elements in the profile and in gray colors.

Grassed waterway. A natural or constructed waterway, typically broad and shallow, seeded to grass as protection against erosion. Conducts surface water away from cropland.

Gravel. Rounded or angular fragments of rock as much as 3 inches (2 millimeters to 7.6 centimeters) in diameter. An individual piece is a pebble.

Gravelly soil material. Material that has 15 to 35 percent, by volume, rounded or angular rock fragments, not prominently flattened, as much as 3 inches (7.6 centimeters) in diameter.

Green manure crop (agronomy). A soil-improving crop grown to be plowed under in an early stage of maturity or soon after maturity.

Ground moraine. An extensive, fairly even layer of till having an uneven or undulating surface; a deposit of rock and mineral debris dragged along in, on, or beneath a glacier.

Ground water. Water filling all the unblocked pores of the material below the water table.

Gully. A miniature valley with steep sides cut by running water and through which water ordinarily runs only after rainfall. The distinction between a gully and a rill is one of depth. A gully generally is an obstacle to farm machinery and is too deep to be obliterated by ordinary tillage; a rill is of lesser depth and can be smoothed over by ordinary tillage.

Hard bedrock. Bedrock that cannot be excavated except by blasting or by the use of special equipment that is not commonly used in construction.

Hardpan. A hardened or cemented soil horizon, or layer. The soil material is sandy, loamy, or clayey and is cemented by iron oxide, silica, calcium carbonate, or other substance.

Head slope. A geomorphic component of hills consisting of a laterally concave area of a hillside, especially at the head of a drainageway. The overland waterflow is converging.

High-chroma zones. Zones having a chroma of 3 or more. Typical color in areas of iron concentrations.

High-residue crops. Such crops as small grain and corn used for grain. If properly managed, residue from these crops can be used to control erosion until the next crop in the rotation is established. These crops return large amounts of organic matter to the soil.

Hill. A natural elevation of the land surface, rising as much as 1,000 feet above surrounding lowlands, commonly of limited summit area and having a well defined outline; hillsides generally have slopes of more than 15 percent. The distinction between a hill and a mountain is arbitrary and is dependent on local usage.

Horizon, soil. A layer of soil, approximately parallel to the surface, having distinct characteristics produced by soil-forming processes. In the identification of soil horizons, an uppercase letter represents the major horizons. Numbers or lowercase letters that follow represent subdivisions of the major horizons. An explanation of the subdivisions is given in the "Soil Survey Manual." The major horizons of mineral soil are as follows:

O horizon.—An organic layer of fresh and decaying plant residue.

A horizon.—The mineral horizon at or near the surface in which an accumulation of humified organic matter is mixed with the mineral material. Also, a plowed surface horizon, most of which was originally part of a B horizon.

E horizon.—The mineral horizon in which the main feature is loss of silicate clay, iron, aluminum, or some combination of these.

B horizon.—The mineral horizon below an A horizon. The B horizon is in part a layer of transition from the overlying A to the underlying C horizon. The B horizon also has distinctive characteristics, such as (1) accumulation of clay, sesquioxides, humus, or a combination of these; (2) prismatic or blocky structure; (3) redder or browner colors than those in the A horizon; or (4) a combination of these.

C horizon.—The mineral horizon or layer, excluding indurated bedrock, that is little affected by soil-forming processes and does not have the properties typical of the overlying soil material. The material of a C horizon may be either like or unlike that in which the solum formed. If the

material is known to differ from that in the solum, an Arabic numeral, commonly a 2, precedes the letter C.

Cr horizon.—Soft, consolidated bedrock beneath the soil.

R layer.—Consolidated bedrock beneath the soil. The bedrock commonly underlies a C horizon, but it can be directly below an A or a B horizon.

Humus. The well decomposed, more or less stable part of the organic matter in mineral soils.

Hydrologic soil groups. Refers to soils grouped according to their runoff potential. The soil properties that influence this potential are those that affect the minimum rate of water infiltration on a bare soil during periods after prolonged wetting when the soil is not frozen. These properties are depth to a seasonal high water table, the infiltration rate and permeability after prolonged wetting, and depth to a very slowly permeable layer. The slope and the kind of plant cover are not considered but are separate factors in predicting runoff.

Illuviation. The movement of soil material from one horizon to another in the soil profile. Generally, material is removed from an upper horizon and deposited in a lower horizon.

Impervious soil. A soil through which water, air, or roots penetrate slowly or not at all. No soil is absolutely impervious to air and water all the time.

Infiltration. The downward entry of water into the immediate surface of soil or other material, as contrasted with percolation, which is movement of water through soil layers or material.

Infiltration rate. The rate at which water penetrates the surface of the soil at any given instant, usually expressed in inches per hour. The rate can be limited by the infiltration capacity of the soil or the rate at which water is applied at the surface.

Intake rate. The average rate of water entering the soil under irrigation. Most soils have a fast initial rate; the rate decreases with application time. Therefore, intake rate for design purposes is not a constant but is a variable depending on the net irrigation application. The rate of water intake, in inches per hour, is expressed as follows:

Less than 0.2	very low
0.2 to 0.4	low
0.4 to 0.75	moderately low
0.75 to 1.25	moderate
1.25 to 1.75	moderately high
1.75 to 2.5	high
More than 2.5	very high

Interfluve. An elevated area between two drainageways that sheds water to those drainageways.

Intermittent stream. A stream, or reach of a stream, that flows for prolonged periods only when it receives ground-water discharge or long, continued contributions from melting snow or other surface and shallow subsurface sources.

Iron concentrations. High-chroma zones having a high content of iron and manganese oxide because of chemical oxidation and accumulation, but having a clay content similar to that of the adjacent matrix. A type of redoximorphic concentration.

Iron depletions. Low-chroma zones having a low content of iron and manganese oxide because of chemical reduction and removal, but having a clay content similar to that of the adjacent matrix. A type of redoximorphic depletion.

Irrigation. Application of water to soils to assist in production of crops. Methods of irrigation are:

Basin.—Water is applied rapidly to nearly level plains surrounded by levees or dikes.

Border.—Water is applied at the upper end of a strip in which the lateral flow of water is controlled by small earth ridges called border dikes, or borders.

Controlled flooding.—Water is released at intervals from closely spaced field ditches and distributed uniformly over the field.

Corrugation.—Water is applied to small, closely spaced furrows or ditches in fields of close-growing crops or in orchards so that it flows in only one direction.

Drip (or trickle).—Water is applied slowly and under low pressure to the surface of the soil or into the soil through such applicators as emitters, porous tubing, or perforated pipe.

Furrow.—Water is applied in small ditches made by cultivation implements. Furrows are used for tree and row crops.

Sprinkler.—Water is sprayed over the soil surface through pipes or nozzles from a pressure system.

Subirrigation.—Water is applied in open ditches or tile lines until the water table is raised enough to wet the soil.

Wild flooding.—Water, released at high points, is allowed to flow onto an area without controlled distribution.

Knoll. A small, low, rounded hill rising above adjacent landforms.

K_{sat} . Saturated hydraulic conductivity. (See Permeability.)

Lamella. A thin (commonly less than 1 centimeter thick), discontinuous or continuous, generally horizontal layer of fine material (especially clay and iron oxides) that has been pedogenically

concentrated (illuviated within a coarser textured eluviated layer several centimeters to several decimeters thick).

Landslide. The rapid downhill movement of a mass of soil and loose rock, generally when wet or saturated. The speed and distance of movement, as well as the amount of soil and rock material, vary greatly.

Large stones (in tables). Rock fragments 3 inches (7.6 centimeters) or more across. Large stones adversely affect the specified use of the soil.

Leaching. The removal of soluble material from soil or other material by percolating water.

Linear extensibility. Refers to the change in length of an unconfined clod as moisture content is decreased from a moist to a dry state. Linear extensibility is used to determine the shrink-swell potential of soils. It is an expression of the volume change between the water content of the clod at $1/3$ - or $1/10$ -bar tension (33kPa or 10kPa tension) and oven dryness. Volume change is influenced by the amount and type of clay minerals in the soil. The volume change is the percent change for the whole soil. If it is expressed as a fraction, the resulting value is COLE, coefficient of linear extensibility.

Liquid limit. The moisture content at which the soil passes from a plastic to a liquid state.

Loam. Soil material that is 7 to 27 percent clay particles, 28 to 50 percent silt particles, and less than 52 percent sand particles.

Loess. Fine grained material, dominantly of silt-sized particles, deposited by wind.

Low strength. The soil is not strong enough to support loads.

Low-chroma zones. Zones having chroma of 2 or less. Typical color in areas of iron depletions.

Low-residue crops. Such crops as corn used for silage, peas, beans, and potatoes. Residue from these crops is not adequate to control erosion until the next crop in the rotation is established. These crops return little organic matter to the soil.

MAP. Mean annual precipitation, expressed in inches.

Masses. Concentrations of substances in the soil matrix that do not have a clearly defined boundary with the surrounding soil material and cannot be removed as a discrete unit. Common compounds making up masses are calcium carbonate, gypsum or other soluble salts, iron oxide, and manganese oxide. Masses consisting of iron oxide or manganese oxide generally are considered a type of redoximorphic concentration.

Mechanical treatment. Use of mechanical equipment

for seeding, brush management, and other management practices.

Medium textured soil. Very fine sandy loam, loam, silt loam, or silt.

Mineral soil. Soil that is mainly mineral material and low in organic material. Its bulk density is more than that of organic soil.

Minimum tillage. Only the tillage essential to crop production and prevention of soil damage.

Miscellaneous area. An area that has little or no natural soil and supports little or no vegetation.

Moderately coarse textured soil. Coarse sandy loam, sandy loam, or fine sandy loam.

Moderately fine textured soil. Clay loam, sandy clay loam, or silty clay loam.

Mollic epipedon. A thick, dark, humus-rich surface horizon (or horizons) that has high base saturation and pedogenic soil structure. It may include the upper part of the subsoil.

Moraine. An accumulation of earth, stones, and other debris deposited by a glacier. Some types are terminal, lateral, medial, and ground.

Morphology, soil. The physical makeup of the soil, including the texture, structure, porosity, consistence, color, and other physical, mineral, and biological properties of the various horizons, and the thickness and arrangement of those horizons in the soil profile.

Mottling, soil. Irregular spots of different colors that vary in number and size. Descriptive terms are as follows: abundance—*few*, *common*, and *many*; size—*fine*, *medium*, and *coarse*; and contrast—*faint*, *distinct*, and *prominent*. The size measurements are of the diameter along the greatest dimension. *Fine* indicates less than 5 millimeters (about 0.2 inch); *medium*, from 5 to 15 millimeters (about 0.2 to 0.6 inch); and *coarse*, more than 15 millimeters (about 0.6 inch).

Munsell notation. A designation of color by degrees of three simple variables—hue, value, and chroma. For example, a notation of 10YR 6/4 is a color with hue of 10YR, value of 6, and chroma of 4.

Neutral soil. A soil having a pH value of 6.6 to 7.3. (See Reaction, soil.)

Nodules. Cemented bodies lacking visible internal structure. Calcium carbonate, iron oxide, and manganese oxide are common compounds making up nodules. If formed in place, nodules of iron oxide or manganese oxide are considered types of redoximorphic concentrations.

Nose slope. A geomorphic component of hills consisting of the projecting end (laterally convex area) of a hillside. The overland waterflow is predominantly divergent.

Nutrient, plant. An element taken in by a plant essential to its growth. Plant nutrients are mainly nitrogen, phosphorus, potassium, calcium, magnesium, sulfur, iron, manganese, copper, boron, and zinc obtained from the soil and carbon, hydrogen, and oxygen obtained from the air and water.

Organic matter. Plant and animal residue in the soil in various stages of decomposition. The content of organic matter in the surface layer is described as follows:

Very low	less than 0.5 percent
Low	0.5 to 1.0 percent
Moderately low	1.0 to 2.0 percent
Moderate	2.0 to 4.0 percent
High	4.0 to 8.0 percent
Very high	more than 8.0 percent

Outwash plain. A landform of mainly sandy or coarse textured material of glaciofluvial origin. An outwash plain is commonly smooth; where pitted, it generally is low in relief.

Parent material. The unconsolidated organic and mineral material in which soil forms.

Ped. An individual natural soil aggregate, such as a granule, a prism, or a block.

Pedisediment. A thin layer of alluvial material that mantles an erosion surface and has been transported to its present position from higher lying areas of the erosion surface.

Pedon. The smallest volume that can be called “a soil.” A pedon is three dimensional and large enough to permit study of all horizons. Its area ranges from about 10 to 100 square feet (1 square meter to 10 square meters), depending on the variability of the soil.

Percolation. The movement of water through the soil.

Permeability. The quality of the soil that enables water or air to move downward through the profile. The rate at which a saturated soil transmits water is accepted as a measure of this quality. In soil physics, the rate is referred to as “saturated hydraulic conductivity,” which is defined in the “Soil Survey Manual.” In line with conventional usage in the engineering profession and with traditional usage in published soil surveys, this rate of flow continues to be expressed as “permeability.” Terms describing permeability, measured in inches per hour, are as follows:

Impermeable	less than 0.0015 inch
Very slow	0.0015 to 0.06 inch
Slow	0.06 to 0.2 inch
Moderately slow	0.2 to 0.6 inch
Moderate	0.6 inch to 2.0 inches
Moderately rapid	2.0 to 6.0 inches
Rapid	6.0 to 20 inches
Very rapid	more than 20 inches

pH value. A numerical designation of acidity and alkalinity in soil. (See Reaction, soil.)

Phase, soil. A subdivision of a soil series based on features that affect its use and management, such as slope, stoniness, and flooding.

Piping (in tables). Formation of subsurface tunnels or pipelike cavities by water moving through the soil.

Plasticity index. The numerical difference between the liquid limit and the plastic limit; the range of moisture content within which the soil remains plastic.

Plastic limit. The moisture content at which a soil changes from semisolid to plastic.

Plowpan. A compacted layer formed in the soil directly below the plowed layer.

Ponding. Standing water on soils in closed depressions. Unless the soils are artificially drained, the water can be removed only by percolation or evapotranspiration.

Poorly graded. Refers to a coarse grained soil or soil material consisting mainly of particles of nearly the same size. Because there is little difference in size of the particles, density can be increased only slightly by compaction.

Potential native plant community. See Climax plant community.

Potential rooting depth (effective rooting depth). Depth to which roots could penetrate if the content of moisture in the soil were adequate. The soil has no properties restricting the penetration of roots to this depth.

Prescribed burning. Deliberately burning an area for specific management purposes, under the appropriate conditions of weather and soil moisture and at the proper time of day.

Productivity, soil. The capability of a soil for producing a specified plant or sequence of plants under specific management.

Profile, soil. A vertical section of the soil extending through all its horizons and into the parent material.

Proper grazing use. Grazing at an intensity that maintains enough cover to protect the soil and maintain or improve the quantity and quality of the desirable vegetation. This practice increases the vigor and reproduction capacity of the key plants and promotes the accumulation of litter and mulch necessary to conserve soil and water.

Reaction, soil. A measure of acidity or alkalinity of a soil, expressed in pH values. A soil that tests to pH 7.0 is described as precisely neutral in reaction because it is neither acid nor alkaline. The

degrees of acidity or alkalinity, expressed as pH values, are:

Ultra acid	less than 3.5
Extremely acid	3.5 to 4.4
Very strongly acid	4.5 to 5.0
Strongly acid	5.1 to 5.5
Moderately acid	5.6 to 6.0
Slightly acid	6.1 to 6.5
Neutral	6.6 to 7.3
Slightly alkaline	7.4 to 7.8
Moderately alkaline	7.9 to 8.4
Strongly alkaline	8.5 to 9.0
Very strongly alkaline	9.1 and higher

Redoximorphic concentrations. Nodules, concretions, soft masses, pore linings, and other features resulting from the accumulation of iron or manganese oxide. An indication of chemical reduction and oxidation resulting from saturation.

Redoximorphic depletions. Low-chroma zones from which iron and manganese oxide or a combination of iron and manganese oxide and clay has been removed. These zones are indications of the chemical reduction of iron resulting from saturation.

Redoximorphic features. Redoximorphic concentrations, redoximorphic depletions, reduced matrices, a positive reaction to alpha,alpha-dipyridyl, and other features indicating the chemical reduction and oxidation of iron and manganese compounds resulting from saturation.

Reduced matrix. A soil matrix that has low chroma in situ because of chemically reduced iron (Fe II). The chemical reduction results from nearly continuous wetness. The matrix undergoes a change in hue or chroma within 30 minutes after exposure to air as the iron is oxidized (Fe III). A type of redoximorphic feature.

Regolith. The unconsolidated mantle of weathered rock and soil material on the earth's surface; the loose earth material above the solid rock.

Relief. The elevations or inequalities of a land surface, considered collectively.

Residuum (residual soil material). Unconsolidated, weathered or partly weathered mineral material that accumulated as consolidated rock disintegrated in place.

Rill. A steep-sided channel resulting from accelerated erosion. A rill generally is a few inches deep and not wide enough to be an obstacle to farm machinery.

Rise. A slight increase in elevation of the land surface, typically with a broad summit and gently sloping sides.

- Riser.** The relatively short, steeply sloping area below a terrace tread that grades to a lower terrace tread or a base level.
- Road cut.** A sloping surface produced by mechanical means during road construction. It is commonly on the uphill side of the road.
- Rock fragments.** Rock or mineral fragments having a diameter of 2 millimeters or more; for example, pebbles, cobbles, stones, and boulders.
- Rock outcrop.** Exposures of bare bedrock other than rock-lined pits.
- Root zone.** The part of the soil that can be penetrated by plant roots.
- Runoff.** The precipitation discharged into stream channels from an area. The water that flows off the surface of the land without sinking into the soil is called surface runoff. Water that enters the soil before reaching surface streams is called ground-water runoff or seepage flow from ground water.
- Sand.** As a soil separate, individual rock or mineral fragments from 0.05 millimeter to 2.0 millimeters in diameter. Most sand grains consist of quartz. As a soil textural class, a soil that is 85 percent or more sand and not more than 10 percent clay.
- Sandstone.** Sedimentary rock containing dominantly sand-sized particles.
- Saturation.** Wetness characterized by zero or positive pressure of the soil water. Under conditions of saturation, the water will flow from the soil matrix into an unlined auger hole.
- Sawtimber.** Hardwood trees more than 11 inches and conifers more than 9 inches in diameter at breast height.
- Second bottom.** The first terrace above the normal flood plain (or first bottom) of a river.
- Sedimentary rock.** Rock made up of particles deposited from suspension in water. The chief kinds of sedimentary rock are conglomerate, formed from gravel; sandstone, formed from sand; shale, formed from clay; and limestone, formed from soft masses of calcium carbonate. There are many intermediate types. Some wind-deposited sand is consolidated into sandstone.
- Sequum.** A sequence consisting of an illuvial horizon and the overlying eluvial horizon. (See Eluviation.)
- Series, soil.** A group of soils that have profiles that are almost alike, except for differences in texture of the surface layer. All the soils of a series have horizons that are similar in composition, thickness, and arrangement.
- Shale.** Sedimentary rock formed by the hardening of a clay deposit.
- Sheet erosion.** The removal of a fairly uniform layer of soil material from the land surface by the action of rainfall and surface runoff.
- Shoulder.** The position that forms the uppermost inclined surface near the top of a hillslope. It is a transition from backslope to summit. The surface is dominantly convex in profile and erosional in origin.
- Shrink-swell** (in tables). The shrinking of soil when dry and the swelling when wet. Shrinking and swelling can damage roads, dams, building foundations, and other structures. It can also damage plant roots.
- Side slope.** A geomorphic component of hills consisting of a laterally planar area of a hillside. The overland waterflow is predominantly parallel.
- Silica.** A combination of silicon and oxygen. The mineral form is called quartz.
- Silt.** As a soil separate, individual mineral particles that range in diameter from the upper limit of clay (0.002 millimeter) to the lower limit of very fine sand (0.05 millimeter). As a soil textural class, soil that is 80 percent or more silt and less than 12 percent clay.
- Siltstone.** Sedimentary rock made up of dominantly silt-sized particles.
- Similar soils.** Soils that share limits of diagnostic criteria, behave and perform in a similar manner, and have similar conservation needs or management requirements for the major land uses in the survey area.
- Site index.** A designation of the quality of a forest site based on the height of the dominant stand at an arbitrarily chosen age. For example, if the average height attained by dominant and codominant trees in a fully stocked stand at the age of 50 years is 75 feet, the site index is 75.
- Slickensides.** Polished and grooved surfaces produced by one mass sliding past another. In soils, slickensides may occur at the bases of slip surfaces on the steeper slopes; on faces of blocks, prisms, and columns; and in swelling clayey soils, where there is marked change in moisture content.
- Slope.** The inclination of the land surface from the horizontal. Percentage of slope is the vertical distance divided by horizontal distance, then multiplied by 100. Thus, a slope of 20 percent is a drop of 20 feet in 100 feet of horizontal distance.
- Slow refill** (in tables). The slow filling of ponds, resulting from restricted permeability in the soil.
- Soft bedrock.** Bedrock that can be excavated with trenching machines, backhoes, small rippers, and other equipment commonly used in construction.

Soil. A natural, three-dimensional body at the earth's surface. It is capable of supporting plants and has properties resulting from the integrated effect of climate and living matter acting on earthy parent material, as conditioned by relief over periods of time.

Soil separates. Mineral particles less than 2 millimeters in equivalent diameter and ranging between specified size limits. The names and sizes, in millimeters, of separates recognized in the United States are as follows:

Very coarse sand	2.0 to 1.0
Coarse sand	1.0 to 0.5
Medium sand	0.5 to 0.25
Fine sand	0.25 to 0.10
Very fine sand	0.10 to 0.05
Silt	0.05 to 0.002
Clay	less than 0.002

Solum. The upper part of a soil profile, above the C horizon, in which the processes of soil formation are active. The solum in soil consists of the A, E, and B horizons. Generally, the characteristics of the material in these horizons are unlike those of the material below the solum. The living roots and plant and animal activities are largely confined to the solum.

Stone line. A concentration of coarse fragments in a soil. Generally, it is indicative of an old weathered surface. In a cross section, the line may be one fragment or more thick. It generally overlies material that weathered in place and is overlain by recent sediment of variable thickness.

Stones. Rock fragments 10 to 24 inches (25 to 60 centimeters) in diameter if rounded or 15 to 24 inches (38 to 60 centimeters) in length if flat.

Stony. Refers to a soil containing stones in numbers that interfere with or prevent tillage.

Steam terrace. One of a series of platforms in a stream valley, flanking and more or less parallel to the stream channel. It is originally formed near the level of the stream and consists of the dissected remnants of an abandoned flood plain, streambed, or valley floor that were produced during a former stage of erosion or deposition.

Structure, soil. The arrangement of primary soil particles into compound particles or aggregates. The principal forms of soil structure are—*platy* (laminated), *prismatic* (vertical axis of aggregates longer than horizontal), *columnar* (prisms with rounded tops), *blocky* (angular or subangular), and *granular*. *Structureless* soils are either *single grain* (each grain by itself, as in dune sand) or *massive* (the particles adhering without any regular cleavage, as in many hardpans).

Stubble mulch. Stubble or other crop residue left on the soil or partly worked into the soil. It protects the soil from wind erosion and water erosion after harvest, during preparation of a seedbed for the next crop, and during the early growing period of the new crop.

Subsoil. Technically, the B horizon; roughly, the part of the solum below plow depth.

Subsoiling. Tilling a soil below normal plow depth, ordinarily to shatter a hardpan or claypan.

Substratum. The part of the soil below the solum.

Subsurface layer. Any surface soil horizon (A, E, AB, or EB) below the surface layer.

Summit. The topographically highest position of a hillslope. It has a nearly level (planar or only slightly convex) surface.

Surface layer. The soil ordinarily moved in tillage, or its equivalent in uncultivated soil, ranging in depth from 4 to 10 inches (10 to 25 centimeters). Frequently designated as the “plow layer,” or the “Ap horizon.”

Surface soil. The A, E, AB, and EB horizons, considered collectively. It includes all subdivisions of these horizons.

Swale. A slight depression in the midst of generally level land. On an undulating ground moraine, a shallow depression resulting from uneven glacial deposition.

Taxadjuncts. Soils that cannot be classified in a series recognized in the classification system. Such soils are named for a series they strongly resemble and are designated as taxadjuncts to that series because they differ in ways too small to be of consequence in interpreting their use and behavior. Soils are recognized as taxadjuncts only when one or more of their characteristics are slightly outside the range defined for the family of the series for which the soils are named.

Terminal moraine. A belt of thick glacial drift that generally marks the termination of important glacial advances.

Terrace. An embankment, or ridge, constructed across sloping soils on the contour or at a slight angle to the contour. The terrace intercepts surface runoff so that water soaks into the soil or flows slowly to a prepared outlet. A terrace in a field generally is built so that the field can be farmed. A terrace intended mainly for drainage has a deep channel that is maintained in permanent sod.

Terrace (geologic). An old alluvial plain, ordinarily flat or undulating, bordering a river, a lake, or the sea.

Texture, soil. The relative proportions of sand, silt, and clay particles in a mass of soil. The basic

textural classes, in order of increasing proportion of fine particles, are *sand*, *loamy sand*, *sandy loam*, *loam*, *silt loam*, *silt*, *sandy clay loam*, *clay loam*, *silty clay loam*, *sandy clay*, *silty clay*, and *clay*. The sand, loamy sand, and sandy loam classes may be further divided by specifying "coarse," "fine," or "very fine."

Thin layer (in tables). Otherwise suitable soil material that is too thin for the specified use.

Till. Unsorted, nonstratified glacial drift consisting of clay, silt, sand, and boulders transported and deposited by glacial ice.

Till plain. An extensive area of nearly level to undulating soils underlain by glacial till.

Tilth, soil. The physical condition of the soil as related to tillage, seedbed preparation, seedling emergence, and root penetration.

Toeslope. The position that forms the gently inclined surface at the base of a hillslope. Toeslopes in profile are commonly gentle and linear and are constructional surfaces forming the lower part of a hillslope continuum that grades to valley or closed-depression floors.

Topsoil. The upper part of the soil, which is the most favorable material for plant growth. It is ordinarily rich in organic matter and is used to topdress roadbanks, lawns, and land affected by mining.

Trace elements. Chemical elements, for example, zinc, cobalt, manganese, copper, and iron, in soils in extremely small amounts. They are essential to plant growth.

Tread. The relatively flat terrace surface that was cut or built by stream or wave action.

Upland. Land at a higher elevation, in general, than

the alluvial plain or stream terrace; land above the lowlands along streams.

Valley fill. In glaciated regions, material deposited in stream valleys by glacial meltwater. In nonglaciated regions, alluvium deposited by heavily loaded streams.

Variagation. Refers to patterns of contrasting colors assumed to be inherited from the parent material rather than to be the result of poor drainage.

Water bars. Smooth, shallow ditches or depression areas that are excavated at an angle across a sloping road. They are used to reduce the downward velocity of water and divert it off and away from the road surface. Water bars can easily be driven over if constructed properly.

Weathering. All physical and chemical changes produced in rocks or other deposits at or near the earth's surface by atmospheric agents. These changes result in disintegration and decomposition of the material.

Well graded. Refers to soil material consisting of coarse grained particles that are well distributed over a wide range in size or diameter. Such soil normally can be easily increased in density and bearing properties by compaction. Contrasts with poorly graded soil.

Wilting point (or permanent wilting point). The moisture content of soil, on an oven-dry basis, at which a plant (specifically a sunflower) wilts so much that it does not recover when placed in a humid, dark chamber.

Windthrow. The uprooting and tipping over of trees by the wind.

Tables

Table 1.--Temperature and Precipitation
(Recorded in the period 1970-99 at Springfield, Illinois)

Month	Temperature						Precipitation				
	Average daily maximum	Average daily minimum	Average daily	2 years in 10 will have--		Average number of growing degree days*	Average	2 years in 10 will have--		Average number of days with 0.10 inch or more	Average snowfall
				Maximum temperature higher than--	Minimum temperature lower than--			Less than--	More than--		
°F	°F	°F	°F	°F	Units	In	In	In	In		
January-----	32.8	16.8	24.8	62	-15	1	1.62	0.63	2.44	4	7.4
February-----	38.7	21.8	30.2	68	-12	5	1.78	.94	2.52	4	5.9
March-----	50.9	32.2	41.5	81	7	49	3.13	1.89	4.24	6	3.4
April-----	63.8	42.4	53.1	86	22	169	3.60	1.87	5.11	6	.8
May-----	74.8	52.7	63.7	91	35	427	4.09	1.90	5.97	7	.0
June-----	83.7	61.9	72.8	97	46	684	3.68	1.59	5.46	6	.0
July-----	87.0	66.0	76.5	99	51	820	3.51	1.78	5.01	5	.0
August-----	84.7	63.8	74.3	97	49	755	3.44	1.66	4.99	5	.0
September---	78.6	55.5	67.1	95	35	517	3.03	1.36	4.65	5	.0
October-----	66.7	44.3	55.5	88	24	220	2.62	1.47	3.64	5	.0
November----	51.2	33.8	42.5	76	11	47	2.79	1.26	4.11	5	1.6
December----	38.6	23.2	30.9	67	-8	6	2.56	1.22	3.71	5	5.0
Yearly:											
Average---	62.6	42.9	52.7	---	---	---	---	---	---	---	---
Extreme---	102	-21	---	100	-18	---	---	---	---	---	---
Total-----	---	---	---	---	---	3,700	35.85	29.40	41.79	63	24.1

* A growing degree day is a unit of heat available for plant growth. It can be calculated by adding the maximum and minimum daily temperatures, dividing the sum by 2, and subtracting the temperature below which growth is minimal for the principal crops in the area (50 degrees F).

Table 2.--Freeze Dates in Spring and Fall
(Recorded in the period 1970-99 at Springfield, Illinois)

Probability	Temperature		
	24 °F or lower	28 °F or lower	32 °F or lower
Last freezing temperature in spring:			
1 year in 10 later than-----	April 12	April 15	April 23
2 years in 10 later than-----	April 7	April 11	April 19
5 years in 10 later than-----	March 27	April 3	April 12
First freezing temperature in fall:			
1 year in 10 earlier than---	Oct. 20	Oct. 12	Oct. 1
2 years in 10 earlier than---	Oct. 26	Oct. 18	Oct. 5
5 years in 10 earlier than---	Nov. 6	Oct. 29	Oct. 13

Table 3.--Growing Season

(Recorded in the period 1970-99 at Springfield, Illinois)

Probability	Daily minimum temperature during growing season		
	Higher than 24 °F	Higher than 28 °F	Higher than 32 °F
	<i>Days</i>	<i>Days</i>	<i>Days</i>
9 years in 10	198	185	169
8 years in 10	207	193	174
5 years in 10	222	207	184
2 years in 10	238	222	193
1 year in 10	247	230	198

Table 4.--Classification of the Soils

(An asterisk in the first column indicates a taxadjunct to the series. See text for a description of those characteristics that are outside the range of the series.)

Soil name	Family or higher taxonomic class
Alvin-----	Coarse-loamy, mixed, superactive, mesic Typic Hapludalfs
*Assumption----	Fine-silty, mixed, superactive, mesic Oxyaquic Argiudolls
Blackberry----	Fine-silty, mixed, superactive, mesic Oxyaquic Argiudolls
*Broadwell----	Fine-silty, mixed, superactive, mesic Typic Argiudolls
Buckhart-----	Fine-silty, mixed, superactive, mesic Oxyaquic Argiudolls
Camden-----	Fine-silty, mixed, superactive, mesic Typic Hapludalfs
Clarksdale----	Fine, smectitic, mesic Udollic Endoaqualfs
Cowden-----	Fine, smectitic, mesic Mollic Albaqualfs
Denny-----	Fine, smectitic, mesic Mollic Albaqualfs
Drury-----	Fine-silty, mixed, superactive, mesic Dystric Eutrudepts
Edinburg-----	Fine, smectitic, mesic Vertic Argiaquolls
Elburn-----	Fine-silty, mixed, superactive, mesic Aquic Argiudolls
Elco-----	Fine-silty, mixed, superactive, mesic Oxyaquic Hapludalfs
*Elkhart-----	Fine-silty, mixed, superactive, mesic Typic Argiudolls
Fayette-----	Fine-silty, mixed, superactive, mesic Typic Hapludalfs
*Harrison-----	Fine-silty, mixed, superactive, mesic Oxyaquic Argiudolls
Hartsburg-----	Fine-silty, mixed, superactive, mesic Typic Endoaquolls
Herrick-----	Fine, smectitic, mesic Aquic Argiudolls
Hickory-----	Fine-loamy, mixed, active, mesic Typic Hapludalfs
Huntsville----	Fine-silty, mixed, superactive, mesic Cumulic Hapludolls
Ipava-----	Fine, smectitic, mesic Aquic Argiudolls
Kendall-----	Fine-silty, mixed, superactive, mesic Aeric Endoaqualfs
Keomah-----	Fine, smectitic, mesic Aeric Endoaqualfs
Lawson-----	Fine-silty, mixed, superactive, mesic Aquic Cumulic Hapludolls
Marseilles----	Fine-silty, mixed, active, mesic Typic Hapludalfs
Middletown----	Fine-silty, mixed, superactive, mesic Typic Hapludalfs
Navlys-----	Fine-silty, mixed, superactive, mesic Typic Hapludalfs
Orthents-----	Fine-silty, mixed, nonacid, mesic Typic Udorthents
*Osco-----	Fine-silty, mixed, superactive, mesic Typic Argiudolls
Plano-----	Fine-silty, mixed, superactive, mesic Typic Argiudolls
Proctor-----	Fine-silty, mixed, superactive, mesic Typic Argiudolls
Radford-----	Fine-silty, mixed, superactive, mesic Fluvaquentic Hapludolls
Ross-----	Fine-loamy, mixed, superactive, mesic Cumulic Hapludolls
Rozetta-----	Fine-silty, mixed, superactive, mesic Typic Hapludalfs
Sable-----	Fine-silty, mixed, superactive, mesic Typic Endoaquolls
Sawmill-----	Fine-silty, mixed, superactive, mesic Cumulic Endoaquolls
Shiloh-----	Fine, smectitic, mesic Cumulic Vertic Endoaquolls
Spaulding----	Fine-silty, mixed, superactive, mesic Typic Calciaquolls
*Tama-----	Fine-silty, mixed, superactive, mesic Typic Argiudolls
Thebes-----	Fine-silty, mixed, superactive, mesic Typic Hapludalfs
Tice-----	Fine-silty, mixed, superactive, mesic Fluvaquentic Hapludolls
Vesser-----	Fine-silty, mixed, superactive, mesic Argiaquic Argialbolls
Virden-----	Fine, smectitic, mesic Vertic Argiaquolls
Worthen-----	Fine-silty, mixed, superactive, mesic Cumulic Hapludolls
Zook-----	Fine, smectitic, mesic Cumulic Vertic Endoaquolls

Table 5.--Acreage and Proportionate Extent of the Soils

Map symbol	Soil name	Acres	Percent
8D	Hickory silt loam, 10 to 18 percent slopes-----	884	0.2
8D2	Hickory loam, 10 to 18 percent slopes, eroded-----	2,184	0.4
8D3	Hickory clay loam, 10 to 18 percent slopes, severely eroded-----	1,415	0.3
8F	Hickory silt loam, 18 to 35 percent slopes-----	5,181	0.9
17A	Keomah silt loam, 0 to 2 percent slopes-----	9,411	1.7
36B	Tama silt loam, 2 to 5 percent slopes-----	1,424	0.3
36C2	Tama silt loam, 5 to 10 percent slopes, eroded-----	1,408	0.3
43A	Ipava silt loam, 0 to 2 percent slopes-----	153,523	27.4
45A	Denny silt loam, 0 to 2 percent slopes-----	1,674	0.3
46A	Herrick silt loam, 0 to 2 percent slopes-----	5,707	1.0
50A	Viriden silty clay loam, 0 to 2 percent slopes-----	35,240	6.3
68A	Sable silty clay loam, 0 to 2 percent slopes-----	62,026	11.1
86B	Osco silt loam, 2 to 5 percent slopes-----	62,473	11.1
86C2	Osco silt loam, 5 to 10 percent slopes, eroded-----	12,550	2.2
112A	Cowden silt loam, 0 to 2 percent slopes-----	1,899	0.3
119C2	Elco silt loam, 5 to 10 percent slopes, eroded-----	1,067	0.2
119D	Elco silt loam, 10 to 18 percent slopes-----	2,375	0.4
119D2	Elco silt loam, 10 to 18 percent slopes, eroded-----	5,683	1.0
119D3	Elco silty clay loam, 10 to 18 percent slopes, severely eroded-----	4,829	0.9
127B	Harrison silt loam, 2 to 5 percent slopes-----	3,740	0.7
127C2	Harrison silt loam, 5 to 10 percent slopes, eroded-----	5,740	1.0
131C2	Alvin fine sandy loam, 5 to 10 percent slopes, eroded-----	1,037	0.2
131D2	Alvin fine sandy loam, 10 to 18 percent slopes, eroded-----	1,640	0.3
134A	Camden silt loam, 0 to 2 percent slopes-----	139	*
134B	Camden silt loam, 2 to 5 percent slopes-----	405	*
134C2	Camden silt loam, 5 to 10 percent slopes, eroded-----	303	*
138A	Shiloh silty clay loam, 0 to 2 percent slopes-----	450	*
198A	Elburn silt loam, 0 to 2 percent slopes-----	89	*
199A	Plano silt loam, 0 to 2 percent slopes-----	85	*
199B	Plano silt loam, 2 to 5 percent slopes-----	56	*
212C2	Thebes silt loam, 5 to 10 percent slopes, eroded-----	1,079	0.2
242A	Kendall silt loam, 0 to 2 percent slopes-----	254	*
244A	Hartsburg silty clay loam, 0 to 2 percent slopes-----	21,444	3.8
249A	Edinburg silty clay loam, 0 to 2 percent slopes-----	1,183	0.2
257A	Clarksdale silt loam, 0 to 2 percent slopes-----	3,790	0.7
259C2	Assumption silt loam, 5 to 10 percent slopes, eroded-----	6,205	1.1
259D2	Assumption silt loam, 10 to 18 percent slopes, eroded-----	5,796	1.0
279B	Rozetta silt loam, 2 to 5 percent slopes-----	27,929	5.0
280C2	Fayette silt loam, 5 to 10 percent slopes, eroded-----	11,627	2.1
280D2	Fayette silt loam, 10 to 18 percent slopes, eroded-----	1,566	0.3
280D3	Fayette silty clay loam, 10 to 18 percent slopes, severely eroded---	392	*
533	Urban land-----	1,908	0.3
536	Dumps, mine-----	134	*
549G	Marseilles silt loam, 35 to 60 percent slopes-----	633	0.1
567C2	Elkhart silt loam, 5 to 10 percent slopes, eroded-----	6,727	1.2
567D2	Elkhart silt loam, 10 to 18 percent slopes, eroded-----	927	0.2
630C2	Navlys silt loam, 5 to 10 percent slopes, eroded-----	2,636	0.5
630D2	Navlys silt loam, 10 to 18 percent slopes, eroded-----	151	*
630D3	Navlys silty clay loam, 10 to 18 percent slopes, severely eroded---	676	0.1
679B	Blackberry silt loam, 2 to 5 percent slopes-----	212	*
684B	Broadwell silt loam, 2 to 5 percent slopes-----	875	0.2
684C2	Broadwell silt loam, 5 to 10 percent slopes, eroded-----	335	*
685B	Middletown silt loam, 2 to 5 percent slopes-----	4,276	0.8
685C2	Middletown silt loam, 5 to 10 percent slopes, eroded-----	919	0.2
705A	Buckhart silt loam, 0 to 2 percent slopes-----	7,391	1.3
705B	Buckhart silt loam, 2 to 5 percent slopes-----	4,866	0.9
712A	Spaulding silty clay loam, 0 to 2 percent slopes-----	856	0.2
801C	Orthents, silty, rolling-----	443	*
830	Landfills-----	49	*
862	Pits, sand-----	401	*
864	Pits, quarries-----	17	*
3073A	Ross silt loam, 0 to 2 percent slopes, frequently flooded-----	1,065	0.2

See footnote at end of table.

Table 5.--Acreage and Proportionate Extent of the Soils--Continued

Map symbol	Soil name	Acres	Percent
3074A	Radford silt loam, 0 to 2 percent slopes, frequently flooded-----	12,852	2.3
3077A	Huntsville silt loam, 0 to 2 percent slopes, frequently flooded-----	324	*
3107A	Sawmill silty clay loam, 0 to 2 percent slopes, frequently flooded--	12,613	2.2
3284A	Tice silty clay loam, 0 to 2 percent slopes, frequently flooded-----	10,192	1.8
3405A	Zook silty clay loam, 0 to 2 percent slopes, frequently flooded-----	511	*
3451A	Lawson silt loam, 0 to 2 percent slopes, frequently flooded-----	8,740	1.6
7037A	Worthen silt loam, 0 to 2 percent slopes, rarely flooded-----	534	*
7075B	Drury silt loam, 2 to 5 percent slopes, rarely flooded-----	1,028	0.2
7148A	Proctor silt loam, 0 to 2 percent slopes, rarely flooded-----	3,195	0.6
7242A	Kendall silt loam, 0 to 2 percent slopes, rarely flooded-----	906	0.2
8396A	Vesser silt loam, 0 to 2 percent slopes, occasionally flooded-----	911	0.2
	Water-----	7,550	1.3
	Total-----	560,755	100.0

* Less than 0.1 percent.

Table 6.--Limitations and Hazards Affecting Cropland and Pasture

(See text for a description of the limitations and hazards listed in this table. Miscellaneous areas and map units generally not available for crop or pasture production are excluded from the table. Absence of an entry indicates that the map unit is generally unsuited to crops or pasture.)

Soil name and map symbol	Limitations and hazards affecting cropland	Limitations and hazards affecting pasture
8D: Hickory-----	Crusting, water erosion	Low pH, water erosion
8D2: Hickory-----	Crusting, water erosion	Low pH, water erosion
8D3: Hickory-----	Poor tilth, crusting, water erosion	Low fertility, low pH, water erosion
8F: Hickory-----	---	Equipment limitation, low pH, water erosion
17A: Keomah-----	Wetness, crusting	Wetness, low pH
36B: Tama-----	Water erosion	Low pH
36C2: Tama-----	Crusting, water erosion	Low pH, water erosion
43A: Ipava-----	Wetness	NA*
45A: Denny-----	Ponding, crusting	NA*
46A: Herrick-----	Wetness	NA*
50A: Virden-----	Ponding, poor tilth	NA*
68A: Sable-----	Ponding, poor tilth	NA*
86B: Osco-----	Water erosion	Low pH
86C2: Osco-----	Crusting, water erosion	Low pH, water erosion
112A: Cowden-----	Ponding, crusting	Frost heave, low pH, ponding
119C2: Elco-----	Crusting, water erosion	Low pH, water erosion
119D: Elco-----	Crusting, water erosion	Low pH, water erosion
119D2: Elco-----	Crusting, water erosion	Low pH, water erosion

See footnotes at end of table.

Table 6.--Limitations and Hazards Affecting Cropland and Pasture--Continued

Soil name and map symbol	Limitations and hazards affecting cropland	Limitations and hazards affecting pasture
119D3: Elco-----	Poor tilth, crusting, water erosion	Low fertility, low pH, water erosion
127B: Harrison-----	Water erosion	Low pH
127C2: Harrison-----	Crusting, water erosion	Low pH, water erosion
131C2: Alvin-----	Water erosion	Low fertility, low pH, water erosion
131D2: Alvin-----	Water erosion	Low fertility, low pH, water erosion
134A: Camden-----	Crusting	Low pH
134B: Camden-----	Crusting, water erosion	Low pH, water erosion
134C2: Camden-----	Crusting, water erosion	Low pH, water erosion
138A: Shiloh-----	Ponding, poor tilth	NA*
198A: Elburn-----	Wetness	NA*
199A: Plano-----	None**	NA*
199B: Plano-----	Water erosion	None***
212C2: Thebes-----	Crusting, water erosion, excessive permeability	Low pH, water erosion
242A: Kendall-----	Wetness, crusting	Wetness, low pH
244A: Hartsburg-----	High pH, ponding, poor tilth	NA*
249A: Edinburg-----	Ponding, poor tilth	NA*
257A: Clarksdale-----	Wetness, crusting	Wetness, low pH
259C2: Assumption-----	Crusting, water erosion	Low pH, water erosion
259D2: Assumption-----	Crusting, water erosion	Low pH, water erosion
279B: Rozetta-----	Crusting, water erosion	Low pH, water erosion

See footnotes at end of table.

Table 6.--Limitations and Hazards Affecting Cropland and Pasture--Continued

Soil name and map symbol	Limitations and hazards affecting cropland	Limitations and hazards affecting pasture
280C2: Fayette-----	Crusting, water erosion	Low pH, water erosion
280D2: Fayette-----	Crusting, water erosion	Low pH, water erosion
280D3: Fayette-----	Poor tilth, crusting, water erosion	Low fertility, low pH, water erosion
549G: Marseilles.		
567C2: Elkhart-----	Crusting, water erosion, high pH	High pH, water erosion
567D2: Elkhart-----	Crusting, water erosion, high pH	High pH, water erosion
630C2: Navlys-----	Crusting, water erosion, high pH	High pH, water erosion
630D2: Navlys-----	Crusting, water erosion, high pH	High pH, water erosion
630D3: Navlys-----	Poor tilth, crusting, water erosion, high pH	High pH, water erosion, low fertility
679B: Blackberry-----	Water erosion	Low pH
684B: Broadwell-----	Water erosion, excessive permeability	None***
684C2: Broadwell-----	Crusting, water erosion, excessive permeability	Water erosion
685B: Middletown-----	Crusting, water erosion, excessive permeability	Low pH, water erosion
685C2: Middletown-----	Crusting, water erosion, excessive permeability	Low pH, water erosion
705A: Buckhart-----	None**	NA*
705B: Buckhart-----	Water erosion	None***
712A: Spaulding-----	Excess lime, high pH, ponding, poor tilth	NA*

See footnotes at end of table.

Table 6.--Limitations and Hazards Affecting Cropland and Pasture

Soil name and map symbol	Limitations and hazards affecting cropland	Limitations and hazards affecting pasture
3073A: Ross-----	Flooding	Flooding
3074A: Radford-----	Flooding, wetness	Flooding, wetness
3077A: Huntsville-----	Flooding	Flooding
3107A: Sawmill-----	Flooding, ponding, poor tilth	Flooding, ponding, frost heave
3284A: Tice-----	Flooding, wetness, poor tilth	NA*
3405A: Zook-----	Flooding, wetness, poor tilth	NA*
3451A: Lawson-----	Flooding, wetness	Flooding, wetness
7037A: Worthen-----	None**	None***
7075B: Drury-----	Crusting, water erosion	Water erosion
7148A: Proctor-----	None**	NA*
7242A: Kendall-----	Wetness, crusting	NA*
8396A: Vesser-----	Flooding, wetness, crusting	NA*

* Pasture is not a major use.

** This soil is well suited to crops.

*** This soil is well suited to pasture.

Table 7.--Land Capability and Yields per Acre of Crops and Pasture--Continued

(Yields are those that can be expected under a high level of management. They are for nonirrigated areas. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil.)

Map symbol and soil name	Land capability	Corn	Soybeans	Winter wheat	Grass-legume pasture	Grass-legume hay
		Bu	Bu	Bu	AUM	Tons
8D: Hickory-----	3e	74	24	27	4.7	2.8
8D2: Hickory-----	3e	71	23	26	4.5	2.7
8D3: Hickory-----	4e	66	22	24	4.2	2.5
8F: Hickory-----	6e	---	---	---	3.7	2.2
17A: Keomah-----	2w	129	39	52	8.5	5.1
36B: Tama-----	2e	153	46	61	9.7	5.8
36C2: Tama-----	3e	146	43	58	9.2	5.5
43A: Ipava-----	1	163	52	66	10.2	6.1
45A: Denny-----	3w	113	37	---	---	---
46A: Herrick-----	2w	141	45	61	---	---
50A: Virdden-----	2w	138	46	---	---	---
68A: Sable-----	2w	156	51	---	---	---
86B: Osco-----	2e	153	46	61	9.7	5.8
86C2: Osco-----	3e	146	43	58	9.2	5.5
112A: Cowden-----	2w	120	37	53	8.0	4.8
119C2: Elco-----	3e	105	35	44	6.9	4.1
119D: Elco-----	3e	104	34	44	6.8	4.0
119D2: Elco-----	3e	100	33	42	6.5	3.9
119D3: Elco-----	4e	93	31	39	6.1	3.7

Table 7.--Land Capability and Yields per Acre of Crops and Pasture--Continued

Map symbol and soil name	Land capability	Corn	Soybeans	Winter wheat	Grass-legume pasture	Grass-legume hay
		Bu	Bu	Bu	AUM	Tons
127B: Harrison-----	2e	135	42	58	8.7	5.2
127C2: Harrison-----	3e	128	39	55	8.3	5.0
131C2: Alvin-----	3e	93	35	44	6.4	3.9
131D2: Alvin-----	4e	88	33	42	6.0	2.1
134A: Camden-----	1	125	39	55	8.3	5.0
134B: Camden-----	2e	124	39	54	8.2	5.0
134C2: Camden-----	3e	118	37	52	7.8	4.7
138A: Shiloh-----	2w	139	46	---	---	---
198A: Elburn-----	1	161	50	63	---	---
199A: Plano-----	1	151	45	60	---	---
199B: Plano-----	2e	149	45	59	9.5	5.7
212C2: Thebes-----	3e	95	33	43	6.3	3.8
242A: Kendall-----	2w	135	41	55	8.7	5.2
244A: Hartsburg-----	2w	145	47	---	---	---
249A: Edinburg-----	3w	132	43	---	---	---
257A: Clarksdale-----	1	140	43	57	8.8	5.3
259C2: Assumption-----	3e	120	37	53	7.8	4.7
259D2: Assumption-----	4e	115	35	50	7.5	4.5
279B: Rozetta-----	2e	130	40	53	8.6	5.1
280C2: Fayette-----	3e	121	37	50	8.1	4.9
280D2: Fayette-----	3e	116	35	48	7.8	4.7

Table 7.--Land Capability and Yields per Acre of Crops and Pasture--Continued

Map symbol and soil name	Land capability	Corn	Soybeans	Winter wheat	Grass-legume pasture	Grass-legume hay
		Bu	Bu	Bu	AUM	Tons
280D3: Fayette-----	4e	107	32	44	7.2	4.3
533: Urban land.						
536: Dumps.						
549G: Marseilles-----	7e	---	---	---	---	---
567C2: Elkhart-----	3e	124	37	50	8.0	4.8
567D2: Elkhart-----	3e	119	32	48	7.7	4.6
630C2: Navlys-----	3e	105	33	50	7.8	4.7
630D2: Navlys-----	3e	101	32	48	7.5	4.5
630D3: Navlys-----	4e	93	29	44	6.9	4.2
679B: Blackberry-----	2e	150	45	59	9.6	5.7
684B: Broadwell-----	2e	144	44	58	9.2	5.5
684C2: Broadwell-----	3e	136	41	55	8.7	5.3
685B: Middletown-----	2e	116	34	52	7.7	4.7
685C2: Middletown-----	3e	110	32	50	7.3	4.4
705A: Buckhart-----	1	158	48	63	---	---
705B: Buckhart-----	2e	158	48	62	9.8	5.9
712A: Spaulding-----	2w	138	44	---	---	---
801C: Orthents-----	3e	---	---	---	---	---
830: Landfills.						
862: Pits, sand.						
864: Pits, quarries.						

Table 7.--Land Capability and Yields per Acre of Crops and Pasture--Continued

Map symbol and soil name	Land capability	Corn	Soybeans	Winter wheat	Grass-legume pasture	Grass-legume hay
		Bu	Bu	Bu	AUM	Tons
3073A: Ross-----	2w	131	41	54	8.2	5.0
3074A: Radford-----	3w	129	41	55	8.4	5.0
3077A: Huntsville-----	2w	137	43	58	8.7	5.2
3107A: Sawmill-----	3w	132	42	---	8.3	5.0
3284A: Tice-----	3w	138	42	55	---	---
3405A: Zook-----	3w	83	32	---	---	---
3451A: Lawson-----	3w	145	43	56	8.6	5.7
7037A: Worthen-----	1	151	46	62	9.8	5.9
7075B: Drury-----	2e	125	40	56	8.2	4.9
7148A: Proctor-----	1	144	44	59	---	---
7242A: Kendall-----	2w	135	41	55	---	---
8396A: Vesser-----	2w	126	42	---	---	---

Table 8.--Prime Farmland

(Only the soils considered prime farmland are listed. Urban or built-up areas of the soils listed are not considered prime farmland. If a soil is prime farmland only under certain conditions, the conditions are specified in parentheses after the soil name.)

Map symbol	Soil name
17A	Keomah silt loam, 0 to 2 percent slopes (where drained)
36B	Tama silt loam, 2 to 5 percent slopes
43A	Ipava silt loam, 0 to 2 percent slopes
45A	Denny silt loam, 0 to 2 percent slopes (where drained)
46A	Herrick silt loam, 0 to 2 percent slopes
50A	Virден silty clay loam, 0 to 2 percent slopes (where drained)
68A	Sable silty clay loam, 0 to 2 percent slopes (where drained)
86B	Osco silt loam, 2 to 5 percent slopes
112A	Cowden silt loam, 0 to 2 percent slopes (where drained)
127B	Harrison silt loam, 2 to 5 percent slopes
131C2	Alvin fine sandy loam, 5 to 10 percent slopes, eroded
134A	Camden silt loam, 0 to 2 percent slopes
134B	Camden silt loam, 2 to 5 percent slopes
138A	Shiloh silty clay loam, 0 to 2 percent slopes (where drained)
198A	Elburn silt loam, 0 to 2 percent slopes
199A	Plano silt loam, 0 to 2 percent slopes
199B	Plano silt loam, 2 to 5 percent slopes
242A	Kendall silt loam, 0 to 2 percent slopes (where drained)
244A	Hartsburg silty clay loam, 0 to 2 percent slopes (where drained)
249A	Edinburg silty clay loam, 0 to 2 percent slopes (where drained)
257A	Clarksdale silt loam, 0 to 2 percent slopes (where drained)
279B	Rozetta silt loam, 2 to 5 percent slopes
679B	Blackberry silt loam, 2 to 5 percent slopes
684B	Broadwell silt loam, 2 to 5 percent slopes
685B	Middletown silt loam, 2 to 5 percent slopes
705A	Buckhart silt loam, 0 to 2 percent slopes
705B	Buckhart silt loam, 2 to 5 percent slopes
712A	Spaulding silty clay loam, 0 to 2 percent slopes (where drained)
3073A	Ross silt loam, 0 to 2 percent slopes, frequently flooded (where protected from flooding or not frequently flooded during the growing season)
3074A	Radford silt loam, 0 to 2 percent slopes, frequently flooded (where protected from flooding or not frequently flooded during the growing season)
3077A	Huntsville silt loam, 0 to 2 percent slopes, frequently flooded (where protected from flooding or not frequently flooded during the growing season)
3107A	Sawmill silty clay loam, 0 to 2 percent slopes, frequently flooded (where drained and either protected from flooding or not frequently flooded during the growing season)
3284A	Tice silty clay loam, 0 to 2 percent slopes, frequently flooded (where protected from flooding or not frequently flooded during the growing season)
3405A	Zook silty clay loam, 0 to 2 percent slopes, frequently flooded (where drained and either protected from flooding or not frequently flooded during the growing season)
3451A	Lawson silt loam, 0 to 2 percent slopes, frequently flooded (where protected from flooding or not frequently flooded during the growing season)
7037A	Worthen silt loam, 0 to 2 percent slopes, rarely flooded
7075B	Drury silt loam, 2 to 5 percent slopes, rarely flooded
7148A	Proctor silt loam, 0 to 2 percent slopes, rarely flooded
7242A	Kendall silt loam, 0 to 2 percent slopes, rarely flooded (where drained)
8396A	Vesser silt loam, 0 to 2 percent slopes, occasionally flooded (where drained)

Table 9.--Hydric Soils

Map symbol and map unit name	Component	Hydric	Local landform
8D: Hickory silt loam, 10 to 18 percent slopes	Hickory	No	Ground moraines
8D2: Hickory loam, 10 to 18 percent slopes, eroded	Hickory	No	Ground moraines
8D3: Hickory clay loam, 10 to 18 percent slopes, severely eroded	Hickory	No	Ground moraines
8F: Hickory silt loam, 18 to 35 percent slopes	Hickory	No	Ground moraines
17A: Keomah silt loam, 0 to 2 percent slopes	Keomah Denny	No Yes	Ground moraines Depressions
36B: Tama silt loam, 2 to 5 percent slopes	Tama	No	Ground moraines, loessial hills
36C2: Tama silt loam, 5 to 10 percent slopes, eroded	Tama	No	Ground moraines, loessial hills
43A: Ipava silt loam, 0 to 2 percent slopes	Ipava Sable Viriden Denny	No Yes Yes Yes	Ground moraines Ground moraines, depressions Ground moraines Depressions
45A: Denny silt loam, 0 to 2 percent slopes	Denny	Yes	Depressions
46A: Herrick silt loam, 0 to 2 percent slopes	Herrick Viriden Cowden	No Yes Yes	Ground moraines Ground moraines Ground moraines
50A: Viriden silty clay loam, 0 to 2 percent slopes	Viriden	Yes	Ground moraines
68A: Sable silty clay loam, 0 to 2 percent slopes	Sable	Yes	Ground moraines
86B: Osco silt loam, 2 to 5 percent slopes	Osco	No	Ground moraines
86C2: Osco silt loam, 5 to 10 percent slopes, eroded	Osco	No	Ground moraines
112A: Cowden silt loam, 0 to 2 percent slopes	Cowden	Yes	Ground moraines
119C2: Elco silt loam, 5 to 10 percent slopes, eroded	Elco	No	Ground moraines
119D: Elco silt loam, 10 to 18 percent slopes	Elco	No	Ground moraines
119D2: Elco silt loam, 10 to 18 percent slopes, eroded	Elco	No	Ground moraines
119D3: Elco silty clay loam, 10 to 18 percent slopes, severely eroded	Elco	No	Ground moraines

Table 9.--Hydric Soils--Continued

Map symbol and map unit name	Component	Hydric	Local landform
127B: Harrison silt loam, 2 to 5 percent slopes	Harrison	No	Ground moraines
	Virden	Yes	Ground moraines
	Cowden	Yes	Ground moraines
127C2: Harrison silt loam, 5 to 10 percent slopes, eroded	Harrison	No	Ground moraines
131C2: Alvin fine sandy loam, 5 to 10 percent slopes, eroded	Alvin	No	Knolls, outwash plains, stream terraces
131D2: Alvin fine sandy loam, 10 to 18 percent slopes, eroded	Alvin	No	Knolls, outwash plains, stream terraces
134A: Camden silt loam, 0 to 2 percent slopes	Camden	No	Outwash plains, stream terraces
	Vesser	Yes	Flood plains
134B: Camden silt loam, 2 to 5 percent slopes	Camden	No	Outwash plains, stream terraces
134C2: Camden silt loam, 5 to 10 percent slopes, eroded	Camden	No	Stream terraces, outwash plains
138A: Shiloh silty clay loam, 0 to 2 percent slopes	Shiloh	Yes	Ground moraines, depressions
198A: Elburn silt loam, 0 to 2 percent slopes	Elburn	No	Outwash plains, stream terraces
	Vesser	Yes	Flood plains
199A: Plano silt loam, 0 to 2 percent slopes	Plano	No	Outwash plains, stream terraces
	Vesser	Yes	Flood plains
199B: Plano silt loam, 2 to 5 percent slopes	Plano	No	Outwash plains, stream terraces
212C2: Thebes silt loam, 5 to 10 percent slopes, eroded	Thebes	No	Knolls, ground moraines, stream terraces
242A: Kendall silt loam, 0 to 2 percent slopes	Kendall	No	Outwash plains, stream terraces
	Vesser	Yes	Flood plains
244A: Hartsburg silty clay loam, 0 to 2 percent slopes	Hartsburg	Yes	Outwash plains, ground moraines
249A: Edinburg silty clay loam, 0 to 2 percent slopes	Edinburg	Yes	Depressions
257A: Clarksdale silt loam, 0 to 2 percent slopes	Clarksdale	No	Ground moraines
	Denny	Yes	Depressions
259C2: Assumption silt loam, 5 to 10 percent slopes, eroded	Assumption	No	Ground moraines

Table 9.--Hydric Soils--Continued

Map symbol and map unit name	Component	Hydric	Local landform
259D2: Assumption silt loam, 10 to 18 percent slopes, eroded	Assumption	No	Ground moraines
279B: Rozetta silt loam, 2 to 5 percent slopes	Rozetta	No	Ground moraines
280C2: Fayette silt loam, 5 to 10 percent slopes, eroded	Fayette	No	Ground moraines
280D2: Fayette silt loam, 10 to 18 percent slopes, eroded	Fayette	No	Ground moraines
280D3: Fayette silty clay loam, 10 to 18 percent slopes, severely eroded	Fayette	No	Ground moraines
533: Urban land	Urban land	Unranked	---
536: Dumps, mine	Dumps	Unranked	---
549G: Marseilles silt loam, 35 to 60 percent slopes	Marseilles	No	Ground moraines
567C2: Elkhart silt loam, 5 to 10 percent slopes, eroded	Elkhart	No	Ground moraines
567D2: Elkhart silt loam, 10 to 18 percent slopes, eroded	Elkhart	No	Ground moraines
630C2: Navlys silt loam, 5 to 10 percent slopes, eroded	Navlys	No	Ground moraines
630D2: Navlys silt loam, 10 to 18 percent slopes, eroded	Navlys	No	Ground moraines
630D3: Navlys silty clay loam, 10 to 18 percent slopes, severely eroded	Navlys	No	Ground moraines
679B: Blackberry silt loam, 2 to 5 percent slopes	Blackberry	No	Outwash plains, stream terraces
684B: Broadwell silt loam, 2 to 5 percent slopes	Broadwell	No	Ground moraines, knolls, outwash plains
684C2: Broadwell silt loam, 5 to 10 percent slopes, eroded	Broadwell	No	Knolls, ground moraines
685B: Middletown silt loam, 2 to 5 percent slopes	Middletown	No	Knolls, ground moraines
685C2: Middletown silt loam, 5 to 10 percent slopes, eroded	Middletown	No	Knolls, ground moraines

Table 9.--Hydric Soils--Continued

Map symbol and map unit name	Component	Hydric	Local landform
705A: Buckhart silt loam, 0 to 2 percent slopes	Buckhart	No	Knolls, ground moraines
	Sable	Yes	Ground moraines, depressions
	Viriden	Yes	Ground moraines
	Denny	Yes	Depressions
705B: Buckhart silt loam, 2 to 5 percent slopes	Buckhart	No	Knolls, ground moraines
	Sable	Yes	Ground moraines, depressions
	Viriden	Yes	Ground moraines
712A: Spaulding silty clay loam, 0 to 2 percent slopes	Spaulding	Yes	Ground moraines, depressions
	Sable	Yes	Ground moraines, depressions
801C: Orthents, silty, rolling	Orthents	No	Ground moraines
830: Landfills	Landfills	Unranked	---
862: Pits, sand	Pits	Unranked	Stream terraces
864: Pits, quarries	Pits	Unranked	---
3073A: Ross silt loam, 0 to 2 percent slopes, frequently flooded	Ross	No	Flood plains
	Sawmill	Yes	Flood plains
3074A: Radford silt loam, 0 to 2 percent slopes, frequently flooded	Radford	No	Flood plains
	Sawmill	Yes	Flood plains
3077A: Huntsville silt loam, 0 to 2 percent slopes, frequently flooded	Huntsville	No	Flood plains
	Sawmill	Yes	Flood plains
3107A: Sawmill silty clay loam, 0 to 2 percent slopes, frequently flooded	Sawmill	Yes	Flood plains
3284A: Tice silty clay loam, 0 to 2 percent slopes, frequently flooded	Tice	No	Flood plains
	Sawmill	Yes	Flood plains
3405A: Zook silty clay loam, 0 to 2 percent slopes, frequently flooded	Zook	Yes	Flood plains
3451A: Lawson silt loam, 0 to 2 percent slopes, frequently flooded	Lawson	No	Flood plains
	Sawmill	Yes	Flood plains
	Zook	Yes	Flood plains
7037A: Worthen silt loam, 0 to 2 percent slopes, rarely flooded	Worthen	No	Alluvial fans, loessial bluffs
	Vesser	Yes	Flood plains
7075B: Drury silt loam, 2 to 5 percent slopes, rarely flooded	Drury	No	Alluvial fans, stream terraces
	Vesser	Yes	Flood plains

Table 9.--Hydric Soils--Continued

Map symbol and map unit name	Component	Hydric	Local landform
7148A: Proctor silt loam, 0 to 2 percent slopes, rarely flooded	Proctor Sawmill	No Yes	Flood-plain steps Flood plains
7242A: Kendall silt loam, 0 to 2 percent slopes, rarely flooded	Kendall Vesser	No Yes	Flood-plain steps Flood plains
8396A: Vesser silt loam, 0 to 2 percent slopes, occasionally flooded	Vesser	Yes	Flood plains

Table 10.--Windbreaks and Environmental Plantings

(Absence of an entry indicates that trees generally do not grow to the given height on the soil.)

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
8D: Hickory-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
8D2: Hickory-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
8D3: Hickory-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
8F: Hickory-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
17A: Keomah-----	American cranberrybush, black chokeberry, buttonbush, common elderberry, common ninebark, common winterberry, gray dogwood, highbush blueberry, northern spicebush, redosier dogwood, silky dogwood	Cockspur hawthorn, hazel alder, nannyberry, roughleaf dogwood	Arborvitae, blackgum, common hackberry, green hawthorn, northern white-cedar, shingle oak	Green ash, red maple, river birch, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak

Table 10.--Windbreaks and Environmental Plantings--Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
36B: Tama-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
36C2: Tama-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
43A: Ipava-----	American cranberrybush, Canada yew, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, northern spicebush, redosier dogwood, silky dogwood	Blackhaw, cockspur hawthorn, common pawpaw, common serviceberry, prairie crabapple, roughleaf dogwood, rusty blackhaw, southern arrowwood, witchhazel	Austrian pine, Douglas fir, arborvitae, blue spruce, common persimmon, eastern redcedar, green hawthorn, nannyberry, pecan, shingle oak	Norway spruce, blackgum, common hackberry, green ash, red maple, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
45A: Denny-----	American cranberrybush, black chokeberry, buttonbush, common elderberry, common ninebark, common winterberry, gray dogwood, highbush blueberry, northern spicebush, redosier dogwood, silky dogwood	Cockspur hawthorn, hazel alder, nannyberry, roughleaf dogwood	Arborvitae, blackgum, common hackberry, green hawthorn, northern white-cedar, shingle oak	Green ash, red maple, river birch, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
46A: Herrick-----	American cranberrybush, Canada yew, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, northern spicebush, redosier dogwood, silky dogwood	Blackhaw, cockspur hawthorn, common pawpaw, common serviceberry, prairie crabapple, roughleaf dogwood, rusty blackhaw, southern arrowwood, witchhazel	Austrian pine, Douglas fir, arborvitae, blue spruce, common persimmon, eastern redcedar, green hawthorn, nannyberry, pecan, shingle oak	Norway spruce, blackgum, common hackberry, green ash, red maple, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak

Table 10.--Windbreaks and Environmental Plantings--Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
50A: Virden-----	American cranberrybush, black chokeberry, buttonbush, common elderberry, common ninebark, common winterberry, gray dogwood, highbush blueberry, northern spicebush, redosier dogwood, silky dogwood	Cockspur hawthorn, hazel alder, nannyberry, roughleaf dogwood	Arborvitae, blackgum, common hackberry, green hawthorn, northern white-cedar, shingle oak	Green ash, red maple, river birch, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
68A: Sable-----	American cranberrybush, black chokeberry, buttonbush, common elderberry, common ninebark, common winterberry, gray dogwood, highbush blueberry, northern spicebush, redosier dogwood, silky dogwood	Cockspur hawthorn, hazel alder, nannyberry, roughleaf dogwood	Arborvitae, blackgum, common hackberry, green hawthorn, northern white-cedar, shingle oak	Green ash, red maple, river birch, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
86B: Osco-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
86C2: Osco-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine

Table 10.--Windbreaks and Environmental Plantings--Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
112A: Cowden-----	American cranberrybush, black chokeberry, buttonbush, common elderberry, common ninebark, common winterberry, gray dogwood, highbush blueberry, northern spicebush, redosier dogwood, silky dogwood	Cockspur hawthorn, hazel alder, nannyberry, roughleaf dogwood	Arborvitae, blackgum, common hackberry, green hawthorn, northern white-cedar, shingle oak	Green ash, red maple, river birch, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
119C2: Elco-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
119D: Elco-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
119D2: Elco-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
119D3: Elco-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine

Table 10.--Windbreaks and Environmental Plantings--Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
127B: Harrison----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
127C2: Harrison----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
131C2: Alvin-----	American hazelnut, black chokeberry, common winterberry, coralberry, gray dogwood, mapleleaf viburnum	American plum, American witchhazel, Arnold hawthorn, blackhaw, common chokecherry, common serviceberry, prairie crabapple	Douglas fir, arborvitae, black walnut, blackgum, blue spruce, bur oak, eastern redcedar, green ash, pecan	Norway spruce, common hackberry, pin oak, tuliptree	Carolina poplar, eastern white pine
131D2: Alvin-----	American hazelnut, black chokeberry, common winterberry, coralberry, gray dogwood, mapleleaf viburnum	American plum, American witchhazel, Arnold hawthorn, blackhaw, common chokecherry, common serviceberry, prairie crabapple	Douglas fir, arborvitae, black walnut, blackgum, blue spruce, bur oak, eastern redcedar, green ash, pecan	Norway spruce, common hackberry, pin oak, tuliptree	Carolina poplar, eastern white pine
134A: Camden-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine

Table 10.--Windbreaks and Environmental Plantings--Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
134B: Camden-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
134C2: Camden-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
138A: Shiloh-----	American cranberrybush, black chokeberry, buttonbush, common elderberry, common ninebark, common winterberry, gray dogwood, highbush blueberry, northern spicebush, redosier dogwood, silky dogwood	Cockspur hawthorn, hazel alder, nannyberry, roughleaf dogwood	Arborvitae, blackgum, common hackberry, green hawthorn, northern white-cedar, shingle oak	Green ash, red maple, river birch, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
198A: Elburn-----	American cranberrybush, Canada yew, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, northern spicebush, redosier dogwood, silky dogwood	Blackhaw, cockspur hawthorn, common pawpaw, common serviceberry, prairie crabapple, roughleaf dogwood, rusty blackhaw, southern arrowwood, witchhazel	Austrian pine, Douglas fir, arborvitae, blue spruce, common persimmon, eastern redcedar, green hawthorn, nannyberry, pecan, shingle oak	Norway spruce, blackgum, common hackberry, green ash, red maple, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
199A: Plano-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine

Table 10.--Windbreaks and Environmental Plantings--Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
199B: Plano-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
212C2: Thebes-----	American cranberrybush, American hazelnut, black chokeberry, common chokecherry, common elderberry, common juniper, coralberry, mapleleaf viburnum, silky dogwood	American plum, bur oak, chinkapin oak, common serviceberry, eastern redcedar, nannyberry, prairie crabapple, roughleaf dogwood, smooth sumac	Black oak, common hackberry, eastern white pine, green ash	Carolina poplar	---
242A: Kendall-----	American cranberrybush, Canada yew, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, northern spicebush, redosier dogwood, silky dogwood	Blackhaw, cockspur hawthorn, common pawpaw, common serviceberry, prairie crabapple, roughleaf dogwood, rusty blackhaw, southern arrowwood, witchhazel	Austrian pine, Douglas fir, arborvitae, blue spruce, common persimmon, eastern redcedar, green hawthorn, nannyberry, pecan, shingle oak	Norway spruce, blackgum, common hackberry, green ash, red maple, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
244A: Hartsburg---	American cranberrybush, black chokeberry, buttonbush, common elderberry, common ninebark, common winterberry, gray dogwood, highbush blueberry, northern spicebush, redosier dogwood, silky dogwood	Cockspur hawthorn, hazel alder, nannyberry, roughleaf dogwood	Arborvitae, blackgum, common hackberry, green hawthorn, northern white-cedar, shingle oak	Green ash, red maple, river birch, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
249A: Edinburg----	American cranberrybush, black chokeberry, buttonbush, common elderberry, common ninebark, common winterberry, gray dogwood, highbush blueberry, northern spicebush, redosier dogwood, silky dogwood	Cockspur hawthorn, hazel alder, nannyberry, roughleaf dogwood	Arborvitae, blackgum, common hackberry, green hawthorn, northern white-cedar, shingle oak	Green ash, red maple, river birch, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak

Table 10.--Windbreaks and Environmental Plantings--Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
257A: Clarksdale--	American cranberrybush, Canada yew, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, northern spicebush, redosier dogwood, silky dogwood	Blackhaw, cockspur hawthorn, common pawpaw, common serviceberry, prairie crabapple, roughleaf dogwood, rusty blackhaw, southern arrowwood, witchhazel	Austrian pine, Douglas fir, arborvitae, blue spruce, common persimmon, eastern redcedar, green hawthorn, nannyberry, pecan, shingle oak	Norway spruce, blackgum, common hackberry, green ash, red maple, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
259C2: Assumption--	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
259D2: Assumption--	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
279B: Rozetta----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
280C2: Fayette----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine

Table 10.--Windbreaks and Environmental Plantings--Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
280D2: Fayette-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
280D3: Fayette-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
533: Urban land.					
536: Dumps.					
549G: Marseilles--	American cranberrybush, American hazelnut, black chokeberry, common chokecherry, common elderberry, common juniper, coralberry, mapleleaf viburnum, silky dogwood	American plum, bur oak, chinkapin oak, common serviceberry, eastern redcedar, nannyberry, prairie crabapple, roughleaf dogwood, smooth sumac	Black oak, common hackberry, eastern white pine, green ash	Carolina poplar	---
567C2: Elkhart-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine

Table 10.--Windbreaks and Environmental Plantings--Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
567D2: Elkhart-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
630C2: Navlys-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
630D2: Navlys-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
630D3: Navlys-----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
679B: Blackberry--	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine

Table 10.--Windbreaks and Environmental Plantings--Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
684B: Broadwell---	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
684C2: Broadwell---	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
685B: Middletown--	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
685C2: Middletown--	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
705A: Buckhart----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine

Table 10.--Windbreaks and Environmental Plantings--Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
705B: Buckhart----	American hazelnut, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, coralberry, mapleleaf viburnum, redosier dogwood, silky dogwood	American plum, American witchhazel, blackhaw, common chokecherry, common serviceberry, prairie crabapple, roughleaf dogwood, smooth sumac, southern arrowwood	Washington hawthorn, arborvitae, blue spruce, common persimmon, eastern redcedar, nannyberry, pecan, white oak	Douglas fir, Norway spruce, black walnut, blackgum, common hackberry, green ash, northern red oak, pin oak, tuliptree	Carolina poplar, eastern cottonwood, eastern white pine
712A: Spaulding---	Common winterberry, gray dogwood, redosier dogwood	Common pawpaw, nannyberry, roughleaf dogwood, silky dogwood	Arborvitae, bur oak, common hackberry, eastern redcedar, green hawthorn	Carolina poplar, eastern cottonwood, green ash	---
801C: Orthents.					
830: Landfills.					
862: Pits, sand.					
864: Pits, quarries.					
3073A: Ross-----	American cranberrybush, Canada yew, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, northern spicebush, redosier dogwood, silky dogwood	Blackhaw, cockspur hawthorn, common pawpaw, common serviceberry, prairie crabapple, roughleaf dogwood, rusty blackhaw, southern arrowwood, witchhazel	Austrian pine, Douglas fir, arborvitae, blue spruce, common persimmon, eastern redcedar, green hawthorn, nannyberry, pecan, shingle oak	Norway spruce, blackgum, common hackberry, green ash, red maple, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
3074A: Radford----	American cranberrybush, Canada yew, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, northern spicebush, redosier dogwood, silky dogwood	Blackhaw, cockspur hawthorn, common pawpaw, common serviceberry, prairie crabapple, roughleaf dogwood, rusty blackhaw, southern arrowwood, witchhazel	Austrian pine, Douglas fir, arborvitae, blue spruce, common persimmon, eastern redcedar, green hawthorn, nannyberry, pecan, shingle oak	Norway spruce, blackgum, common hackberry, green ash, red maple, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak

Table 10.--Windbreaks and Environmental Plantings--Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
3077A: Huntsville--	American cranberrybush, Canada yew, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, northern spicebush, redosier dogwood, silky dogwood	Blackhaw, cockspur hawthorn, common pawpaw, common serviceberry, prairie crabapple, roughleaf dogwood, rusty blackhaw, southern arrowwood, witchhazel	Austrian pine, Douglas fir, arborvitae, blue spruce, common persimmon, eastern redcedar, green hawthorn, nannyberry, pecan, shingle oak	Norway spruce, blackgum, common hackberry, green ash, red maple, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
3107A: Sawmill-----	American cranberrybush, black chokeberry, buttonbush, common elderberry, common ninebark, common winterberry, gray dogwood, highbush blueberry, northern spicebush, redosier dogwood, silky dogwood	Cockspur hawthorn, hazel alder, nannyberry, roughleaf dogwood	Arborvitae, blackgum, common hackberry, green hawthorn, northern white-cedar, shingle oak	Green ash, red maple, river birch, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
3284A: Tice-----	American cranberrybush, Canada yew, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, northern spicebush, redosier dogwood, silky dogwood	Blackhaw, cockspur hawthorn, common pawpaw, common serviceberry, prairie crabapple, roughleaf dogwood, rusty blackhaw, southern arrowwood, witchhazel	Austrian pine, Douglas fir, arborvitae, blue spruce, common persimmon, eastern redcedar, green hawthorn, nannyberry, pecan, shingle oak	Norway spruce, blackgum, common hackberry, green ash, red maple, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
3405A: Zook-----	American cranberrybush, black chokeberry, buttonbush, common elderberry, common ninebark, common winterberry, gray dogwood, highbush blueberry, northern spicebush, redosier dogwood, silky dogwood	Cockspur hawthorn, hazel alder, nannyberry, roughleaf dogwood	Arborvitae, blackgum, common hackberry, green hawthorn, northern white-cedar, shingle oak	Green ash, red maple, river birch, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak

Table 10.--Windbreaks and Environmental Plantings--Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
3451A: Lawson-----	American cranberrybush, Canada yew, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, northern spicebush, redosier dogwood, silky dogwood	Blackhaw, cockspur hawthorn, common pawpaw, common serviceberry, prairie crabapple, roughleaf dogwood, rusty blackhaw, southern arrowwood, witchhazel	Austrian pine, Douglas fir, arborvitae, blue spruce, common persimmon, eastern redcedar, green hawthorn, nannyberry, pecan, shingle oak	Norway spruce, blackgum, common hackberry, green ash, red maple, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
7037A: Worthen-----	American cranberrybush, Canada yew, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, northern spicebush, redosier dogwood, silky dogwood	Blackhaw, cockspur hawthorn, common pawpaw, common serviceberry, prairie crabapple, roughleaf dogwood, rusty blackhaw, southern arrowwood, witchhazel	Austrian pine, Douglas fir, arborvitae, blue spruce, common persimmon, eastern redcedar, green hawthorn, nannyberry, pecan, shingle oak	Norway spruce, blackgum, common hackberry, green ash, red maple, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
7075B: Drury-----	American cranberrybush, Canada yew, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, northern spicebush, redosier dogwood, silky dogwood	Blackhaw, cockspur hawthorn, common pawpaw, common serviceberry, prairie crabapple, roughleaf dogwood, rusty blackhaw, southern arrowwood, witchhazel	Austrian pine, Douglas fir, arborvitae, blue spruce, common persimmon, eastern redcedar, green hawthorn, nannyberry, pecan, shingle oak	Norway spruce, blackgum, common hackberry, green ash, red maple, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
7148A: Proctor-----	American cranberrybush, Canada yew, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, northern spicebush, redosier dogwood, silky dogwood	Blackhaw, cockspur hawthorn, common pawpaw, common serviceberry, prairie crabapple, roughleaf dogwood, rusty blackhaw, southern arrowwood, witchhazel	Austrian pine, Douglas fir, arborvitae, blue spruce, common persimmon, eastern redcedar, green hawthorn, nannyberry, pecan, shingle oak	Norway spruce, blackgum, common hackberry, green ash, red maple, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak
7242A: Kendall-----	American cranberrybush, Canada yew, black chokeberry, common elderberry, common juniper, common ninebark, common winterberry, northern spicebush, redosier dogwood, silky dogwood	Blackhaw, cockspur hawthorn, common pawpaw, common serviceberry, prairie crabapple, roughleaf dogwood, rusty blackhaw, southern arrowwood, witchhazel	Austrian pine, Douglas fir, arborvitae, blue spruce, common persimmon, eastern redcedar, green hawthorn, nannyberry, pecan, shingle oak	Norway spruce, blackgum, common hackberry, green ash, red maple, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak

Table 10.--Windbreaks and Environmental Plantings--Continued

Map symbol and soil name	Trees having predicted 20-year average height, in feet, of--				
	<8	8-15	16-25	26-35	>35
8396A: Vesser-----	American cranberrybush, black chokeberry, buttonbush, common elderberry, common ninebark, common winterberry, gray dogwood, highbush blueberry, northern spicebush, redosier dogwood, silky dogwood	Cockspur hawthorn, hazel alder, nannyberry, roughleaf dogwood	Arborvitae, blackgum, common hackberry, green hawthorn, northern white-cedar, shingle oak	Green ash, red maple, river birch, swamp white oak, sweetgum	Carolina poplar, eastern cottonwood, pin oak

Table 11.--Forestland Productivity

Map symbol and soil name	Potential productivity			Suggested trees to plant
	Common trees	Site index	Volume of wood fiber	
8D:				
Hickory-----	Bitternut hickory-----	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
	Black oak-----	---	---	
	Green ash-----	---	---	
	Northern red oak-----	85	72	
	White oak-----	85	72	
8D2:				
Hickory-----	Northern red oak-----	85	72	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
	White oak-----	85	72	
	Black oak-----	---	---	
	Green ash-----	---	---	
	Bitternut hickory-----	---	---	
8D3:				
Hickory-----	Bitternut hickory-----	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
	Black oak-----	---	---	
	Green ash-----	---	---	
	Northern red oak-----	85	72	
	White oak-----	85	72	
8F:				
Hickory-----	Bitternut hickory-----	---	---	Eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
	Black oak-----	---	---	
	Green ash-----	---	---	
	Northern red oak-----	85	72	
	White oak-----	85	72	
17A:				
Keomah-----	Northern red oak-----	70	57	Common hackberry, eastern cottonwood, green ash, pin oak, river birch, swamp white oak, sweetgum.
	White oak-----	65	43	
36B:				
Tama-----	---	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
36C2:				
Tama-----	---	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
43A:				
Ipava-----	---	---	---	Common hackberry, common persimmon, eastern cottonwood, green ash, pecan, pin oak, swamp white oak.

Table 11.--Forestland Productivity--Continued

Map symbol and soil name	Potential productivity			Suggested trees to plant
	Common trees	Site index	Volume of wood fiber	
45A: Denny-----	---	---	---	Common hackberry, eastern cottonwood, green ash, pin oak, river birch, swamp white oak, sweetgum.
46A: Herrick-----	---	---	---	Common hackberry, common persimmon, eastern cottonwood, green ash, pecan, pin oak, swamp white oak.
50A: Virden-----	---	---	---	Common hackberry, eastern cottonwood, green ash, pin oak, river birch, swamp white oak, sweetgum.
68A: Sable-----	---	---	---	Common hackberry, eastern cottonwood, green ash, pin oak, river birch, swamp white oak, sweetgum.
86B: Osco-----	---	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
86C2: Osco-----	---	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
112A: Cowden-----	---	---	---	Common hackberry, eastern cottonwood, green ash, pin oak, river birch, swamp white oak, sweetgum.
119C2: Elco-----	Black walnut----- Northern red oak----- White oak-----	--- --- 80	--- --- 57	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
119D: Elco-----	Black walnut----- Northern red oak----- White oak-----	--- --- 80	--- --- 57	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.

Table 11.--Forestland Productivity--Continued

Map symbol and soil name	Potential productivity			Suggested trees to plant
	Common trees	Site index	Volume of wood fiber	
119D2: Elco-----	Northern red oak-----	85	72	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
	White oak-----	85	72	
	Black walnut-----	---	---	
119D3: Elco-----	Black walnut-----	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
	Northern red oak-----	---	---	
	White oak-----	80	57	
127B: Harrison-----	---	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
127C2: Harrison-----	---	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
131C2: Alvin-----	Black walnut-----	---	---	Black walnut, bur oak, eastern white pine, pecan, pin oak, tuliptree.
	Northern red oak-----	80	57	
	White oak-----	80	57	
131D2: Alvin-----	Black walnut-----	---	---	Black walnut, bur oak, eastern white pine, pecan, pin oak, tuliptree.
	Northern red oak-----	80	57	
	White oak-----	80	57	
134A: Camden-----	White oak-----	85	72	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
	Green ash-----	76	72	
	Northern red oak-----	85	72	
	Black walnut-----	---	---	
	Basswood-----	---	---	
134B: Camden-----	White oak-----	85	72	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
	Green ash-----	76	72	
	Northern red oak-----	85	72	
	Basswood-----	---	---	
	Black walnut-----	---	---	
134C2: Camden-----	Northern red oak-----	85	72	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
	White oak-----	85	72	
	Green ash-----	76	72	
	Black walnut-----	---	---	
	Basswood-----	---	---	

Table 11.--Forestland Productivity--Continued

Map symbol and soil name	Potential productivity			Suggested trees to plant
	Common trees	Site index	Volume of wood fiber	
138A: Shiloh-----	---	---	---	Common hackberry, eastern cottonwood, green ash, pin oak, river birch, swamp white oak, sweetgum.
198A: Elburn-----	---	---	---	Common hackberry, common persimmon, eastern cottonwood, green ash, pecan, pin oak, swamp white oak.
199A: Plano-----	---	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
199B: Plano-----	---	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
212C2: Thebes-----	Black walnut----- Northern red oak----- White oak-----	--- --- 80	--- --- 57	Black oak, common hackberry, eastern white pine, green ash.
242A: Kendall-----	White oak----- Northern red oak-----	80 80	57 57	Common hackberry, common persimmon, eastern cottonwood, green ash, pecan, pin oak, swamp white oak.
244A: Hartsburg-----	---	---	---	Common hackberry, eastern cottonwood, green ash, pin oak, river birch, swamp white oak, sweetgum.
249A: Edinburg-----	---	---	---	Common hackberry, eastern cottonwood, green ash, pin oak, river birch, swamp white oak, sweetgum.
257A: Clarksdale-----	White oak----- Northern red oak-----	80 80	57 57	Common hackberry, common persimmon, eastern cottonwood, green ash, pecan, pin oak, swamp white oak.

Table 11.--Forestland Productivity--Continued

Map symbol and soil name	Potential productivity			Suggested trees to plant
	Common trees	Site index	Volume of wood fiber	
259C2: Assumption-----	---	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
259D2: Assumption-----	---	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
279B: Rozetta-----	White oak----- Northern red oak----- Black walnut-----	80 80 ---	57 57 ---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
280C2: Fayette-----	Northern red oak----- White oak----- Black walnut-----	80 80 ---	57 57 ---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
280D2: Fayette-----	Black walnut----- Northern red oak----- White oak-----	--- 80 80	--- 57 57	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
280D3: Fayette-----	Black walnut----- Northern red oak----- White oak-----	--- 80 80	--- 57 57	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
533: Urban land.				
536: Dumps.				
549G: Marseilles-----	White oak-----	66	29	Black oak, common hackberry, eastern white pine, green ash.
567C2: Elkhart-----	---	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.

Table 11.--Forestland Productivity--Continued

Map symbol and soil name	Potential productivity			Suggested trees to plant
	Common trees	Site index	Volume of wood fiber	
567D2: Elkhart-----	---	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
630C2: Navlys-----	White oak----- Northern red oak----- Black walnut----- Yellow poplar-----	85 85 --- ---	72 72 --- ---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
630D2: Navlys-----	White oak----- Northern red oak----- Black walnut----- Yellow poplar-----	85 85 --- ---	72 72 --- ---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
630D3: Navlys-----	White oak----- Northern red oak----- Black walnut----- Yellow poplar-----	85 85 --- ---	72 72 --- ---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
679B: Blackberry-----	---	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
684B: Broadwell-----	---	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
684C2: Broadwell-----	---	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
685B: Middletown-----	Black walnut----- Northern red oak----- White oak-----	--- 80 80	--- 57 57	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
685C2: Middletown-----	Black walnut----- Northern red oak----- White oak-----	--- 80 80	--- 57 57	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.

Table 11.--Forestland Productivity--Continued

Map symbol and soil name	Potential productivity			Suggested trees to plant
	Common trees	Site index	Volume of wood fiber	
705A: Buckhart-----	---	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
705B: Buckhart-----	---	---	---	Black walnut, eastern cottonwood, eastern white pine, green ash, northern red oak, pecan, pin oak, tuliptree, white oak.
712A: Spaulding-----	---	---	---	Bur oak, common hackberry, eastern cottonwood, eastern redcedar, green ash.
801C: Orthents.				
830: Landfills.				
862: Pits, sand				
864: Pits, quarries.				
3073A: Ross-----	Black cherry----- Black walnut----- Northern red oak----- White ash----- White oak-----	--- --- 86 --- ---	--- --- 72 --- ---	Common hackberry, common persimmon, eastern cottonwood, green ash, pecan, pin oak, swamp white oak.
3074A: Radford-----	Pin oak----- Eastern cottonwood----- Sweetgum----- Tuliptree----- White ash-----	96 --- 86 90 ---	72 --- 100 86 ---	Common hackberry, common persimmon, eastern cottonwood, green ash, pecan, pin oak, swamp white oak.
3077A: Huntsville-----	American sycamore----- Eastern cottonwood----- Green ash-----	--- 110 ---	--- 157 ---	Common hackberry, common persimmon, eastern cottonwood, green ash, pecan, pin oak, swamp white oak.
3107A: Sawmill-----	Pin oak----- American sycamore----- Eastern cottonwood----- Sweetgum-----	90 --- --- ---	72 --- --- ---	Common hackberry, eastern cottonwood, green ash, pin oak, river birch, swamp white oak, sweetgum.

Table 11.--Forestland Productivity--Continued

Map symbol and soil name	Potential productivity			Suggested trees to plant
	Common trees	Site index	Volume of wood fiber	
3284A: Tice-----	Virginia pine-----	90	90	Common hackberry, common persimmon, eastern cottonwood, green ash, pecan, pin oak, swamp white oak.
	Eastern cottonwood-----	---	---	
	Pin oak-----	96	78	
	Sweetgum-----	86	95	
	Tuliptree-----	---	---	
	White ash-----	---	---	
3405A: Zook-----	---	---	---	Common hackberry, eastern cottonwood, green ash, pin oak, river birch, swamp white oak, sweetgum.
3451A: Lawson-----	Silver maple-----	70	29	Common hackberry, common persimmon, eastern cottonwood, green ash, pecan, pin oak, swamp white oak.
	White ash-----	---	---	
	American elm-----	---	---	
7037A: Worthen-----	---	---	---	Common hackberry, common persimmon, eastern cottonwood, green ash, pecan, pin oak, swamp white oak.
7075B: Drury-----	White oak-----	85	72	Common hackberry, common persimmon, eastern cottonwood, green ash, pecan, pin oak, swamp white oak.
	Northern red oak-----	85	72	
	Green ash-----	---	---	
7148A: Proctor-----	---	---	---	Common hackberry, common persimmon, eastern cottonwood, green ash, pecan, pin oak, swamp white oak.
7242A: Kendall-----	Green ash-----	---	---	Common hackberry, common persimmon, eastern cottonwood, green ash, pecan, pin oak, swamp white oak.
	Bur oak-----	---	---	
8396A: Vesser-----	---	---	---	Common hackberry, eastern cottonwood, green ash, pin oak, river birch, swamp white oak, sweetgum.

Table 12.--Forestland Management (Part 1)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
8D: Hickory-----	Moderate Low strength	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Low strength	1.00
8D2: Hickory-----	Moderate Low strength	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Low strength	1.00
8D3: Hickory-----	Moderate Low strength	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Low strength	1.00
8F: Hickory-----	Moderate Slope Low strength	0.50 0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Low strength	1.00
17A: Keomah-----	Moderate Low strength	0.50	Moderately suited Wetness Low strength	0.50 0.50	Severe Low strength	1.00
36B: Tama-----	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Low strength	1.00
36C2: Tama-----	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Low strength	1.00
43A: Ipava-----	Moderate Low strength	0.50	Moderately suited Low strength Wetness	0.50 0.50	Severe Low strength	1.00
45A: Denny-----	Moderate Low strength	0.50	Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50	Severe Low strength	1.00
46A: Herrick-----	Moderate Low strength	0.50	Moderately suited Low strength Wetness	0.50 0.50	Severe Low strength	1.00

Table 12.--Forestland Management (Part 1)--Continued

Map symbol and soil name	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
50A: Virden-----	Moderate Low strength	0.50	Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50	Severe Low strength	1.00
68A: Sable-----	Moderate Low strength	0.50	Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50	Severe Low strength	1.00
86B: Osco-----	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Low strength	1.00
86C2: Osco-----	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Low strength	1.00
112A: Cowden-----	Moderate Low strength	0.50	Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50	Severe Low strength	1.00
119C2: Elco-----	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Low strength	1.00
119D: Elco-----	Moderate Low strength	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Low strength	1.00
119D2: Elco-----	Moderate Low strength	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Low strength	1.00
119D3: Elco-----	Moderate Low strength	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Low strength	1.00
127B: Harrison-----	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Low strength	1.00
127C2: Harrison-----	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Low strength	1.00

Table 12.--Forestland Management (Part 1)--Continued

Map symbol and soil name	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
131C2: Alvin-----	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Low strength	1.00
131D2: Alvin-----	Moderate Low strength	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Low strength	1.00
134A: Camden-----	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Low strength	1.00
134B: Camden-----	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Low strength	1.00
134C2: Camden-----	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Low strength	1.00
138A: Shiloh-----	Moderate Low strength	0.50	Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50	Severe Low strength	1.00
198A: Elburn-----	Moderate Low strength	0.50	Moderately suited Low strength Wetness	0.50 0.50	Severe Low strength	1.00
199A: Plano-----	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Low strength	1.00
199B: Plano-----	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Low strength	1.00
212C2: Thebes-----	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Low strength	1.00
242A: Kendall-----	Moderate Low strength	0.50	Moderately suited Wetness Low strength	0.50 0.50	Severe Low strength	1.00
244A: Hartsburg-----	Moderate Low strength	0.50	Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50	Severe Low strength	1.00

Table 12.--Forestland Management (Part 1)--Continued

Map symbol and soil name	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
249A: Edinburg-----	Moderate Low strength	0.50	Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50	Severe Low strength	1.00
257A: Clarksdale-----	Moderate Low strength	0.50	Moderately suited Wetness Low strength	0.50 0.50	Severe Low strength	1.00
259C2: Assumption-----	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Low strength	1.00
259D2: Assumption-----	Moderate Low strength	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Low strength	1.00
279B: Rozetta-----	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Low strength	1.00
280C2: Fayette-----	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Low strength	1.00
280D2: Fayette-----	Moderate Low strength	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Low strength	1.00
280D3: Fayette-----	Moderate Low strength	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Low strength	1.00
533: Urban land-----	Not rated		Not rated		Not rated	
536: Dumps-----	Not rated		Not rated		Not rated	
549G: Marseilles-----	Severe Slope Low strength	1.00 0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Low strength	1.00
567C2: Elkhart-----	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Low strength	1.00

Table 12.--Forestland Management (Part 1)--Continued

Map symbol and soil name	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
567D2: Elkhart-----	Moderate Low strength	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Low strength	1.00
630C2: Navlys-----	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Low strength	1.00
630D2: Navlys-----	Moderate Low strength	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Low strength	1.00
630D3: Navlys-----	Moderate Low strength	0.50	Poorly suited Slope Low strength	1.00 0.50	Severe Low strength	1.00
679B: Blackberry-----	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Low strength	1.00
684B: Broadwell-----	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Low strength	1.00
684C2: Broadwell-----	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Low strength	1.00
685B: Middletown-----	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Low strength	1.00
685C2: Middletown-----	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Low strength	1.00
705A: Buckhart-----	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Low strength	1.00
705B: Buckhart-----	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Low strength	1.00
712A: Spaulding-----	Moderate Low strength	0.50	Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50	Severe Low strength	1.00

Table 12.--Forestland Management (Part 1)--Continued

Map symbol and soil name	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
801C: Orthents-----	Moderate Low strength	0.50	Moderately suited Low strength Slope	0.50 0.50	Severe Low strength	1.00
830: Landfills-----	Not rated		Not rated		Not rated	
862: Pits, sand-----	Not rated		Not rated		Not rated	
864: Pits, quarries----	Not rated		Not rated		Not rated	
3073A: Ross-----	Severe Flooding Low strength	1.00 0.50	Poorly suited Flooding Low strength	1.00 0.50	Severe Low strength	1.00
3074A: Radford-----	Severe Flooding Low strength	1.00 0.50	Poorly suited Flooding Low strength Wetness	1.00 0.50 0.50	Severe Low strength	1.00
3077A: Huntsville-----	Severe Flooding Low strength	1.00 0.50	Poorly suited Flooding Low strength	1.00 0.50	Severe Low strength	1.00
3107A: Sawmill-----	Severe Flooding Low strength	1.00 0.50	Poorly suited Ponding Flooding Wetness Low strength	1.00 1.00 1.00 0.50	Severe Low strength	1.00
3284A: Tice-----	Severe Flooding Low strength	1.00 0.50	Poorly suited Flooding Low strength Wetness	1.00 0.50 0.50	Severe Low strength	1.00
3405A: Zook-----	Severe Flooding Low strength	1.00 0.50	Poorly suited Flooding Wetness Low strength	1.00 1.00 0.50	Severe Low strength	1.00
3451A: Lawson-----	Severe Flooding Low strength	1.00 0.50	Poorly suited Flooding Low strength Wetness	1.00 0.50 0.50	Severe Low strength	1.00

Table 12.--Forestland Management (Part 1)--Continued

Map symbol and soil name	Limitations affecting construction of haul roads and log landings		Suitability for log landings		Soil rutting hazard	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
7037A: Worthen-----	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Low strength	1.00
7075B: Drury-----	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Low strength	1.00
7148A: Proctor-----	Moderate Low strength	0.50	Moderately suited Low strength	0.50	Severe Low strength	1.00
7242A: Kendall-----	Moderate Low strength	0.50	Moderately suited Wetness Low strength	0.50 0.50	Severe Low strength	1.00
8396A: Vesser-----	Severe Flooding Low strength	1.00 0.50	Poorly suited Flooding Wetness Low strength	1.00 1.00 0.50	Severe Low strength	1.00

Table 13.--Forestland Management (Part 2)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
8D: Hickory-----	Moderate Slope/erodibility	0.27	Severe Slope/erodibility	1.00	Poorly suited Slope Low strength	1.00 0.50
8D2: Hickory-----	Moderate Slope/erodibility	0.27	Severe Slope/erodibility	1.00	Poorly suited Slope Low strength	1.00 0.50
8D3: Hickory-----	Moderate Slope/erodibility	0.27	Severe Slope/erodibility	1.00	Poorly suited Slope Low strength	1.00 0.50
8F: Hickory-----	Moderate Slope/erodibility	0.52	Severe Slope/erodibility	1.00	Poorly suited Slope Low strength	1.00 0.50
17A: Keomah-----	Slight Slope/erodibility	0.02	Slight Slope/erodibility	0.11	Moderately suited Wetness Low strength	0.50 0.50
36B: Tama-----	Slight Slope/erodibility	0.07	Moderate Slope/erodibility	0.39	Moderately suited Low strength	0.50
36C2: Tama-----	Slight Slope/erodibility	0.18	Moderate Slope/erodibility	0.83	Moderately suited Low strength Slope	0.50 0.50
43A: Ipava-----	Slight Slope/erodibility	0.02	Slight Slope/erodibility	0.11	Moderately suited Low strength Wetness	0.50 0.50
45A: Denny-----	Slight Slope/erodibility	0.01	Slight Slope/erodibility	0.06	Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50
46A: Herrick-----	Slight Slope/erodibility	0.01	Slight Slope/erodibility	0.06	Moderately suited Low strength Wetness	0.50 0.50

Table 13.--Forestland Management (Part 2)--Continued

Map symbol and soil name	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
50A: Virden-----	Slight Slope/erodibility	0.01	Slight Slope/erodibility	0.06	Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50
68A: Sable-----	Slight Slope/erodibility	0.01	Slight Slope/erodibility	0.06	Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50
86B: Osc-----	Slight Slope/erodibility	0.07	Moderate Slope/erodibility	0.39	Moderately suited Low strength	0.50
86C2: Osc-----	Slight Slope/erodibility	0.18	Moderate Slope/erodibility	0.83	Moderately suited Low strength Slope	0.50 0.50
112A: Cowden-----	Slight Slope/erodibility	0.01	Slight Slope/erodibility	0.06	Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50
119C2: Elco-----	Slight Slope/erodibility	0.18	Moderate Slope/erodibility	0.83	Moderately suited Low strength Slope	0.50 0.50
119D: Elco-----	Moderate Slope/erodibility	0.34	Severe Slope/erodibility	1.00	Poorly suited Slope Low strength	1.00 0.50
119D2: Elco-----	Moderate Slope/erodibility	0.34	Severe Slope/erodibility	1.00	Poorly suited Slope Low strength	1.00 0.50
119D3: Elco-----	Moderate Slope/erodibility	0.34	Severe Slope/erodibility	1.00	Poorly suited Slope Low strength	1.00 0.50
127B: Harrison-----	Slight Slope/erodibility	0.07	Moderate Slope/erodibility	0.39	Moderately suited Low strength	0.50
127C2: Harrison-----	Slight Slope/erodibility	0.18	Moderate Slope/erodibility	0.83	Moderately suited Low strength Slope	0.50 0.50

Table 13.--Forestland Management (Part 2)--Continued

Map symbol and soil name	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
131C2: Alvin-----	Slight Slope/erodibility	0.15	Moderate Slope/erodibility	0.47	Moderately suited Low strength Slope	0.50 0.50
131D2: Alvin-----	Moderate Slope/erodibility	0.27	Moderate Slope/erodibility	0.88	Poorly suited Slope Low strength	1.00 0.50
134A: Camden-----	Slight Slope/erodibility	0.02	Slight Slope/erodibility	0.11	Moderately suited Low strength	0.50
134B: Camden-----	Slight Slope/erodibility	0.09	Moderate Slope/erodibility	0.39	Moderately suited Low strength	0.50
134C2: Camden-----	Slight Slope/erodibility	0.18	Moderate Slope/erodibility	0.83	Moderately suited Low strength Slope	0.50 0.50
138A: Shiloh-----	Slight Slope/erodibility	0.01	Slight Slope/erodibility	0.06	Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50
198A: Elburn-----	Slight Slope/erodibility	0.02	Slight Slope/erodibility	0.11	Moderately suited Low strength Wetness	0.50 0.50
199A: Plano-----	Slight Slope/erodibility	0.02	Slight Slope/erodibility	0.11	Moderately suited Low strength	0.50
199B: Plano-----	Slight Slope/erodibility	0.07	Moderate Slope/erodibility	0.39	Moderately suited Low strength	0.50
212C2: Thebes-----	Slight Slope/erodibility	0.18	Moderate Slope/erodibility	0.83	Moderately suited Low strength Slope	0.50 0.50
242A: Kendall-----	Slight Slope/erodibility	0.02	Slight Slope/erodibility	0.11	Moderately suited Wetness Low strength	0.50 0.50
244A: Hartsburg-----	Slight Slope/erodibility	0.01	Slight Slope/erodibility	0.06	Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50

Table 13.--Forestland Management (Part 2)--Continued

Map symbol and soil name	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
249A: Edinburg-----	Slight Slope/erodibility	0.01	Slight Slope/erodibility	0.06	Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50
257A: Clarksdale-----	Slight Slope/erodibility	0.02	Slight Slope/erodibility	0.11	Moderately suited Wetness Low strength	0.50 0.50
259C2: Assumption-----	Slight Slope/erodibility	0.18	Moderate Slope/erodibility	0.83	Moderately suited Low strength Slope	0.50 0.50
259D2: Assumption-----	Moderate Slope/erodibility	0.34	Severe Slope/erodibility	1.00	Poorly suited Slope Low strength	1.00 0.50
279B: Rozetta-----	Slight Slope/erodibility	0.09	Moderate Slope/erodibility	0.39	Moderately suited Low strength	0.50
280C2: Fayette-----	Slight Slope/erodibility	0.18	Moderate Slope/erodibility	0.83	Moderately suited Low strength Slope	0.50 0.50
280D2: Fayette-----	Moderate Slope/erodibility	0.34	Severe Slope/erodibility	1.00	Poorly suited Slope Low strength	1.00 0.50
280D3: Fayette-----	Moderate Slope/erodibility	0.34	Severe Slope/erodibility	1.00	Poorly suited Slope Low strength	1.00 0.50
533: Urban land-----	Not rated		Not rated		Not rated	
536: Dumps-----	Not rated		Not rated		Not rated	
549G: Marseilles-----	Severe Slope/erodibility	0.93	Severe Slope/erodibility	1.00	Poorly suited Slope Low strength	1.00 0.50
567C2: Elkhart-----	Slight Slope/erodibility	0.18	Moderate Slope/erodibility	0.83	Moderately suited Low strength Slope	0.50 0.50

Table 13.--Forestland Management (Part 2)--Continued

Map symbol and soil name	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
567D2: Elkhart-----	Moderate Slope/erodibility	0.34	Severe Slope/erodibility	1.00	Poorly suited Slope Low strength	1.00 0.50
630C2: Navlys-----	Slight Slope/erodibility	0.18	Moderate Slope/erodibility	0.83	Moderately suited Low strength Slope	0.50 0.50
630D2: Navlys-----	Moderate Slope/erodibility	0.34	Severe Slope/erodibility	1.00	Poorly suited Slope Low strength	1.00 0.50
630D3: Navlys-----	Moderate Slope/erodibility	0.34	Severe Slope/erodibility	1.00	Poorly suited Slope Low strength	1.00 0.50
679B: Blackberry-----	Slight Slope/erodibility	0.07	Moderate Slope/erodibility	0.39	Moderately suited Low strength	0.50
684B: Broadwell-----	Slight Slope/erodibility	0.07	Moderate Slope/erodibility	0.39	Moderately suited Low strength	0.50
684C2: Broadwell-----	Slight Slope/erodibility	0.18	Moderate Slope/erodibility	0.83	Moderately suited Low strength Slope	0.50 0.50
685B: Middletown-----	Slight Slope/erodibility	0.09	Moderate Slope/erodibility	0.39	Moderately suited Low strength	0.50
685C2: Middletown-----	Slight Slope/erodibility	0.18	Moderate Slope/erodibility	0.83	Moderately suited Low strength Slope	0.50 0.50
705A: Buckhart-----	Slight Slope/erodibility	0.02	Slight Slope/erodibility	0.11	Moderately suited Low strength	0.50
705B: Buckhart-----	Slight Slope/erodibility	0.07	Moderate Slope/erodibility	0.39	Moderately suited Low strength	0.50
712A: Spaulding-----	Slight Slope/erodibility	0.01	Slight Slope/erodibility	0.06	Poorly suited Ponding Wetness Low strength	1.00 1.00 0.50

Table 13.--Forestland Management (Part 2)--Continued

Map symbol and soil name	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
801C: Orthents-----	Slight Slope/erodibility	0.18	Moderate Slope/erodibility	0.83	Moderately suited Low strength Slope	0.50 0.50
830: Landfills-----	Not rated		Not rated		Not rated	
862: Pits, sand-----	Not rated		Not rated		Not rated	
864: Pits, quarries-----	Not rated		Not rated		Not rated	
3073A: Ross-----	Slight Slope/erodibility	0.02	Slight Slope/erodibility	0.11	Poorly suited Flooding Low strength	1.00 0.50
3074A: Radford-----	Slight Slope/erodibility	0.02	Slight Slope/erodibility	0.11	Poorly suited Flooding Low strength Wetness	1.00 0.50 0.50
3077A: Huntsville-----	Slight Slope/erodibility	0.02	Slight Slope/erodibility	0.11	Poorly suited Flooding Low strength	1.00 0.50
3107A: Sawmill-----	Slight Slope/erodibility	0.01	Slight Slope/erodibility	0.06	Poorly suited Ponding Flooding Wetness Low strength	1.00 1.00 1.00 0.50
3284A: Tice-----	Slight Slope/erodibility	0.02	Slight Slope/erodibility	0.11	Poorly suited Flooding Low strength Wetness	1.00 0.50 0.50
3405A: Zook-----	Slight Slope/erodibility	0.01	Slight Slope/erodibility	0.06	Poorly suited Flooding Wetness Low strength	1.00 1.00 0.50
3451A: Lawson-----	Slight Slope/erodibility	0.02	Slight Slope/erodibility	0.11	Poorly suited Flooding Low strength Wetness	1.00 0.50 0.50

Table 13.--Forestland Management (Part 2)--Continued

Map symbol and soil name	Hazard of off-road or off-trail erosion		Hazard of erosion on roads and trails		Suitability for roads (natural surface)	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
7037A: Worthen-----	Slight Slope/erodibility	0.02	Slight Slope/erodibility	0.11	Moderately suited Low strength	0.50
7075B: Drury-----	Slight Slope/erodibility	0.09	Moderate Slope/erodibility	0.39	Moderately suited Low strength	0.50
7148A: Proctor-----	Slight Slope/erodibility	0.02	Slight Slope/erodibility	0.11	Moderately suited Low strength	0.50
7242A: Kendall-----	Slight Slope/erodibility	0.02	Slight Slope/erodibility	0.11	Moderately suited Wetness Low strength	0.50 0.50
8396A: Vesser-----	Slight Slope/erodibility	0.01	Slight Slope/erodibility	0.06	Poorly suited Flooding Wetness Low strength	1.00 1.00 0.50

Table 14.--Forestland Management (Part 3)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Suitability for hand planting		Suitability for mechanical planting		Suitability for use of harvesting equipment	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
8D: Hickory-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
8D2: Hickory-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
8D3: Hickory-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
8F: Hickory-----	Moderately suited Stickiness	0.50	Unsuited Slope Stickiness	1.00 0.50	Moderately suited Low strength Slope	0.50 0.50
17A: Keomah-----	Well suited		Well suited		Moderately suited Low strength	0.50
36B: Tama-----	Well suited		Well suited		Moderately suited Low strength	0.50
36C2: Tama-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
43A: Ipava-----	Well suited		Well suited		Moderately suited Low strength	0.50
45A: Denny-----	Well suited		Well suited		Moderately suited Low strength	0.50
46A: Herrick-----	Well suited		Well suited		Moderately suited Low strength	0.50
50A: Virden-----	Moderately suited Stickiness	0.50	Moderately suited Stickiness	0.50	Moderately suited Low strength	0.50
68A: Sable-----	Moderately suited Stickiness	0.50	Moderately suited Stickiness	0.50	Moderately suited Low strength	0.50

Table 14.--Forestland Management (Part 3)--Continued

Map symbol and soil name	Suitability for hand planting		Suitability for mechanical planting		Suitability for use of harvesting equipment	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
86B: Osco-----	Moderately suited Stickiness	0.50	Moderately suited Stickiness	0.50	Moderately suited Low strength	0.50
86C2: Osco-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
112A: Cowden-----	Well suited		Well suited		Moderately suited Low strength	0.50
119C2: Elco-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
119D: Elco-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
119D2: Elco-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
119D3: Elco-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
127B: Harrison-----	Moderately suited Stickiness	0.50	Moderately suited Stickiness	0.50	Moderately suited Low strength	0.50
127C2: Harrison-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
131C2: Alvin-----	Well suited		Moderately suited Slope	0.50	Moderately suited Low strength	0.50
131D2: Alvin-----	Well suited		Moderately suited Slope	0.50	Moderately suited Low strength	0.50
134A: Camden-----	Well suited		Well suited		Moderately suited Low strength	0.50
134B: Camden-----	Well suited		Well suited		Moderately suited Low strength	0.50

Table 14.--Forestland Management (Part 3)--Continued

Map symbol and soil name	Suitability for hand planting		Suitability for mechanical planting		Suitability for use of harvesting equipment	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
134C2: Camden-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
138A: Shiloh-----	Moderately suited Stickiness	0.50	Moderately suited Stickiness	0.50	Moderately suited Low strength	0.50
198A: Elburn-----	Well suited		Well suited		Moderately suited Low strength	0.50
199A: Plano-----	Well suited		Well suited		Moderately suited Low strength	0.50
199B: Plano-----	Well suited		Well suited		Moderately suited Low strength	0.50
212C2: Thebes-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
242A: Kendall-----	Well suited		Well suited		Moderately suited Low strength	0.50
244A: Hartsburg-----	Moderately suited Stickiness	0.50	Moderately suited Stickiness	0.50	Moderately suited Low strength	0.50
249A: Edinburg-----	Moderately suited Stickiness	0.50	Moderately suited Stickiness	0.50	Moderately suited Low strength	0.50
257A: Clarksdale-----	Well suited		Well suited		Moderately suited Low strength	0.50
259C2: Assumption-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
259D2: Assumption-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
279B: Rozetta-----	Moderately suited Stickiness	0.50	Moderately suited Stickiness	0.50	Moderately suited Low strength	0.50

Table 14.--Forestland Management (Part 3)--Continued

Map symbol and soil name	Suitability for hand planting		Suitability for mechanical planting		Suitability for use of harvesting equipment	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
280C2: Fayette-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
280D2: Fayette-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
280D3: Fayette-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
533: Urban land-----	Not rated		Not rated		Not rated	
536: Dumps-----	Not rated		Not rated		Not rated	
549G: Marseilles-----	Moderately suited Slope Stickiness	0.50 0.50	Unsuited Slope Stickiness	1.00 0.50	Poorly suited Slope Low strength	1.00 0.50
567C2: Elkhart-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
567D2: Elkhart-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
630C2: Navlys-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
630D2: Navlys-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
630D3: Navlys-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
679B: Blackberry-----	Well suited		Well suited		Moderately suited Low strength	0.50

Table 14.--Forestland Management (Part 3)--Continued

Map symbol and soil name	Suitability for hand planting		Suitability for mechanical planting		Suitability for use of harvesting equipment	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
684B: Broadwell-----	Well suited		Well suited		Moderately suited Low strength	0.50
684C2: Broadwell-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
685B: Middletown-----	Moderately suited Stickiness	0.50	Moderately suited Stickiness	0.50	Moderately suited Low strength	0.50
685C2: Middletown-----	Moderately suited Stickiness	0.50	Moderately suited Slope Stickiness	0.50 0.50	Moderately suited Low strength	0.50
705A: Buckhart-----	Well suited		Well suited		Moderately suited Low strength	0.50
705B: Buckhart-----	Well suited		Well suited		Moderately suited Low strength	0.50
712A: Spaulding-----	Moderately suited Stickiness	0.50	Moderately suited Stickiness	0.50	Moderately suited Low strength	0.50
801C: Orthents-----	Well suited		Moderately suited Slope	0.50	Moderately suited Low strength	0.50
830: Landfills-----	Not rated		Not rated		Not rated	
862: Pits, sand-----	Not rated		Not rated		Not rated	
864: Pits, quarries-----	Not rated		Not rated		Not rated	
3073A: Ross-----	Well suited		Well suited		Moderately suited Low strength	0.50
3074A: Radford-----	Well suited		Well suited		Moderately suited Low strength	0.50
3077A: Huntsville-----	Well suited		Well suited		Moderately suited Low strength	0.50

Table 14.--Forestland Management (Part 3)--Continued

Map symbol and soil name	Suitability for hand planting		Suitability for mechanical planting		Suitability for use of harvesting equipment	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
3107A: Sawmill-----	Moderately suited Stickiness	0.50	Moderately suited Stickiness	0.50	Moderately suited Low strength	0.50
3284A: Tice-----	Moderately suited Stickiness	0.50	Moderately suited Stickiness	0.50	Moderately suited Low strength	0.50
3405A: Zook-----	Poorly suited Stickiness	0.75	Poorly suited Stickiness	0.75	Moderately suited Low strength	0.50
3451A: Lawson-----	Well suited		Well suited		Moderately suited Low strength	0.50
7037A: Worthen-----	Well suited		Well suited		Moderately suited Low strength	0.50
7075B: Drury-----	Well suited		Well suited		Moderately suited Low strength	0.50
7148A: Proctor-----	Well suited		Well suited		Moderately suited Low strength	0.50
7242A: Kendall-----	Well suited		Well suited		Moderately suited Low strength	0.50
8396A: Vesser-----	Well suited		Well suited		Moderately suited Low strength	0.50

Table 15.--Forestland Management (Part 4)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Suitability for mechanical site preparation (surface)		Suitability for mechanical site preparation (deep)	
	Rating class and limiting features	Value	Rating class and limiting features	Value
8D: Hickory-----	Well suited		Well suited	
8D2: Hickory-----	Well suited		Well suited	
8D3: Hickory-----	Well suited		Well suited	
8F: Hickory-----	Poorly suited Slope	0.50	Poorly suited Slope	0.50
17A: Keomah-----	Well suited		Well suited	
36B: Tama-----	Well suited		Well suited	
36C2: Tama-----	Well suited		Well suited	
43A: Ipava-----	Well suited		Well suited	
45A: Denny-----	Well suited		Well suited	
46A: Herrick-----	Well suited		Well suited	
50A: Virden-----	Well suited		Well suited	
68A: Sable-----	Well suited		Well suited	
86B: Osco-----	Well suited		Well suited	
86C2: Osco-----	Well suited		Well suited	
112A: Cowden-----	Well suited		Well suited	
119C2: Elco-----	Well suited		Well suited	
119D: Elco-----	Well suited		Well suited	

Table 15.--Forestland Management (Part 4)--Continued

Map symbol and soil name	Suitability for mechanical site preparation (surface)		Suitability for mechanical site preparation (deep)	
	Rating class and limiting features	Value	Rating class and limiting features	Value
119D2: Elco-----	Well suited		Well suited	
119D3: Elco-----	Well suited		Well suited	
127B: Harrison-----	Well suited		Well suited	
127C2: Harrison-----	Well suited		Well suited	
131C2: Alvin-----	Well suited		Well suited	
131D2: Alvin-----	Well suited		Well suited	
134A: Camden-----	Well suited		Well suited	
134B: Camden-----	Well suited		Well suited	
134C2: Camden-----	Well suited		Well suited	
138A: Shiloh-----	Well suited		Well suited	
198A: Elburn-----	Well suited		Well suited	
199A: Plano-----	Well suited		Well suited	
199B: Plano-----	Well suited		Well suited	
212C2: Thebes-----	Well suited		Well suited	
242A: Kendall-----	Well suited		Well suited	
244A: Hartsburg-----	Well suited		Well suited	
249A: Edinburg-----	Well suited		Well suited	
257A: Clarksdale-----	Well suited		Well suited	
259C2: Assumption-----	Well suited		Well suited	
259D2: Assumption-----	Well suited		Well suited	

Table 15.--Forestland Management (Part 4)--Continued

Map symbol and soil name	Suitability for mechanical site preparation (surface)		Suitability for mechanical site preparation (deep)	
	Rating class and limiting features	Value	Rating class and limiting features	Value
279B: Rozetta-----	Well suited		Well suited	
280C2: Fayette-----	Well suited		Well suited	
280D2: Fayette-----	Well suited		Well suited	
280D3: Fayette-----	Well suited		Well suited	
533: Urban land-----	Not rated		Not rated	
536: Dumps-----	Not rated		Not rated	
549G: Marseilles-----	Unsuited Slope	1.00	Unsuited Slope	1.00
567C2: Elkhart-----	Well suited		Well suited	
567D2: Elkhart-----	Well suited		Well suited	
630C2: Navlys-----	Well suited		Well suited	
630D2: Navlys-----	Well suited		Well suited	
630D3: Navlys-----	Well suited		Well suited	
679B: Blackberry-----	Well suited		Well suited	
684B: Broadwell-----	Well suited		Well suited	
684C2: Broadwell-----	Well suited		Well suited	
685B: Middletown-----	Well suited		Well suited	
685C2: Middletown-----	Well suited		Well suited	
705A: Buckhart-----	Well suited		Well suited	
705B: Buckhart-----	Well suited		Well suited	
712A: Spaulding-----	Well suited		Well suited	

Table 15.--Forestland Management (Part 4)--Continued

Map symbol and soil name	Suitability for mechanical site preparation (surface)		Suitability for mechanical site preparation (deep)	
	Rating class and limiting features	Value	Rating class and limiting features	Value
801C: Orthents-----	Well suited		Well suited	
830: Landfills-----	Not rated		Not rated	
862: Pits, sand-----	Not rated		Not rated	
864: Pits, quarries-----	Not rated		Not rated	
3073A: Ross-----	Well suited		Well suited	
3074A: Radford-----	Well suited		Well suited	
3077A: Huntsville-----	Well suited		Well suited	
3107A: Sawmill-----	Well suited		Well suited	
3284A: Tice-----	Well suited		Well suited	
3405A: Zook-----	Poorly suited Stickiness	0.50	Well suited	
3451A: Lawson-----	Well suited		Well suited	
7037A: Worthen-----	Well suited		Well suited	
7075B: Drury-----	Well suited		Well suited	
7148A: Proctor-----	Well suited		Well suited	
7242A: Kendall-----	Well suited		Well suited	
8396A: Vesser-----	Well suited		Well suited	

Table 16.--Forestland Management (Part 5)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Potential for damage to soil by fire		Potential for seedling mortality	
	Rating class and limiting features	Value	Rating class and limiting features	Value
8D: Hickory-----	None		Low	
8D2: Hickory-----	Moderate Texture/rock fragments	0.50	Low	
8D3: Hickory-----	Moderate Texture/rock fragments	0.50	Low	
8F: Hickory-----	Moderate Texture/rock fragments	0.50	Low	
17A: Keomah-----	Low Texture/rock fragments	0.10	High Wetness	1.00
36B: Tama-----	Low Texture/rock fragments	0.10	Low	
36C2: Tama-----	Low Texture/rock fragments	0.10	Low	
43A: Ipava-----	Low Texture/rock fragments	0.10	Low	
45A: Denny-----	Low Texture/rock fragments	0.10	High Wetness	1.00
46A: Herrick-----	Low Texture/rock fragments	0.10	Low	
50A: Viriden-----	None		High Wetness	1.00

Table 16.--Forestland Management (Part 5)--Continued

Map symbol and soil name	Potential for damage to soil by fire		Potential for seedling mortality	
	Rating class and limiting features	Value	Rating class and limiting features	Value
68A: Sable-----	None		High Wetness	1.00
86B: Osco-----	Low Texture/rock fragments	0.10	Low	
86C2: Osco-----	Low Texture/rock fragments	0.10	Low	
112A: Cowden-----	Low Texture/rock fragments	0.10	High Wetness	1.00
119C2: Elco-----	Moderate Texture/rock fragments	0.50	Low	
119D: Elco-----	None		Low	
119D2: Elco-----	Moderate Texture/rock fragments	0.50	Low	
119D3: Elco-----	None		Low	
127B: Harrison-----	Low Texture/rock fragments	0.10	Low	
127C2: Harrison-----	Low Texture/rock fragments	0.10	Low	
131C2: Alvin-----	Moderate Texture/rock fragments	0.50	Low	
131D2: Alvin-----	Moderate Texture/rock fragments	0.50	Low	
134A: Camden-----	Low Texture/rock fragments	0.10	Low	

Table 16.--Forestland Management (Part 5)--Continued

Map symbol and soil name	Potential for damage to soil by fire		Potential for seedling mortality	
	Rating class and limiting features	Value	Rating class and limiting features	Value
134B: Camden-----	Low Texture/rock fragments	0.10	Low	
134C2: Camden-----	Moderate Texture/rock fragments	0.50	Low	
138A: Shiloh-----	None		High Wetness	1.00
198A: Elburn-----	Low Texture/rock fragments	0.10	Low	
199A: Plano-----	Low Texture/rock fragments	0.10	Low	
199B: Plano-----	Low Texture/rock fragments	0.10	Low	
212C2: Thebes-----	Moderate Texture/rock fragments	0.50	Low	
242A: Kendall-----	Low Texture/rock fragments	0.10	High Wetness	1.00
244A: Hartsburg-----	None		High Wetness	1.00
249A: Edinburg-----	None		High Wetness	1.00
257A: Clarksdale-----	Low Texture/rock fragments	0.10	High Wetness	1.00
259C2: Assumption-----	Low Texture/rock fragments	0.10	Low	

Table 16.--Forestland Management (Part 5)--Continued

Map symbol and soil name	Potential for damage to soil by fire		Potential for seedling mortality	
	Rating class and limiting features	Value	Rating class and limiting features	Value
259D2: Assumption-----	Low Texture/rock fragments	0.10	Low	
279B: Rozetta-----	Low Texture/rock fragments	0.10	Low	
280C2: Fayette-----	Moderate Texture/rock fragments	0.50	Low	
280D2: Fayette-----	Moderate Texture/rock fragments	0.50	Low	
280D3: Fayette-----	None		Low	
533: Urban land-----	Not rated		Not rated	
536: Dumps-----	Not rated		Not rated	
549G: Marseilles-----	Low Texture/rock fragments	0.10	Low	
567C2: Elkhart-----	Low Texture/rock fragments	0.10	Low	
567D2: Elkhart-----	Low Texture/rock fragments	0.10	Low	
630C2: Navlys-----	Moderate Texture/rock fragments	0.50	Low	
630D2: Navlys-----	Moderate Texture/rock fragments	0.50	Low	
630D3: Navlys-----	None		Low	

Table 16.--Forestland Management (Part 5)--Continued

Map symbol and soil name	Potential for damage to soil by fire		Potential for seedling mortality	
	Rating class and limiting features	Value	Rating class and limiting features	Value
679B: Blackberry-----	Low Texture/rock fragments	0.10	Low	
684B: Broadwell-----	Low Texture/rock fragments	0.10	Low	
684C2: Broadwell-----	Low Texture/rock fragments	0.10	Low	
685B: Middletown-----	Low Texture/rock fragments	0.10	Low	
685C2: Middletown-----	Moderate Texture/rock fragments	0.50	Low	
705A: Buckhart-----	Low Texture/rock fragments	0.10	Low	
705B: Buckhart-----	Low Texture/rock fragments	0.10	Low	
712A: Spaulding-----	None		High Wetness Lime Soil reaction	1.00 0.50 0.50
801C: Orthents-----	None		Low	
830: Landfills-----	Not rated		Not rated	
862: Pits, sand-----	Not rated		Not rated	
864: Pits, quarries-----	Not rated		Not rated	
3073A: Ross-----	Low Texture/rock fragments	0.10	Low	

Table 16.--Forestland Management (Part 5)--Continued

Map symbol and soil name	Potential for damage to soil by fire		Potential for seedling mortality	
	Rating class and limiting features	Value	Rating class and limiting features	Value
3074A: Radford-----	Low Texture/rock fragments	0.10	Low	
3077A: Huntsville-----	Low Texture/rock fragments	0.10	Low	
3107A: Sawmill-----	None		High Wetness	1.00
3284A: Tice-----	None		Low	
3405A: Zook-----	None		High Wetness	1.00
3451A: Lawson-----	Low Texture/rock fragments	0.10	Low	
7037A: Worthen-----	Low Texture/rock fragments	0.10	Low	
7075B: Drury-----	Moderate Texture/rock fragments	0.50	Low	
7148A: Proctor-----	Low Texture/rock fragments	0.10	Low	
7242A: Kendall-----	Low Texture/rock fragments	0.10	High Wetness	1.00
8396A: Vesser-----	Low Texture/rock fragments	0.10	High Wetness	1.00

Table 17.--Recreation (Part 1)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Camp areas		Picnic areas		Playgrounds	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
8D: Hickory-----	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
8D2: Hickory-----	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
8D3: Hickory-----	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
8F: Hickory-----	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
17A: Keomah-----	Very limited Depth to saturated zone Restricted permeability	1.00 0.96	Somewhat limited Restricted permeability Depth to saturated zone	0.96 0.94	Very limited Depth to saturated zone Restricted permeability	1.00 0.96
36B: Tama-----	Not limited		Not limited		Somewhat limited Slope	0.28
36C2: Tama-----	Not limited		Not limited		Very limited Slope	1.00
43A: Ipava-----	Somewhat limited Depth to saturated zone Restricted permeability	0.98 0.21	Somewhat limited Depth to saturated zone Restricted permeability	0.75 0.21	Somewhat limited Depth to saturated zone Restricted permeability	0.98 0.21
45A: Denny-----	Very limited Depth to saturated zone Ponding Restricted permeability	1.00 1.00 0.96	Very limited Ponding Depth to saturated zone Restricted permeability	1.00 1.00 0.96	Very limited Depth to saturated zone Ponding Restricted permeability	1.00 1.00 0.96
46A: Herrick-----	Somewhat limited Depth to saturated zone Restricted permeability	0.98 0.21	Somewhat limited Depth to saturated zone Restricted permeability	0.75 0.21	Somewhat limited Depth to saturated zone Restricted permeability	0.98 0.21

Table 17.--Recreation (Part 1)--Continued

Map symbol and soil name	Camp areas		Picnic areas		Playgrounds	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
50A:						
Virden-----	Very limited		Very limited		Very limited	
	Depth to	1.00	Ponding	1.00	Depth to	1.00
	saturated zone		Depth to	1.00	saturated zone	
	Ponding	1.00	saturated zone		Ponding	1.00
	Restricted	0.21	permeability	0.21	Restricted	0.21
	permeability				permeability	
68A:						
Sable-----	Very limited		Very limited		Very limited	
	Depth to	1.00	Ponding	1.00	Depth to	1.00
	saturated zone		Depth to	1.00	saturated zone	
	Ponding	1.00	saturated zone	1.00	Ponding	1.00
86B:						
Oscos-----	Not limited		Not limited		Somewhat limited	
					Slope	0.28
86C2:						
Oscos-----	Not limited		Not limited		Very limited	
					Slope	1.00
112A:						
Cowden-----	Very limited		Very limited		Very limited	
	Depth to	1.00	Ponding	1.00	Depth to	1.00
	saturated zone		Depth to	1.00	saturated zone	
	Ponding	1.00	saturated zone		Ponding	1.00
	Restricted	0.96	permeability	0.96	Restricted	0.96
	permeability				permeability	
119C2:						
Elco-----	Somewhat limited		Somewhat limited		Very limited	
	Restricted	0.43	Restricted	0.43	Slope	1.00
	permeability		permeability		Restricted	0.43
					permeability	
119D:						
Elco-----	Somewhat limited		Somewhat limited		Very limited	
	Slope	0.96	Slope	0.96	Slope	1.00
	Restricted	0.43	Restricted	0.43	Restricted	0.43
	permeability		permeability		permeability	
119D2:						
Elco-----	Somewhat limited		Somewhat limited		Very limited	
	Slope	0.96	Slope	0.96	Slope	1.00
	Restricted	0.43	Restricted	0.43	Restricted	0.43
	permeability		permeability		permeability	
119D3:						
Elco-----	Somewhat limited		Somewhat limited		Very limited	
	Slope	0.96	Slope	0.96	Slope	1.00
	Restricted	0.43	Restricted	0.43	Restricted	0.43
	permeability		permeability		permeability	
127B:						
Harrison-----	Not limited		Not limited		Somewhat limited	
					Slope	0.28
127C2:						
Harrison-----	Not limited		Not limited		Very limited	
					Slope	1.00

Table 17.--Recreation (Part 1)--Continued

Map symbol and soil name	Camp areas		Picnic areas		Playgrounds	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
131C2: Alvin-----	Not limited		Not limited		Very limited Slope	1.00
131D2: Alvin-----	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
134A: Camden-----	Not limited		Not limited		Not limited	
134B: Camden-----	Not limited		Not limited		Somewhat limited Slope	0.28
134C2: Camden-----	Not limited		Not limited		Very limited Slope	1.00
138A: Shiloh-----	Very limited Depth to saturated zone Ponding Restricted permeability	1.00 1.00 0.21	Very limited Ponding Depth to saturated zone Restricted permeability	1.00 1.00 0.21	Very limited Depth to saturated zone Ponding Restricted permeability	1.00 1.00 0.21
198A: Elburn-----	Somewhat limited Depth to saturated zone	0.98	Somewhat limited Depth to saturated zone	0.75	Somewhat limited Depth to saturated zone	0.98
199A: Plano-----	Not limited		Not limited		Not limited	
199B: Plano-----	Not limited		Not limited		Somewhat limited Slope	0.28
212C2: Thebes-----	Not limited		Not limited		Very limited Slope	1.00
242A: Kendall-----	Very limited Depth to saturated zone	1.00	Somewhat limited Depth to saturated zone	0.94	Very limited Depth to saturated zone	1.00
244A: Hartsburg-----	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
249A: Edinburg-----	Very limited Depth to saturated zone Ponding Restricted permeability	1.00 1.00 0.96	Very limited Ponding Depth to saturated zone Restricted permeability	1.00 1.00 0.96	Very limited Depth to saturated zone Ponding Restricted permeability	1.00 1.00 0.96

Table 17.--Recreation (Part 1)--Continued

Map symbol and soil name	Camp areas		Picnic areas		Playgrounds	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
257A: Clarksdale-----	Very limited Depth to saturated zone Restricted permeability	1.00 0.21	Somewhat limited Depth to saturated zone Restricted permeability	0.94 0.21	Very limited Depth to saturated zone Restricted permeability	1.00 0.21
259C2: Assumption-----	Somewhat limited Restricted permeability	0.43	Somewhat limited Restricted permeability	0.43	Very limited Slope Restricted permeability	1.00 0.43
259D2: Assumption-----	Somewhat limited Slope Restricted permeability	0.96 0.43	Somewhat limited Slope Restricted permeability	0.96 0.43	Very limited Slope Restricted permeability	1.00 0.43
279B: Rozetta-----	Not limited		Not limited		Somewhat limited Slope	0.28
280C2: Fayette-----	Not limited		Not limited		Very limited Slope	1.00
280D2: Fayette-----	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
280D3: Fayette-----	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
533: Urban land-----	Not rated		Not rated		Not rated	
536: Dumps-----	Not rated		Not rated		Not rated	
549G: Marseilles-----	Very limited Slope Restricted permeability	1.00 0.96	Very limited Slope Restricted permeability	1.00 0.96	Very limited Slope Restricted permeability Depth to bedrock	1.00 0.96 0.10
567C2: Elkhart-----	Not limited		Not limited		Very limited Slope	1.00
567D2: Elkhart-----	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
630C2: Navlys-----	Not limited		Not limited		Very limited Slope	1.00

Table 17.--Recreation (Part 1)--Continued

Map symbol and soil name	Camp areas		Picnic areas		Playgrounds	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
630D2: Navlys-----	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
630D3: Navlys-----	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
679B: Blackberry-----	Not limited		Not limited		Somewhat limited Slope	0.28
684B: Broadwell-----	Not limited		Not limited		Somewhat limited Slope	0.28
684C2: Broadwell-----	Not limited		Not limited		Very limited Slope	1.00
685B: Middletown-----	Not limited		Not limited		Somewhat limited Slope	0.28
685C2: Middletown-----	Not limited		Not limited		Very limited Slope	1.00
705A: Buckhart-----	Not limited		Not limited		Not limited	
705B: Buckhart-----	Not limited		Not limited		Somewhat limited	
712A: Spaulding-----	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00
801C: Orthents-----	Not limited		Not limited		Very limited Slope	1.00
830: Landfills-----	Not rated		Not rated		Not rated	
862: Pits, sand-----	Not rated		Not rated		Not rated	
864: Pits, quarries-----	Not rated		Not rated		Not rated	
3073A: Ross-----	Very limited Flooding	1.00	Somewhat limited Flooding	0.40	Very limited Flooding	1.00

Table 17.--Recreation (Part 1)--Continued

Map symbol and soil name	Camp areas		Picnic areas		Playgrounds	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
3074A: Radford-----	Very limited Flooding Depth to saturated zone	1.00 0.98	Somewhat limited Depth to saturated zone Flooding	0.75 0.40	Very limited Flooding Depth to saturated zone	1.00 0.98
3077A: Huntsville-----	Very limited Flooding	1.00	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
3107A: Sawmill-----	Very limited Depth to saturated zone Flooding Ponding	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Flooding	1.00 1.00 0.40	Very limited Depth to saturated zone Flooding Ponding	1.00 1.00 1.00
3284A: Tice-----	Very limited Flooding Depth to saturated zone	1.00 0.98	Somewhat limited Depth to saturated zone Flooding	0.75 0.40	Very limited Flooding Depth to saturated zone	1.00 0.98
3405A: Zook-----	Very limited Depth to saturated zone Flooding Restricted permeability	1.00 1.00 0.96	Very limited Depth to saturated zone Restricted permeability Flooding	1.00 0.96 0.40	Very limited Depth to saturated zone Flooding Restricted permeability	1.00 1.00 0.96
3451A: Lawson-----	Very limited Flooding Depth to saturated zone	1.00 0.98	Somewhat limited Depth to saturated zone Flooding	0.75 0.40	Very limited Flooding Depth to saturated zone	1.00 0.98
7037A: Worthen-----	Very limited Flooding	1.00	Not limited		Not limited	
7075B: Drury-----	Very limited Flooding	1.00	Not limited		Somewhat limited Slope	0.28
7148A: Proctor-----	Very limited Flooding	1.00	Not limited		Not limited	
7242A: Kendall-----	Very limited Flooding Depth to saturated zone	1.00 1.00	Somewhat limited Depth to saturated zone	0.94	Very limited Depth to saturated zone	1.00
8396A: Vesser-----	Very limited Depth to saturated zone Flooding	1.00 1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Flooding	1.00 0.60

Table 18.--Recreation (Part 2)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Paths and trails		Off-road motorcycle trails		Golf fairways	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
8D: Hickory-----	Not limited		Not limited		Somewhat limited Slope	0.96
8D2: Hickory-----	Not limited		Not limited		Somewhat limited Slope	0.96
8D3: Hickory-----	Not limited		Not limited		Somewhat limited Slope	0.96
8F: Hickory-----	Very limited Slope	1.00	Somewhat limited Slope	0.02	Very limited Slope	1.00
17A: Keomah-----	Somewhat limited Depth to saturated zone	0.86	Somewhat limited Depth to saturated zone	0.86	Somewhat limited Depth to saturated zone	0.94
36B: Tama-----	Not limited		Not limited		Not limited	
36C2: Tama-----	Not limited		Not limited		Not limited	
43A: Ipava-----	Somewhat limited Depth to saturated zone	0.44	Somewhat limited Depth to saturated zone	0.44	Somewhat limited Depth to saturated zone	0.75
45A: Denny-----	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00
46A: Herrick-----	Somewhat limited Depth to saturated zone	0.44	Somewhat limited Depth to saturated zone	0.44	Somewhat limited Depth to saturated zone	0.75
50A: Virden-----	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00
68A: Sable-----	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00

Table 18.--Recreation (Part 2)--Continued

Map symbol and soil name	Paths and trails		Off-road motorcycle trails		Golf fairways	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
86B: Osco-----	Not limited		Not limited		Not limited	
86C2: Osco-----	Not limited		Not limited		Not limited	
112A: Cowden-----	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00
119C2: Elco-----	Not limited		Not limited		Not limited	
119D: Elco-----	Not limited		Not limited		Somewhat limited Slope	0.96
119D2: Elco-----	Not limited		Not limited		Somewhat limited Slope	0.96
119D3: Elco-----	Not limited		Not limited		Somewhat limited Slope	0.96
127B: Harrison-----	Not limited		Not limited		Not limited	
127C2: Harrison-----	Not limited		Not limited		Not limited	
131C2: Alvin-----	Not limited		Not limited		Not limited	
131D2: Alvin-----	Not limited		Not limited		Somewhat limited Slope	0.96
134A: Camden-----	Not limited		Not limited		Not limited	
134B: Camden-----	Not limited		Not limited		Not limited	
134C2: Camden-----	Not limited		Not limited		Not limited	
138A: Shiloh-----	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00
198A: Elburn-----	Somewhat limited Depth to saturated zone	0.44	Somewhat limited Depth to saturated zone	0.44	Somewhat limited Depth to saturated zone	0.75

Table 18.--Recreation (Part 2)--Continued

Map symbol and soil name	Paths and trails		Off-road motorcycle trails		Golf fairways	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
199A: Plano-----	Not limited		Not limited		Not limited	
199B: Plano-----	Not limited		Not limited		Not limited	
212C2: Thebes-----	Not limited		Not limited		Not limited	
242A: Kendall-----	Somewhat limited Depth to saturated zone	0.86	Somewhat limited Depth to saturated zone	0.86	Somewhat limited Depth to saturated zone	0.94
244A: Hartsburg-----	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00
249A: Edinburg-----	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00
257A: Clarksdale-----	Somewhat limited Depth to saturated zone	0.86	Somewhat limited Depth to saturated zone	0.86	Somewhat limited Depth to saturated zone	0.94
259C2: Assumption-----	Not limited		Not limited		Not limited	
259D2: Assumption-----	Not limited		Not limited		Somewhat limited Slope	0.96
279B: Rozetta-----	Not limited		Not limited		Not limited	
280C2: Fayette-----	Not limited		Not limited		Not limited	
280D2: Fayette-----	Not limited		Not limited		Somewhat limited Slope	0.96
280D3: Fayette-----	Not limited		Not limited		Somewhat limited Slope	0.96
533: Urban land-----	Not rated		Not rated		Not rated	
536: Dumps-----	Not rated		Not rated		Not rated	

Table 18.--Recreation (Part 2)--Continued

Map symbol and soil name	Paths and trails		Off-road motorcycle trails		Golf fairways	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
549G: Marseilles-----	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope Depth to bedrock	1.00 0.10
567C2: Elkhart-----	Not limited		Not limited		Not limited	
567D2: Elkhart-----	Not limited		Not limited		Somewhat limited Slope	0.96
630C2: Navlys-----	Not limited		Not limited		Not limited	
630D2: Navlys-----	Not limited		Not limited		Somewhat limited Slope	0.96
630D3: Navlys-----	Not limited		Not limited		Somewhat limited Slope	0.96
679B: Blackberry-----	Not limited		Not limited		Not limited	
684B: Broadwell-----	Not limited		Not limited		Not limited	
684C2: Broadwell-----	Not limited		Not limited		Not limited	
685B: Middletown-----	Not limited		Not limited		Not limited	
685C2: Middletown-----	Not limited		Not limited		Not limited	
705A: Buckhart-----	Not limited		Not limited		Not limited	
705B: Buckhart-----	Not limited		Not limited		Not limited	
712A: Spaulding-----	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00
801C: Orthents-----	Not limited		Not limited		Not limited	
830: Landfills-----	Not rated		Not rated		Not rated	
862: Pits, sand-----	Not rated		Not rated		Not rated	
864: Pits, quarries-----	Not rated		Not rated		Not rated	

Table 18.--Recreation (Part 2)--Continued

Map symbol and soil name	Paths and trails		Off-road motorcycle trails		Golf fairways	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
3073A: Ross-----	Somewhat limited Flooding	0.40	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
3074A: Radford-----	Somewhat limited Depth to saturated zone Flooding	0.44 0.40	Somewhat limited Depth to saturated zone Flooding	0.44 0.40	Very limited Flooding Depth to saturated zone	1.00 0.75
3077A: Huntsville-----	Somewhat limited Flooding	0.40	Somewhat limited Flooding	0.40	Very limited Flooding	1.00
3107A: Sawmill-----	Very limited Depth to saturated zone Ponding Flooding	1.00 1.00 0.40	Very limited Depth to saturated zone Ponding Flooding	1.00 1.00 0.40	Very limited Ponding Flooding Depth to saturated zone	1.00 1.00 1.00
3284A: Tice-----	Somewhat limited Depth to saturated zone Flooding	0.44 0.40	Somewhat limited Depth to saturated zone Flooding	0.44 0.40	Very limited Flooding Depth to saturated zone	1.00 0.75
3405A: Zook-----	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Depth to saturated zone Flooding	1.00 0.40	Very limited Flooding Depth to saturated zone	1.00 1.00
3451A: Lawson-----	Somewhat limited Depth to saturated zone Flooding	0.44 0.40	Somewhat limited Depth to saturated zone Flooding	0.44 0.40	Very limited Flooding Depth to saturated zone	1.00 0.75
7037A: Worthen-----	Not limited		Not limited		Not limited	
7075B: Drury-----	Not limited		Not limited		Not limited	
7148A: Proctor-----	Not limited		Not limited		Not limited	
7242A: Kendall-----	Somewhat limited Depth to saturated zone	0.86	Somewhat limited Depth to saturated zone	0.86	Somewhat limited Depth to saturated zone	0.94
8396A: Vesser-----	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Flooding	1.00 0.60

Table 19.--Wildlife Habitat

(See text for definitions of terms used in this table. Absence of an entry indicates that no rating is applicable.)

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hardwood trees	Conif- erous plants	Wetland plants	Shallow water areas	Openland wildlife	Woodland wildlife	Wetland wildlife
8D: Hickory-----	Fair	Good	Good	Good	Good	Very poor.	Very poor.	Good	Good	Very poor.
8D2: Hickory-----	Fair	Good	Good	Good	Good	Very poor.	Very poor.	Good	Good	Very poor.
8D3: Hickory-----	Fair	Good	Good	Good	Good	Very poor.	Very poor.	Good	Good	Very poor.
8F: Hickory-----	Very poor.	Fair	Good	Good	Good	Very poor.	Very poor.	Fair	Good	Very poor.
17A: Keomah-----	Fair	Good	Good	Good	Good	Fair	Fair	Good	Good	Fair.
36B: Tama-----	Good	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
36C2: Tama-----	Fair	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
43A: Ipava-----	Fair	Good	Good	Good	Good	Fair	Fair	Good	Good	Fair.
45A: Denny-----	Fair	Fair	Fair	Fair	Fair	Good	Good	Fair	Fair	Good.
46A: Herrick-----	Fair	Good	Good	Good	Good	Fair	Fair	Good	Good	Fair.
50A: Virden-----	Fair	Fair	Fair	Fair	Fair	Good	Good	Fair	Fair	Good.
68A: Sable-----	Fair	Fair	Fair	Fair	Fair	Good	Good	Fair	Fair	Good.
86B: Osco-----	Good	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
86C2: Osco-----	Fair	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
112A: Cowden-----	Fair	Fair	Fair	Fair	Fair	Good	Good	Fair	Fair	Good.
119C2: Elco-----	Fair	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.

Table 19.--Wildlife Habitat--Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hardwood trees	Conif- erous plants	Wetland plants	Shallow water areas	Openland wildlife	Woodland wildlife	Wetland wildlife
119D: Elco-----	Fair	Good	Good	Good	Good	Very poor.	Very poor.	Good	Good	Very poor.
119D2: Elco-----	Fair	Good	Good	Good	Good	Very poor.	Very poor.	Good	Good	Very poor.
119D3: Elco-----	Fair	Good	Good	Good	Good	Very poor.	Very poor.	Good	Good	Very poor.
127B: Harrison-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor.
127C2: Harrison-----	Fair	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
131C2: Alvin-----	Fair	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
131D2: Alvin-----	Fair	Good	Good	Good	Good	Very poor.	Very poor.	Good	Good	Very poor.
134A: Camden-----	Good	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
134B: Camden-----	Good	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
134C2: Camden-----	Fair	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
138A: Shiloh-----	Fair	Fair	Fair	Fair	Poor	Good	Good	Fair	Fair	Good.
198A: Elburn-----	Fair	Good	Good	Good	Good	Fair	Fair	Good	Good	Fair.
199A: Plano-----	Good	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
199B: Plano-----	Good	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
212C2: Thebes-----	Fair	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
242A: Kendall-----	Fair	Good	Good	Good	Good	Fair	Fair	Good	Good	Fair.

Table 19.--Wildlife Habitat--Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hardwood trees	Conif- erous plants	Wetland plants	Shallow water areas	Openland wildlife	Woodland wildlife	Wetland wildlife
244A: Hartsburg-----	Fair	Fair	Fair	Fair	Fair	Good	Good	Fair	Fair	Good.
249A: Edinburg-----	Fair	Fair	Fair	Fair	Fair	Good	Good	Fair	Fair	Good.
257A: Clarksdale-----	Fair	Good	Good	Good	Good	Fair	Fair	Good	Good	Fair.
259C2: Assumption-----	Fair	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
259D2: Assumption-----	Fair	Good	Good	Good	Good	Very poor.	Very poor.	Good	Good	Very poor.
279B: Rozetta-----	Good	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
280C2: Fayette-----	Fair	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
280D2: Fayette-----	Fair	Good	Good	Good	Good	Very poor.	Very poor.	Good	Good	Very poor.
280D3: Fayette-----	Fair	Good	Good	Good	Good	Very poor.	Very poor.	Good	Good	Very poor.
533: Urban land.										
536: Dumps.										
549G: Marseilles-----	Very poor.	Poor	Good	Good	Good	Very poor.	Very poor.	Poor	Good	Very poor.
567C2: Elkhart-----	Fair	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
567D2: Elkhart-----	Fair	Good	Good	Good	Good	Very poor.	Very poor.	Good	Good	Very poor.
630C2: Navlys-----	Fair	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
630D2: Navlys-----	Fair	Good	Good	Good	Good	Very poor.	Very poor.	Good	Good	Very poor.

Table 19.--Wildlife Habitat--Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hardwood trees	Conif- erous plants	Wetland plants	Shallow water areas	Openland wildlife	Woodland wildlife	Wetland wildlife
630D3: Navlys-----	Fair	Good	Good	Good	Good	Very poor.	Very poor.	Good	Good	Very poor.
679B: Blackberry-----	Good	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
684B: Broadwell-----	Good	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
684C2: Broadwell-----	Fair	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
685B: Middletown-----	Good	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
685C2: Middletown-----	Fair	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
705A: Buckhart-----	Good	Good	Good	Good	Good	Poor	Poor	Good	Good	Poor.
705B: Buckhart-----	Good	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
712A: Spaulding-----	Fair	Fair	Fair	Fair	Fair	Good	Good	Fair	Fair	Good.
801C: Orthents-----	Fair	Good	Good	Good	Good	Very poor.	Very poor.	Good	Good	Very poor.
830: Landfills.										
862: Pits, sand.										
864: Pits, quarries.										
3073A: Ross-----	Poor	Fair	Fair	Good	Good	Poor	Very poor.	Fair	Good	Very poor.
3074A: Radford-----	Poor	Fair	Fair	Good	Good	Fair	Fair	Fair	Good	Fair.
3077A: Huntsville-----	Poor	Fair	Fair	Good	Good	Poor	Very poor.	Fair	Good	Very poor.
3107A: Sawmill-----	Poor	Fair	Fair	Fair	Fair	Good	Good	Fair	Fair	Good.

Table 19.--Wildlife Habitat--Continued

Map symbol and soil name	Potential for habitat elements							Potential as habitat for--		
	Grain and seed crops	Grasses and legumes	Wild herba- ceous plants	Hardwood trees	Conif- erous plants	Wetland plants	Shallow water areas	Openland wildlife	Woodland wildlife	Wetland wildlife
3284A: Tice-----	Poor	Fair	Fair	Good	Good	Fair	Fair	Fair	Good	Fair.
3405A: Zook-----	Poor	Fair	Fair	Fair	Fair	Good	Good	Fair	Fair	Good.
3451A: Lawson-----	Poor	Fair	Fair	Good	Good	Fair	Fair	Fair	Good	Fair.
7037A: Worthen-----	Good	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
7075B: Drury-----	Good	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
7148A: Proctor-----	Good	Good	Good	Good	Good	Poor	Very poor.	Good	Good	Very poor.
7242A: Kendall-----	Fair	Good	Good	Good	Good	Fair	Fair	Good	Good	Fair.
8396A: Vesser-----	Fair	Fair	Fair	Fair	Fair	Good	Good	Fair	Fair	Good.

Table 20.--Building Site Development (Part 1)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Dwellings without basements		Dwellings with basements		Small commercial buildings	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
8D, 8D2, 8D3: Hickory-----	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00
	Shrink-swell	0.50	Shrink-swell	0.50	Shrink-swell	0.50
8F: Hickory-----	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
	Shrink-swell	0.50	Shrink-swell	0.50	Shrink-swell	0.50
17A: Keomah-----	Very limited Shrink-swell	1.00	Very limited Depth to	1.00	Very limited Shrink-swell	1.00
	Depth to saturated zone	1.00	saturated zone		Depth to saturated zone	1.00
36B: Tama-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
36C2: Tama-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Slope	0.96
					Shrink-swell	0.50
43A: Ipava-----	Very limited Shrink-swell	1.00	Very limited Depth to	1.00	Very limited Shrink-swell	1.00
	Depth to saturated zone	0.98	saturated zone		Depth to saturated zone	0.98
			Shrink-swell	0.50		
45A: Denny-----	Very limited Ponding	1.00	Very limited Ponding	1.00	Very limited Ponding	1.00
	Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
	Shrink-swell	1.00	Shrink-swell	0.50	Shrink-swell	1.00
46A: Herrick-----	Very limited Shrink-swell	1.00	Very limited Depth to	1.00	Very limited Shrink-swell	1.00
	Depth to saturated zone	0.98	saturated zone		Depth to saturated zone	0.98
			Shrink-swell	1.00		
50A: Virden-----	Very limited Ponding	1.00	Very limited Ponding	1.00	Very limited Ponding	1.00
	Depth to saturated zone	1.00	Depth to saturated zone	1.00	Depth to saturated zone	1.00
	Shrink-swell	1.00	Shrink-swell	1.00	Shrink-swell	1.00

Table 20.--Building Site Development (Part 1)--Continued

Map symbol and soil name	Dwellings without basements		Dwellings with basements		Small commercial buildings	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
68A: Sable-----	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 0.50
86B: Osco-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Depth to saturated zone	0.50 0.15	Somewhat limited Shrink-swell	0.50
86C2: Osco-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Depth to saturated zone	0.50 0.15	Somewhat limited Slope Shrink-swell	0.96 0.50
112A: Cowden-----	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 1.00
119C2: Elco-----	Somewhat limited Shrink-swell	0.50	Very limited Shrink-swell Depth to saturated zone	1.00 0.99	Somewhat limited Slope Shrink-swell	0.96 0.50
119D, 119D2, 119D3: Elco-----	Somewhat limited Slope Shrink-swell	0.96 0.50	Very limited Shrink-swell Depth to saturated zone Slope	1.00 0.99 0.96	Very limited Slope Shrink-swell	1.00 0.50
127B: Harrison-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to saturated zone Shrink-swell	0.99 0.50	Somewhat limited Shrink-swell	0.50
127C2: Harrison-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to saturated zone Shrink-swell	0.99 0.50	Somewhat limited Slope Shrink-swell	0.96 0.50
131C2: Alvin-----	Not limited		Not limited		Somewhat limited Slope	0.96
131D2: Alvin-----	Somewhat limited Slope	0.96	Somewhat limited Slope	0.96	Very limited Slope	1.00

Table 20.--Building Site Development (Part 1)--Continued

Map symbol and soil name	Dwellings without basements		Dwellings with basements		Small commercial buildings	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
134A: Camden-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
134B: Camden-----	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Shrink-swell	0.50
134C2: Camden-----	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Slope Shrink-swell	0.96 0.50
138A: Shiloh-----	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 1.00
198A: Elburn-----	Somewhat limited Depth to saturated zone Shrink-swell	0.98 0.50	Very limited Depth to saturated zone Shrink-swell	1.00 0.50	Somewhat limited Depth to saturated zone Shrink-swell	0.98 0.50
199A: Plano-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
199B: Plano-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
212C2: Thebes-----	Somewhat limited Shrink-swell	0.50	Not limited		Somewhat limited Slope Shrink-swell	0.96 0.50
242A: Kendall-----	Very limited Depth to saturated zone Shrink-swell	1.00 0.50	Very limited Depth to saturated zone Shrink-swell	1.00 0.50	Very limited Depth to saturated zone Shrink-swell	1.00 0.50
244A: Hartsburg-----	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 0.50
249A: Edinburg-----	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 1.00

Table 20.--Building Site Development (Part 1)--Continued

Map symbol and soil name	Dwellings without basements		Dwellings with basements		Small commercial buildings	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
257A: Clarksdale-----	Very limited Shrink-swell Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Shrink-swell	1.00 1.00	Very limited Shrink-swell Depth to saturated zone	1.00 1.00
259C2: Assumption-----	Very limited Shrink-swell	1.00	Very limited Shrink-swell Depth to saturated zone	1.00 0.99	Very limited Shrink-swell Slope	1.00 0.96
259D2: Assumption-----	Somewhat limited Slope Shrink-swell	0.96 0.50	Very limited Shrink-swell Depth to saturated zone Slope	1.00 0.99 0.96	Very limited Slope Shrink-swell	1.00 0.50
279B: Rozetta-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell Depth to saturated zone	0.50 0.15	Somewhat limited Shrink-swell	0.50
280C2: Fayette-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Slope Shrink-swell	0.96 0.50
280D2, 280D3: Fayette-----	Somewhat limited Slope Shrink-swell	0.96 0.50	Somewhat limited Slope Shrink-swell	0.96 0.50	Very limited Slope Shrink-swell	1.00 0.50
533: Urban land-----	Not rated		Not rated		Not rated	
536: Dumps-----	Not rated		Not rated		Not rated	
549G: Marseilles-----	Very limited Slope Shrink-swell	1.00 0.50	Very limited Slope Shrink-swell Depth to soft bedrock	1.00 0.50 0.10	Very limited Slope Shrink-swell	1.00 0.50
567C2: Elkhart-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to saturated zone	0.16	Somewhat limited Slope Shrink-swell	0.96 0.50
567D2: Elkhart-----	Somewhat limited Slope Shrink-swell	0.96 0.50	Somewhat limited Slope Depth to saturated zone	0.96 0.16	Very limited Slope Shrink-swell	1.00 0.50

Table 20.--Building Site Development (Part 1)--Continued

Map symbol and soil name	Dwellings without basements		Dwellings with basements		Small commercial buildings	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
630C2: Navlys-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to saturated zone	0.15	Somewhat limited Slope Shrink-swell	0.96 0.50
630D2, 630D3: Navlys-----	Somewhat limited Slope Shrink-swell	0.96 0.50	Somewhat limited Slope Depth to saturated zone	0.96 0.15	Very limited Slope Shrink-swell	1.00 0.50
679B: Blackberry-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to saturated zone Shrink-swell	0.99 0.50	Somewhat limited Shrink-swell	0.50
684B: Broadwell-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
684C2: Broadwell-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Slope Shrink-swell	0.96 0.50
685B: Middletown-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50
685C2: Middletown-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Slope Shrink-swell	0.96 0.50
705A: Buckhart-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to saturated zone Shrink-swell	0.99 0.50	Somewhat limited Shrink-swell	0.50
705B: Buckhart-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Depth to saturated zone Shrink-swell	0.99 0.50	Somewhat limited Shrink-swell	0.50
712A: Spaulding-----	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Ponding Depth to saturated zone Shrink-swell	1.00 1.00 0.50
801C: Orthents-----	Somewhat limited Shrink-swell	0.50	Somewhat limited Shrink-swell	0.50	Somewhat limited Slope Shrink-swell	0.96 0.50
830: Landfills-----	Not rated		Not rated		Not rated	

Table 20.--Building Site Development (Part 1)--Continued

Map symbol and soil name	Dwellings without basements		Dwellings with basements		Small commercial buildings	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
862: Pits, sand-----	Not rated		Not rated		Not rated	
864: Pits, quarries-----	Not rated		Not rated		Not rated	
3073A: Ross-----	Very limited Flooding	1.00	Very limited Flooding Depth to saturated zone	1.00 0.15	Very limited Flooding	1.00
3074A: Radford-----	Very limited Flooding Depth to saturated zone	1.00 0.98	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Flooding Depth to saturated zone	1.00 0.98
3077A: Huntsville-----	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding Shrink-swell Depth to saturated zone	1.00 0.50 0.15	Very limited Flooding Shrink-swell	1.00 0.50
3107A: Sawmill-----	Very limited Ponding Flooding Depth to saturated zone Shrink-swell	1.00 1.00 1.00 0.50	Very limited Ponding Flooding Depth to saturated zone Shrink-swell	1.00 1.00 1.00 0.50	Very limited Ponding Flooding Depth to saturated zone Shrink-swell	1.00 1.00 1.00 0.50
3284A: Tice-----	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 0.98 0.50	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 0.98 0.50
3405A: Zook-----	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 1.00
3451A: Lawson-----	Very limited Flooding Depth to saturated zone	1.00 0.98	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Flooding Depth to saturated zone	1.00 0.98
7037A: Worthen-----	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00

Table 20.--Building Site Development (Part 1)--Continued

Map symbol and soil name	Dwellings without basements		Dwellings with basements		Small commercial buildings	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
7075B: Drury-----	Very limited Flooding	1.00	Very limited Flooding	1.00	Very limited Flooding	1.00
7148A: Proctor-----	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding Shrink-swell	1.00 0.50	Very limited Flooding Shrink-swell	1.00 0.50
7242A: Kendall-----	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50
8396A: Vesser-----	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50	Very limited Flooding Depth to saturated zone Shrink-swell	1.00 1.00 0.50

Table 21.--Building Site Development (Part 2)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Local roads and streets		Shallow excavations		Lawns and landscaping	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
8D: Hickory-----	Very limited		Somewhat limited		Somewhat limited	
	Low strength	1.00	Slope	0.96	Slope	0.96
	Slope	0.96	Cutbanks cave	0.10		
	Shrink-swell	0.50				
	Frost action	0.50				
8D2: Hickory-----	Very limited		Somewhat limited		Somewhat limited	
	Low strength	1.00	Slope	0.96	Slope	0.96
	Slope	0.96	Cutbanks cave	0.10		
	Shrink-swell	0.50				
	Frost action	0.50				
8D3: Hickory-----	Very limited		Somewhat limited		Somewhat limited	
	Low strength	1.00	Slope	0.96	Slope	0.96
	Slope	0.96	Cutbanks cave	0.10		
	Shrink-swell	0.50				
	Frost action	0.50				
8F: Hickory-----	Very limited		Very limited		Very limited	
	Slope	1.00	Slope	1.00	Slope	1.00
	Low strength	1.00	Cutbanks cave	0.10		
	Shrink-swell	0.50				
	Frost action	0.50				
17A: Keomah-----	Very limited		Very limited		Somewhat limited	
	Frost action	1.00	Depth to	1.00	Depth to	0.94
	Low strength	1.00	saturated zone		saturated zone	
	Shrink-swell	1.00	Cutbanks cave	0.10		
	Depth to	0.94				
	saturated zone					
36B: Tama-----	Very limited		Somewhat limited		Not limited	
	Frost action	1.00	Cutbanks cave	0.10		
	Low strength	1.00				
	Shrink-swell	0.50				
36C2: Tama-----	Very limited		Somewhat limited		Not limited	
	Frost action	1.00	Cutbanks cave	0.10		
	Low strength	1.00				
	Shrink-swell	0.50				
43A: Ipava-----	Very limited		Very limited		Somewhat limited	
	Frost action	1.00	Depth to	1.00	Depth to	0.75
	Low strength	1.00	saturated zone		saturated zone	
	Shrink-swell	1.00	Cutbanks cave	0.10		
	Depth to	0.75				
	saturated zone					

Table 21.--Building Site Development (Part 2)--Continued

Map symbol and soil name	Local roads and streets		Shallow excavations		Lawns and landscaping	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
45A: Denny-----	Very limited Ponding Depth to saturated zone Frost action Low strength Shrink-swell	 1.00 1.00 1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Cutbanks cave	 1.00 1.00 0.10	Very limited Ponding Depth to saturated zone	 1.00 1.00
46A: Herrick-----	Very limited Frost action Low strength Shrink-swell Depth to saturated zone	 1.00 1.00 1.00 0.75 	Very limited Depth to saturated zone Cutbanks cave	 1.00 0.10	Somewhat limited Depth to saturated zone	 0.75
50A: Virden-----	Very limited Ponding Depth to saturated zone Frost action Low strength Shrink-swell	 1.00 1.00 1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Cutbanks cave	 1.00 1.00 0.10	Very limited Ponding Depth to saturated zone	 1.00 1.00
68A: Sable-----	Very limited Ponding Depth to saturated zone Frost action Low strength Shrink-swell	 1.00 1.00 1.00 1.00 0.50	Very limited Ponding Depth to saturated zone Cutbanks cave	 1.00 1.00 0.10	Very limited Ponding Depth to saturated zone	 1.00 1.00
86B: Osco-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Somewhat limited Depth to saturated zone Cutbanks cave	 0.15 0.10	Not limited	
86C2: Osco-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Somewhat limited Depth to saturated zone Cutbanks cave	 0.15 0.10	Not limited	
112A: Cowden-----	Very limited Ponding Depth to saturated zone Frost action Low strength Shrink-swell	 1.00 1.00 1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Cutbanks cave	 1.00 1.00 0.10	Very limited Ponding Depth to saturated zone	 1.00 1.00
119C2: Elco-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Somewhat limited Depth to saturated zone Cutbanks cave	 0.99 0.10	Not limited	

Table 21.--Building Site Development (Part 2)--Continued

Map symbol and soil name	Local roads and streets		Shallow excavations		Lawns and landscaping	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
119D: Elco-----	Very limited Frost action Low strength Slope Shrink-swell	 1.00 1.00 0.96 0.50	Somewhat limited Depth to saturated zone Slope Cutbanks cave	 0.99 0.96 0.10	Somewhat limited Slope	 0.96
119D2: Elco-----	Very limited Frost action Low strength Slope Shrink-swell	 1.00 1.00 0.96 0.50	Somewhat limited Depth to saturated zone Slope Cutbanks cave	 0.99 0.96 0.10	Somewhat limited Slope	 0.96
119D3: Elco-----	Very limited Frost action Low strength Slope Shrink-swell	 1.00 1.00 0.96 0.50	Somewhat limited Depth to saturated zone Slope Cutbanks cave	 0.99 0.96 0.10	Somewhat limited Slope	 0.96
127B: Harrison-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Somewhat limited Depth to saturated zone Cutbanks cave	 0.99 0.10	Not limited	
127C2: Harrison-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Somewhat limited Depth to saturated zone Cutbanks cave	 0.99 0.10	Not limited	
131C2: Alvin-----	Somewhat limited Frost action	 0.50	Very limited Cutbanks cave	 1.00	Not limited	
131D2: Alvin-----	Somewhat limited Slope Frost action	 0.96 0.50	Very limited Cutbanks cave Slope	 1.00 0.96	Somewhat limited Slope	 0.96
134A: Camden-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Very limited Cutbanks cave	 1.00	Not limited	
134B: Camden-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Somewhat limited Cutbanks cave	 0.10	Not limited	
134C2: Camden-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Very limited Cutbanks cave	 1.00	Not limited	

Table 21.--Building Site Development (Part 2)--Continued

Map symbol and soil name	Local roads and streets		Shallow excavations		Lawns and landscaping	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
138A: Shiloh-----	Very limited Ponding Depth to saturated zone Frost action Low strength Shrink-swell	 1.00 1.00 1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Cutbanks cave	 1.00 1.00 0.10	Very limited Ponding Depth to saturated zone	 1.00 1.00
198A: Elburn-----	Very limited Frost action Low strength Depth to saturated zone Shrink-swell	 1.00 1.00 0.75 0.50	Very limited Depth to saturated zone Cutbanks cave	 1.00 1.00	Somewhat limited Depth to saturated zone	 0.75
199A: Plano-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Very limited Cutbanks cave	 1.00	Not limited	
199B: Plano-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Very limited Cutbanks cave	 1.00	Not limited	
212C2: Thebes-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Very limited Cutbanks cave	 1.00	Not limited	
242A: Kendall-----	Very limited Frost action Low strength Depth to saturated zone Shrink-swell	 1.00 1.00 0.94 0.50	Very limited Depth to saturated zone Cutbanks cave	 1.00 0.10	Somewhat limited Depth to saturated zone	 0.94
244A: Hartsburg-----	Very limited Ponding Depth to saturated zone Frost action Low strength Shrink-swell	 1.00 1.00 1.00 1.00 0.50	Very limited Ponding Depth to saturated zone Cutbanks cave	 1.00 1.00 0.10	Very limited Ponding Depth to saturated zone	 1.00 1.00
249A: Edinburg-----	Very limited Ponding Depth to saturated zone Frost action Low strength Shrink-swell	 1.00 1.00 1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Cutbanks cave	 1.00 1.00 0.10	Very limited Ponding Depth to saturated zone	 1.00 1.00

Table 21.--Building Site Development (Part 2)--Continued

Map symbol and soil name	Local roads and streets		Shallow excavations		Lawns and landscaping	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
257A: Clarksdale-----	Very limited Frost action Low strength Shrink-swell Depth to saturated zone	 1.00 1.00 1.00 0.94 0.94	Very limited Depth to saturated zone Cutbanks cave	 1.00 0.10	Somewhat limited Depth to saturated zone	 0.94
259C2: Assumption-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 1.00	Somewhat limited Depth to saturated zone Cutbanks cave	 0.99 0.10	Not limited	
259D2: Assumption-----	Very limited Frost action Low strength Slope Shrink-swell	 1.00 1.00 0.96 0.50	Somewhat limited Depth to saturated zone Slope Cutbanks cave	 0.99 0.96 0.10	Somewhat limited Slope	 0.96
279B: Rozetta-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Somewhat limited Depth to saturated zone Cutbanks cave	 0.15 0.10	Not limited	
280C2: Fayette-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Somewhat limited Cutbanks cave	 0.10	Not limited	
280D2: Fayette-----	Very limited Frost action Low strength Slope Shrink-swell	 1.00 1.00 0.96 0.50	Somewhat limited Slope Cutbanks cave	 0.96 0.10	Somewhat limited Slope	 0.96
280D3: Fayette-----	Very limited Frost action Low strength Slope Shrink-swell	 1.00 1.00 0.96 0.50	Somewhat limited Slope Cutbanks cave	 0.96 0.10	Somewhat limited Slope	 0.96
533: Urban land-----	Not rated		Not rated		Not rated	
536: Dumps-----	Not rated		Not rated		Not rated	
549G: Marseilles-----	Very limited Slope Frost action Low strength Shrink-swell	 1.00 1.00 1.00 0.50	Very limited Slope Cutbanks cave Depth to soft bedrock	 1.00 0.10 0.10	Very limited Slope Depth to bedrock	 1.00 0.10

Table 21.--Building Site Development (Part 2)--Continued

Map symbol and soil name	Local roads and streets		Shallow excavations		Lawns and landscaping	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
567C2: Elkhart-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Somewhat limited Depth to saturated zone Cutbanks cave	 0.16 0.10	Not limited	
567D2: Elkhart-----	Very limited Frost action Low strength Slope Shrink-swell	 1.00 1.00 0.96 0.50	Somewhat limited Slope Depth to saturated zone Cutbanks cave	 0.96 0.16 0.10	Somewhat limited Slope	 0.96
630C2: Navlys-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Somewhat limited Depth to saturated zone Cutbanks cave	 0.15 0.10	Not limited	
630D2: Navlys-----	Very limited Frost action Low strength Slope Shrink-swell	 1.00 1.00 0.96 0.50	Somewhat limited Slope Depth to saturated zone Cutbanks cave	 0.96 0.15 0.10	Somewhat limited Slope	 0.96
630D3: Navlys-----	Very limited Frost action Low strength Slope Shrink-swell	 1.00 1.00 0.96 0.50	Somewhat limited Slope Depth to saturated zone Cutbanks cave	 0.96 0.15 0.10	Somewhat limited Slope	 0.96
679B: Blackberry-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Somewhat limited Depth to saturated zone Cutbanks cave	 0.99 0.10	Not limited	
684B: Broadwell-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Very limited Cutbanks cave	 1.00	Not limited	
684C2: Broadwell-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Very limited Cutbanks cave	 1.00	Not limited	
685B: Middletown-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Very limited Cutbanks cave	 1.00	Not limited	

Table 21.--Building Site Development (Part 2)--Continued

Map symbol and soil name	Local roads and streets		Shallow excavations		Lawns and landscaping	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
685C2: Middletown-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Very limited Cutbanks cave	 1.00	Not limited	
705A: Buckhart-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Somewhat limited Depth to saturated zone Cutbanks cave	 0.99 0.10	Not limited	
705B: Buckhart-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Somewhat limited Depth to saturated zone Cutbanks cave	 0.99 0.10	Not limited	
712A: Spaulding-----	Very limited Ponding Depth to saturated zone Frost action Low strength Shrink-swell	 1.00 1.00 1.00 1.00 0.50	Very limited Ponding Depth to saturated zone Cutbanks cave	 1.00 1.00 0.10	Very limited Ponding Depth to saturated zone	 1.00 1.00
801C: Orthents-----	Very limited Frost action Low strength Shrink-swell	 1.00 1.00 0.50	Somewhat limited Cutbanks cave	 0.10	Not limited	
830: Landfills-----	Not rated		Not rated		Not rated	
862: Pits, sand-----	Not rated		Not rated		Not rated	
864: Pits, quarries-----	Not rated		Not rated		Not rated	
3073A: Ross-----	Very limited Flooding Low strength Frost action	 1.00 0.78 0.50	Somewhat limited Flooding Depth to saturated zone Cutbanks cave	 0.80 0.15 0.10	Very limited Flooding	 1.00
3074A: Radford-----	Very limited Frost action Flooding Low strength Depth to saturated zone	 1.00 1.00 1.00 0.75	Very limited Depth to saturated zone Flooding Cutbanks cave	 1.00 0.80 0.10	Very limited Flooding Depth to saturated zone	 1.00 0.75

Table 21.--Building Site Development (Part 2)--Continued

Map symbol and soil name	Local roads and streets		Shallow excavations		Lawns and landscaping	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
3077A: Huntsville-----	Very limited Frost action Flooding Low strength Shrink-swell	 1.00 1.00 1.00 0.50	Somewhat limited Flooding Depth to saturated zone Cutbanks cave	 0.80 0.15 0.10	Very limited Flooding	 1.00
3107A: Sawmill-----	Very limited Ponding Depth to saturated zone Frost action Flooding Low strength	 1.00 1.00 1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Flooding Cutbanks cave	 1.00 1.00 0.80 0.10	Very limited Ponding Flooding Depth to saturated zone	 1.00 1.00 1.00
3284A: Tice-----	Very limited Frost action Flooding Low strength Depth to saturated zone Shrink-swell	 1.00 1.00 1.00 0.75 0.50	Very limited Depth to saturated zone Flooding Cutbanks cave	 1.00 0.80 0.10	Very limited Flooding Depth to saturated zone	 1.00 0.75
3405A: Zook-----	Very limited Depth to saturated zone Frost action Flooding Low strength Shrink-swell	 1.00 1.00 1.00 1.00 1.00	Very limited Depth to saturated zone Flooding Cutbanks cave Too clayey	 1.00 0.80 0.10 0.01	Very limited Flooding Depth to saturated zone	 1.00 1.00
3451A: Lawson-----	Very limited Frost action Flooding Depth to saturated zone	 1.00 1.00 0.75 	Very limited Depth to saturated zone Flooding Cutbanks cave	 1.00 0.80 0.10	Very limited Flooding Depth to saturated zone	 1.00 0.75
7037A: Worthen-----	Very limited Frost action Low strength Flooding	 1.00 1.00 0.40	Somewhat limited Cutbanks cave	 0.10	Not limited	
7075B: Drury-----	Very limited Frost action Low strength Flooding	 1.00 1.00 0.40	Somewhat limited Cutbanks cave	 0.10	Not limited	
7148A: Proctor-----	Very limited Frost action Low strength Shrink-swell Flooding	 1.00 1.00 0.50 0.40	Somewhat limited Cutbanks cave	 0.10	Not limited	

Table 21.--Building Site Development (Part 2)--Continued

Map symbol and soil name	Local roads and streets		Shallow excavations		Lawns and landscaping	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
7242A: Kendall-----	Very limited		Very limited		Somewhat limited	
	Frost action	1.00	Depth to	1.00	Depth to	0.94
	Low strength	1.00	saturated zone		saturated zone	
	Depth to	0.94	Cutbanks cave	0.10		
	saturated zone					
	Shrink-swell	0.50				
	Flooding	0.40				
8396A: Vesser-----	Very limited		Very limited		Very limited	
	Depth to	1.00	Depth to	1.00	Depth to	1.00
	saturated zone		saturated zone		saturated zone	
	Frost action	1.00	Flooding	0.60	Flooding	0.60
	Flooding	1.00	Cutbanks cave	0.10		
	Low strength	1.00				
	Shrink-swell	0.50				

Table 22.--Sanitary Facilities (Part 1)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Septic tank absorption fields		Sewage lagoons	
	Rating class and limiting features	Value	Rating class and limiting features	Value
8D: Hickory-----	Somewhat limited Slope	0.96	Very limited Slope	1.00
	Restricted permeability	0.46	Seepage	0.53
8D2: Hickory-----	Somewhat limited Slope	0.96	Very limited Slope	1.00
	Restricted permeability	0.46	Seepage	0.53
8D3: Hickory-----	Somewhat limited Slope	0.96	Very limited Slope	1.00
	Restricted permeability	0.46	Seepage	0.53
8F: Hickory-----	Very limited Slope	1.00	Very limited Slope	1.00
	Restricted permeability	0.46	Seepage	0.53
17A: Keomah-----	Very limited Restricted permeability	1.00	Very limited Depth to saturated zone	1.00
	Depth to saturated zone	1.00	Seepage	0.53
36B: Tama-----	Somewhat limited Restricted permeability	0.46	Somewhat limited Seepage	0.53
			Slope	0.19
36C2: Tama-----	Somewhat limited Restricted permeability	0.46	Very limited Slope	1.00
			Seepage	0.53
43A: Ipava-----	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
	Restricted permeability	1.00		

Table 22.--Sanitary Facilities (Part 1)--Continued

Map symbol and soil name	Septic tank absorption fields		Sewage lagoons	
	Rating class and limiting features	Value	Rating class and limiting features	Value
45A: Denny-----	Very limited Restricted permeability Ponding Depth to saturated zone	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00
46A: Herrick-----	Very limited Depth to saturated zone Restricted permeability	1.00 1.00	Very limited Depth to saturated zone	1.00
50A: Virden-----	Very limited Ponding Depth to saturated zone Restricted permeability	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00
68A: Sable-----	Very limited Ponding Depth to saturated zone Restricted permeability	1.00 1.00 0.46	Very limited Ponding Depth to saturated zone Seepage	1.00 1.00 0.53
86B: Osco-----	Somewhat limited Restricted permeability Depth to saturated zone	0.46 0.40	Somewhat limited Seepage Slope	0.53 0.19
86C2: Osco-----	Somewhat limited Restricted permeability Depth to saturated zone	0.46 0.40	Very limited Slope Seepage	1.00 0.53
112A: Cowden-----	Very limited Restricted permeability Ponding Depth to saturated zone	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00
119C2: Elco-----	Very limited Depth to saturated zone Restricted permeability	1.00 1.00	Very limited Slope Depth to saturated zone Seepage	1.00 0.96 0.53

Table 22.--Sanitary Facilities (Part 1)--Continued

Map symbol and soil name	Septic tank absorption fields		Sewage lagoons	
	Rating class and limiting features	Value	Rating class and limiting features	Value
119D: Elco-----	Very limited		Very limited	
	Depth to saturated zone	1.00	Slope	1.00
	Restricted permeability	1.00	Depth to saturated zone	0.96
	Slope	0.96	Seepage	0.53
119D2: Elco-----	Very limited		Very limited	
	Depth to saturated zone	1.00	Slope	1.00
	Restricted permeability	1.00	Depth to saturated zone	0.96
	Slope	0.96	Seepage	0.53
119D3: Elco-----	Very limited		Very limited	
	Depth to saturated zone	1.00	Slope	1.00
	Restricted permeability	1.00	Depth to saturated zone	0.96
	Slope	0.96	Seepage	0.53
127B: Harrison-----	Very limited		Somewhat limited	
	Restricted permeability	1.00	Depth to saturated zone	0.96
	Depth to saturated zone	1.00	Seepage	0.53
			Slope	0.19
127C2: Harrison-----	Very limited		Very limited	
	Restricted permeability	1.00	Slope	1.00
	Depth to saturated zone	1.00	Depth to saturated zone	0.96
			Seepage	0.53
131C2: Alvin-----	Not limited		Very limited	
			Seepage	1.00
			Slope	1.00
131D2: Alvin-----	Somewhat limited		Very limited	
	Slope	0.96	Slope	1.00
			Seepage	1.00
134A: Camden-----	Somewhat limited		Very limited	
	Restricted permeability	0.46	Seepage	1.00
134B: Camden-----	Somewhat limited		Very limited	
	Restricted permeability	0.46	Seepage	1.00
			Slope	0.19

Table 22.--Sanitary Facilities (Part 1)--Continued

Map symbol and soil name	Septic tank absorption fields		Sewage lagoons	
	Rating class and limiting features	Value	Rating class and limiting features	Value
134C2: Camden-----	Somewhat limited Restricted permeability	0.46	Very limited Slope Seepage	1.00 1.00
138A: Shiloh-----	Very limited Ponding Depth to saturated zone Restricted permeability	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00
198A: Elburn-----	Very limited Depth to saturated zone Restricted permeability	1.00 0.46	Very limited Seepage Depth to saturated zone	1.00 1.00
199A: Plano-----	Somewhat limited Restricted permeability	0.46	Very limited Seepage	1.00
199B: Plano-----	Somewhat limited Restricted permeability	0.46	Very limited Seepage Slope	1.00 0.19
212C2: Thebes-----	Very limited Poor filtering capacity Restricted permeability	1.00 0.46	Very limited Seepage Slope	1.00 1.00
242A: Kendall-----	Very limited Depth to saturated zone Restricted permeability	1.00 0.46	Very limited Depth to saturated zone Seepage	1.00 0.53
244A: Hartsburg-----	Very limited Ponding Depth to saturated zone Restricted permeability	1.00 1.00 0.46	Very limited Ponding Depth to saturated zone Seepage	1.00 1.00 0.53
249A: Edinburg-----	Very limited Restricted permeability Ponding Depth to saturated zone	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Seepage	1.00 1.00 0.28

Table 22.--Sanitary Facilities (Part 1)--Continued

Map symbol and soil name	Septic tank absorption fields		Sewage lagoons	
	Rating class and limiting features	Value	Rating class and limiting features	Value
257A: Clarksdale-----	Very limited Depth to saturated zone Restricted permeability	1.00 1.00	Very limited Depth to saturated zone	1.00
259C2: Assumption-----	Very limited Depth to saturated zone Restricted permeability	1.00 1.00	Very limited Slope Depth to saturated zone Seepage	1.00 1.00 0.53
259D2: Assumption-----	Very limited Depth to saturated zone Restricted permeability Slope	1.00 1.00 0.96	Very limited Slope Depth to saturated zone Seepage	1.00 0.96 0.53
279B: Rozetta-----	Somewhat limited Restricted permeability Depth to saturated zone	0.46 0.40	Somewhat limited Seepage Slope	0.53 0.19
280C2: Fayette-----	Somewhat limited Restricted permeability	0.46	Very limited Slope Seepage	1.00 0.53
280D2: Fayette-----	Somewhat limited Slope Restricted permeability	0.96 0.46	Very limited Slope Seepage	1.00 0.53
280D3: Fayette-----	Somewhat limited Slope Restricted permeability	0.96 0.46	Very limited Slope Seepage	1.00 0.53
533: Urban land-----	Not rated		Not rated	
536: Dumps-----	Not rated		Not rated	
549G: Marseilles-----	Very limited Restricted permeability Depth to bedrock Slope	1.00 1.00 1.00	Very limited Depth to soft bedrock Slope	1.00 1.00

Table 22.--Sanitary Facilities (Part 1)--Continued

Map symbol and soil name	Septic tank absorption fields		Sewage lagoons	
	Rating class and limiting features	Value	Rating class and limiting features	Value
567C2: Elkhart-----	Somewhat limited		Very limited	
	Restricted permeability	0.46	Slope	1.00
	Depth to saturated zone	0.43	Seepage	0.53
567D2: Elkhart-----	Somewhat limited		Very limited	
	Slope	0.96	Slope	1.00
	Restricted permeability	0.46	Seepage	0.53
	Depth to saturated zone	0.43		
630C2: Navlys-----	Somewhat limited		Very limited	
	Restricted permeability	0.46	Slope	1.00
	Depth to saturated zone	0.40	Seepage	0.53
630D2: Navlys-----	Somewhat limited		Very limited	
	Slope	0.96	Slope	1.00
	Restricted permeability	0.46	Seepage	0.53
	Depth to saturated zone	0.40		
630D3: Navlys-----	Somewhat limited		Very limited	
	Slope	0.96	Slope	1.00
	Restricted permeability	0.46	Seepage	0.53
	Depth to saturated zone	0.40		
679B: Blackberry-----	Very limited		Very limited	
	Depth to saturated zone	1.00	Depth to saturated zone	1.00
	Restricted permeability	0.46	Seepage	0.53
			Slope	0.19
684B: Broadwell-----	Very limited		Very limited	
	Poor filtering capacity	1.00	Seepage	1.00
	Restricted permeability	0.46	Slope	0.19
684C2: Broadwell-----	Very limited		Very limited	
	Poor filtering capacity	1.00	Seepage	1.00
	Restricted permeability	0.46	Slope	1.00

Table 22.--Sanitary Facilities (Part 1)--Continued

Map symbol and soil name	Septic tank absorption fields		Sewage lagoons	
	Rating class and limiting features	Value	Rating class and limiting features	Value
685B: Middletown-----	Very limited Poor filtering capacity Restricted permeability	1.00 0.46	Very limited Seepage Slope	1.00 0.19
685C2: Middletown-----	Very limited Poor filtering capacity Restricted permeability	1.00 0.46	Very limited Seepage Slope	1.00 1.00
705A: Buckhart-----	Very limited Depth to saturated zone Restricted permeability	1.00 0.46	Very limited Depth to saturated zone Seepage	1.00 0.53
705B: Buckhart-----	Very limited Depth to saturated zone Restricted permeability	1.00 0.46	Very limited Depth to saturated zone Seepage Slope	1.00 0.53 0.19
712A: Spaulding-----	Very limited Ponding Depth to saturated zone Restricted permeability	1.00 1.00 0.46	Very limited Ponding Depth to saturated zone Seepage	1.00 1.00 0.53
801C: Orthents-----	Somewhat limited Restricted permeability	0.72	Very limited Slope	1.00
830: Landfills-----	Not rated		Not rated	
862: Pits, sand-----	Not rated		Not rated	
864: Pits, quarries-----	Not rated		Not rated	
3073A: Ross-----	Very limited Flooding Restricted permeability Depth to saturated zone	1.00 0.46 0.40	Very limited Flooding Seepage	1.00 1.00

Table 22.--Sanitary Facilities (Part 1)--Continued

Map symbol and soil name	Septic tank absorption fields		Sewage lagoons	
	Rating class and limiting features	Value	Rating class and limiting features	Value
3074A: Radford-----	Very limited Flooding Depth to saturated zone Restricted permeability	 1.00 1.00 0.46	Very limited Flooding Depth to saturated zone Seepage	 1.00 1.00 0.53
3077A: Huntsville-----	Very limited Flooding Restricted permeability Depth to saturated zone	 1.00 0.46 0.40	Very limited Flooding Seepage	 1.00 0.53
3107A: Sawmill-----	Very limited Flooding Ponding Depth to saturated zone Restricted permeability	 1.00 1.00 1.00 0.46	Very limited Ponding Flooding Depth to saturated zone Seepage	 1.00 1.00 1.00 0.53
3284A: Tice-----	Very limited Flooding Depth to saturated zone Restricted permeability	 1.00 1.00 0.46	Very limited Flooding Depth to saturated zone Seepage	 1.00 1.00 0.53
3405A: Zook-----	Very limited Flooding Restricted permeability Depth to saturated zone	 1.00 1.00 1.00	Very limited Flooding Depth to saturated zone	 1.00 1.00
3451A: Lawson-----	Very limited Flooding Depth to saturated zone Restricted permeability	 1.00 1.00 0.46	Very limited Flooding Depth to saturated zone Seepage	 1.00 1.00 0.53
7037A: Worthen-----	Somewhat limited Restricted permeability Flooding	 0.46 0.40	Somewhat limited Seepage Flooding	 0.53 0.40

Table 22.--Sanitary Facilities (Part 1)--Continued

Map symbol and soil name	Septic tank absorption fields		Sewage lagoons	
	Rating class and limiting features	Value	Rating class and limiting features	Value
7075B: Drury-----	Somewhat limited		Somewhat limited	
	Restricted permeability	0.46	Seepage	0.53
	Flooding	0.40	Flooding	0.40
			Slope	0.19
7148A: Proctor-----	Somewhat limited		Very limited	
	Restricted permeability	0.46	Seepage	1.00
	Flooding	0.40	Flooding	0.40
7242A: Kendall-----	Very limited		Very limited	
	Depth to saturated zone	1.00	Depth to saturated zone	1.00
	Restricted permeability	0.46	Seepage	0.53
	Flooding	0.40	Flooding	0.40
8396A: Vesser-----	Very limited		Very limited	
	Flooding	1.00	Flooding	1.00
	Depth to saturated zone	1.00	Depth to saturated zone	1.00
	Restricted permeability	0.46	Seepage	0.53

Table 23.--Sanitary Facilities (Part 2)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
8D: Hickory-----	Somewhat limited Slope Too clayey	0.96 0.50	Somewhat limited Slope	0.96	Somewhat limited Slope Too clayey	0.96 0.50
8D2: Hickory-----	Somewhat limited Slope Too clayey	0.96 0.50	Somewhat limited Slope	0.96	Somewhat limited Slope Too clayey	0.96 0.50
8D3: Hickory-----	Somewhat limited Slope Too clayey	0.96 0.50	Somewhat limited Slope	0.96	Somewhat limited Slope Too clayey	0.96 0.50
8F: Hickory-----	Very limited Slope Too clayey	1.00 0.50	Very limited Slope	1.00	Very limited Slope Too clayey	1.00 0.50
17A: Keomah-----	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Too clayey	1.00 0.50
36B: Tama-----	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
36C2: Tama-----	Not limited		Not limited		Not limited	
43A: Ipava-----	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00
45A: Denny-----	Very limited Depth to saturated zone Ponding Too clayey	1.00 1.00 0.50	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Ponding Depth to saturated zone Too clayey	1.00 1.00 0.50
46A: Herrick-----	Very limited Depth to saturated zone Too clayey	1.00 0.50	Very limited Depth to saturated zone	1.00	Very limited Hard to compact Depth to saturated zone Too clayey	1.00 1.00 0.50

Table 23.--Sanitary Facilities (Part 2)--Continued

Map symbol and soil name	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
50A:						
Virден-----	Very limited		Very limited		Very limited	
	Depth to	1.00	Ponding	1.00	Ponding	1.00
	saturated zone		Depth to	1.00	Depth to	1.00
	Ponding	1.00	saturated zone		saturated zone	
	Too clayey	0.50			Hard to compact	1.00
					Too clayey	0.50
68A:						
Sable-----	Very limited		Very limited		Very limited	
	Depth to	1.00	Ponding	1.00	Ponding	1.00
	saturated zone		Depth to	1.00	Depth to	1.00
	Ponding	1.00	saturated zone		saturated zone	
	Too clayey	0.50			Hard to compact	1.00
					Too clayey	0.50
86B:						
Oscо-----	Very limited		Very limited		Somewhat limited	
	Depth to	1.00	Depth to	1.00	Too clayey	0.50
	saturated zone		saturated zone			
	Too clayey	0.50				
86C2:						
Oscо-----	Very limited		Very limited		Not limited	
	Depth to	1.00	Depth to	1.00		
	saturated zone		saturated zone			
112A:						
Cowden-----	Very limited		Very limited		Very limited	
	Depth to	1.00	Ponding	1.00	Ponding	1.00
	saturated zone		Depth to	1.00	Depth to	1.00
	Ponding	1.00	saturated zone		saturated zone	
	Too clayey	0.50			Hard to compact	1.00
					Too clayey	0.50
119C2:						
Elco-----	Somewhat limited		Somewhat limited		Somewhat limited	
	Depth to	0.68	Depth to	0.04	Too clayey	0.50
	saturated zone		saturated zone		Depth to	0.25
	Too clayey	0.50			saturated zone	
119D:						
Elco-----	Somewhat limited		Somewhat limited		Somewhat limited	
	Slope	0.96	Slope	0.96	Slope	0.96
	Depth to	0.68	Depth to	0.04	Too clayey	0.50
	saturated zone		saturated zone		Depth to	0.25
	Too clayey	0.50			saturated zone	
119D2:						
Elco-----	Somewhat limited		Somewhat limited		Somewhat limited	
	Slope	0.96	Slope	0.96	Slope	0.96
	Depth to	0.68	Depth to	0.04	Too clayey	0.50
	saturated zone		saturated zone		Depth to	0.25
	Too clayey	0.50			saturated zone	
119D3:						
Elco-----	Somewhat limited		Somewhat limited		Somewhat limited	
	Slope	0.96	Slope	0.96	Slope	0.96
	Depth to	0.68	Depth to	0.04	Too clayey	0.50
	saturated zone		saturated zone		Depth to	0.25
	Too clayey	0.50			saturated zone	

Table 23.--Sanitary Facilities (Part 2)--Continued

Map symbol and soil name	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
127B: Harrison-----	Somewhat limited Depth to saturated zone Too clayey	0.68 0.50	Somewhat limited Depth to saturated zone	0.04	Somewhat limited Too clayey Depth to saturated zone	0.50 0.25
127C2: Harrison-----	Somewhat limited Depth to saturated zone Too clayey	0.68 0.50	Somewhat limited Depth to saturated zone	0.04	Somewhat limited Too clayey Depth to saturated zone	0.50 0.25
131C2: Alvin-----	Very limited Seepage	1.00	Very limited Seepage	1.00	Somewhat limited Seepage	0.52
131D2: Alvin-----	Very limited Seepage Slope	1.00 0.96	Very limited Seepage Slope	1.00 0.96	Somewhat limited Slope Seepage	0.96 0.52
134A: Camden-----	Very limited Seepage Too clayey	1.00 0.50	Not limited		Somewhat limited Too clayey	0.50
134B: Camden-----	Very limited Seepage	1.00	Not limited		Somewhat limited Seepage	0.22
134C2: Camden-----	Very limited Seepage Too clayey	1.00 0.50	Not limited		Somewhat limited Too clayey Seepage	0.50 0.22
138A: Shiloh-----	Very limited Depth to saturated zone Ponding Too clayey	1.00 1.00 0.50	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Ponding Depth to saturated zone Too clayey	1.00 1.00 0.50
198A: Elburn-----	Very limited Depth to saturated zone Seepage Too clayey	1.00 1.00 0.50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Too clayey	1.00 0.50
199A: Plano-----	Very limited Seepage Too clayey	1.00 0.50	Not limited		Somewhat limited Too clayey	0.50
199B: Plano-----	Very limited Seepage Too clayey	1.00 0.50	Not limited		Somewhat limited Too clayey	0.50

Table 23.--Sanitary Facilities (Part 2)--Continued

Map symbol and soil name	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
212C2: Thebes-----	Very limited Seepage Too sandy	1.00 1.00	Not limited		Very limited Seepage Too sandy Too clayey	1.00 0.50 0.50
242A: Kendall-----	Very limited Depth to saturated zone Too clayey	1.00 0.50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Too clayey	1.00 0.50
244A: Hartsburg-----	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00
249A: Edinburg-----	Very limited Depth to saturated zone Ponding Too clayey	1.00 1.00 0.50	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Ponding Depth to saturated zone Hard to compact Too clayey	1.00 1.00 1.00 0.50
257A: Clarksdale-----	Very limited Depth to saturated zone Too clayey	1.00 0.50	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone Hard to compact Too clayey	1.00 1.00 1.00 0.50
259C2: Assumption-----	Somewhat limited Depth to saturated zone Too clayey	0.68 0.50	Somewhat limited Depth to saturated zone	0.04	Somewhat limited Too clayey Depth to saturated zone	0.50 0.25
259D2: Assumption-----	Somewhat limited Slope Depth to saturated zone Too clayey	0.96 0.68 0.50	Somewhat limited Slope Depth to saturated zone	0.96 0.04	Somewhat limited Slope Too clayey Depth to saturated zone	0.96 0.50 0.25
279B: Rozetta-----	Very limited Depth to saturated zone Too clayey	1.00 0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Too clayey	0.50
280C2: Fayette-----	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
280D2: Fayette-----	Somewhat limited Slope Too clayey	0.96 0.50	Somewhat limited Slope	0.96	Somewhat limited Slope Too clayey	0.96 0.50

Table 23.--Sanitary Facilities (Part 2)--Continued

Map symbol and soil name	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
280D3: Fayette-----	Somewhat limited Slope Too clayey	0.96 0.50	Somewhat limited Slope	0.96	Somewhat limited Slope Too clayey	0.96 0.50
533: Urban land-----	Not rated		Not rated		Not rated	
536: Dumps-----	Not rated		Not rated		Not rated	
549G: Marseilles-----	Very limited Slope Depth to bedrock Too clayey	1.00 1.00 0.50	Very limited Slope Depth to bedrock	1.00 1.00	Very limited Depth to bedrock Slope Hard to compact Too clayey	1.00 1.00 1.00 0.50
567C2: Elkhart-----	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Not limited	
567D2: Elkhart-----	Very limited Depth to saturated zone Slope	1.00 0.96	Very limited Depth to saturated zone Slope	1.00 0.96	Somewhat limited Slope	0.96
630C2: Navlys-----	Very limited Depth to saturated zone	1.00	Very limited Depth to saturated zone	1.00	Not limited	
630D2: Navlys-----	Very limited Depth to saturated zone Slope	1.00 0.96	Very limited Depth to saturated zone Slope	1.00 0.96	Somewhat limited Slope	0.96
630D3: Navlys-----	Very limited Depth to saturated zone Slope	1.00 0.96	Very limited Depth to saturated zone Slope	1.00 0.96	Somewhat limited Slope	0.96
679B: Blackberry-----	Very limited Depth to saturated zone Too clayey	1.00 0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Too clayey Depth to saturated zone	0.50 0.25
684B: Broadwell-----	Very limited Seepage Too clayey	1.00 0.50	Not limited		Somewhat limited Too clayey	0.50
684C2: Broadwell-----	Very limited Seepage Too clayey	1.00 0.50	Not limited		Somewhat limited Too clayey	0.50

Table 23.--Sanitary Facilities (Part 2)--Continued

Map symbol and soil name	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
685B: Middletown-----	Very limited Seepage Too clayey	1.00 0.50	Not limited		Somewhat limited Too clayey	0.50
685C2: Middletown-----	Very limited Seepage Too clayey	1.00 0.50	Not limited		Somewhat limited Too clayey	0.50
705A: Buckhart-----	Very limited Depth to saturated zone Too clayey	1.00 0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Too clayey Depth to saturated zone	0.50 0.25
705B: Buckhart-----	Very limited Depth to saturated zone Too clayey	1.00 0.50	Very limited Depth to saturated zone	1.00	Somewhat limited Too clayey Depth to saturated zone	0.50 0.25
712A: Spaulding-----	Very limited Depth to saturated zone Ponding	1.00 1.00	Very limited Ponding Depth to saturated zone	1.00 1.00	Very limited Ponding Depth to saturated zone Too clayey	1.00 1.00 0.50
801C: Orthents-----	Somewhat limited Too clayey	0.50	Not limited		Somewhat limited Too clayey	0.50
830: Landfills-----	Not rated		Not rated		Not rated	
862: Pits, sand-----	Not rated		Not rated		Not rated	
864: Pits, quarries-----	Not rated		Not rated		Not rated	
3073A: Ross-----	Very limited Flooding Depth to saturated zone Seepage	1.00 1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Not limited	
3074A: Radford-----	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone	1.00
3077A: Huntsville-----	Very limited Flooding Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone	1.00 1.00	Not limited	

Table 23.--Sanitary Facilities (Part 2)--Continued

Map symbol and soil name	Trench sanitary landfill		Area sanitary landfill		Daily cover for landfill	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
3107A: Sawmill-----	Very limited Flooding Depth to saturated zone Ponding Too clayey	 1.00 1.00 1.00 0.50	Very limited Flooding Ponding Depth to saturated zone	 1.00 1.00 1.00 	Very limited Ponding Depth to saturated zone Too clayey	 1.00 1.00 1.00 0.50
3284A: Tice-----	Very limited Flooding Depth to saturated zone Too clayey	 1.00 1.00 0.50	Very limited Flooding Depth to saturated zone	 1.00 1.00 1.00 	Very limited Hard to compact Depth to saturated zone Too clayey	 1.00 1.00 1.00 0.50
3405A: Zook-----	Very limited Flooding Depth to saturated zone Too clayey	 1.00 1.00 1.00	Very limited Flooding Depth to saturated zone	 1.00 1.00 1.00 	Very limited Depth to saturated zone Too clayey Hard to compact	 1.00 1.00 1.00 1.00
3451A: Lawson-----	Very limited Flooding Depth to saturated zone	 1.00 1.00 	Very limited Flooding Depth to saturated zone	 1.00 1.00 1.00 	Very limited Depth to saturated zone	 1.00
7037A: Worthen-----	Somewhat limited Flooding	 0.40	Somewhat limited Flooding	 0.40	Not limited	
7075B: Drury-----	Somewhat limited Flooding	 0.40	Somewhat limited Flooding	 0.40	Not limited	
7148A: Proctor-----	Very limited Seepage Flooding	 1.00 0.40	Somewhat limited Flooding	 0.40	Not limited	
7242A: Kendall-----	Very limited Depth to saturated zone Too clayey Flooding	 1.00 0.50 0.40	Very limited Depth to saturated zone Flooding	 1.00 0.40	Very limited Depth to saturated zone Too clayey	 1.00 0.50
8396A: Vesser-----	Very limited Flooding Depth to saturated zone Too clayey	 1.00 1.00 0.50	Very limited Flooding Depth to saturated zone	 1.00 1.00 1.00 	Very limited Depth to saturated zone Too clayey	 1.00 0.50

Table 24.--Construction Materials (Part 1)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

Map symbol and soil name	Potential as source of gravel		Potential as source of sand	
	Rating class	Value	Rating class	Value
8D: Hickory-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
8D2: Hickory-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
8D3: Hickory-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
8F: Hickory-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
17A: Keomah-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
36B: Tama-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
36C2: Tama-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
43A: Ipava-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
45A: Denny-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
46A: Herrick-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00

Table 24.--Construction Materials (Part 1)--Continued

Map symbol and soil name	Potential as source of gravel		Potential as source of sand	
	Rating class	Value	Rating class	Value
50A: Virден-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
68A: Sable-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
86B: Osco-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
86C2: Osco-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
112A: Cowden-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
119C2: Elco-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
119D: Elco-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
119D2: Elco-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
119D3: Elco-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
127B: Harrison-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
127C2: Harrison-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
131C2: Alvin-----	Poor		Fair	
	Bottom layer	0.00	Thickest layer	0.04
	Thickest layer	0.00	Bottom layer	0.15

Table 24.--Construction Materials (Part 1)--Continued

Map symbol and soil name	Potential as source of gravel		Potential as source of sand	
	Rating class	Value	Rating class	Value
131D2:				
Alvin-----	Poor		Fair	
	Bottom layer	0.00	Thickest layer	0.04
	Thickest layer	0.00	Bottom layer	0.22
134A:				
Camden-----	Poor		Fair	
	Bottom layer	0.00	Thickest layer	0.00
	Thickest layer	0.00	Bottom layer	0.10
134B:				
Camden-----	Poor		Fair	
	Bottom layer	0.00	Thickest layer	0.00
	Thickest layer	0.00	Bottom layer	0.02
134C2:				
Camden-----	Poor		Fair	
	Bottom layer	0.00	Thickest layer	0.00
	Thickest layer	0.00	Bottom layer	0.10
138A:				
Shiloh-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
198A:				
Elburn-----	Poor		Fair	
	Bottom layer	0.00	Thickest layer	0.00
	Thickest layer	0.00	Bottom layer	0.10
199A:				
Plano-----	Poor		Fair	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.06
199B:				
Plano-----	Poor		Fair	
	Bottom layer	0.00	Thickest layer	0.00
	Thickest layer	0.00	Bottom layer	0.06
212C2:				
Thebes-----	Poor		Fair	
	Bottom layer	0.00	Thickest layer	0.00
	Thickest layer	0.00	Bottom layer	0.17
242A:				
Kendall-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
244A:				
Hartsburg-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
249A:				
Edinburg-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00

Table 24.--Construction Materials (Part 1)--Continued

Map symbol and soil name	Potential as source of gravel		Potential as source of sand	
	Rating class	Value	Rating class	Value
257A: Clarksdale-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
259C2: Assumption-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
259D2: Assumption-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
279B: Rozetta-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
280C2: Fayette-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
280D2: Fayette-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
280D3: Fayette-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
533: Urban land-----	Not rated		Not rated	
536: Dumps-----	Not rated		Not rated	
549G: Marseilles-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
567C2: Elkhart-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
567D2: Elkhart-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
630C2: Navlys-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00

Table 24.--Construction Materials (Part 1)--Continued

Map symbol and soil name	Potential as source of gravel		Potential as source of sand	
	Rating class	Value	Rating class	Value
630D2: Navlys-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
630D3: Navlys-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
679B: Blackberry-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
684B: Broadwell-----	Poor		Fair	
	Bottom layer	0.00	Thickest layer	0.00
	Thickest layer	0.00	Bottom layer	0.10
684C2: Broadwell-----	Poor		Fair	
	Bottom layer	0.00	Thickest layer	0.00
	Thickest layer	0.00	Bottom layer	0.10
685B: Middletown-----	Poor		Fair	
	Bottom layer	0.00	Thickest layer	0.00
	Thickest layer	0.00	Bottom layer	0.10
685C2: Middletown-----	Poor		Fair	
	Bottom layer	0.00	Thickest layer	0.00
	Thickest layer	0.00	Bottom layer	0.10
705A: Buckhart-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
705B: Buckhart-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
712A: Spaulding-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
801C: Orthents-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
830: Landfills-----	Not rated		Not rated	
862: Pits, sand-----	Not rated		Not rated	

Table 24.--Construction Materials (Part 1)--Continued

Map symbol and soil name	Potential as source of gravel		Potential as source of sand	
	Rating class	Value	Rating class	Value
864: Pits, quarries-----	Not rated		Not rated	
3073A: Ross-----	Poor		Fair	
	Bottom layer	0.00	Thickest layer	0.00
	Thickest layer	0.00	Bottom layer	0.03
3074A: Radford-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
3077A: Huntsville-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
3107A: Sawmill-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
3284A: Tice-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
3405A: Zook-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
3451A: Lawson-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
7037A: Worthen-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
7075B: Drury-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00
7148A: Proctor-----	Poor		Fair	
	Bottom layer	0.00	Thickest layer	0.00
	Thickest layer	0.00	Bottom layer	0.10
7242A: Kendall-----	Poor		Fair	
	Bottom layer	0.00	Thickest layer	0.00
	Thickest layer	0.00	Bottom layer	0.10
8396A: Vesser-----	Poor		Poor	
	Bottom layer	0.00	Bottom layer	0.00
	Thickest layer	0.00	Thickest layer	0.00

Table 25.--Construction Materials (Part 2)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Potential as source of topsoil		Potential as source of roadfill	
	Rating class and limiting features	Value	Rating class and limiting features	Value
8D: Hickory-----	Fair		Poor	
	Slope	0.04	Low strength	0.00
	Too clayey	0.57	Shrink-swell	0.98
	Rock fragments	0.97		
8D2: Hickory-----	Fair		Poor	
	Slope	0.04	Low strength	0.00
	Too clayey	0.57	Shrink-swell	0.97
	Rock fragments	0.97		
8D3: Hickory-----	Fair		Poor	
	Slope	0.04	Low strength	0.00
	Too clayey	0.57	Shrink-swell	0.97
	Rock fragments	0.97		
8F: Hickory-----	Poor		Poor	
	Slope	0.00	Slope	0.00
	Too clayey	0.57	Low strength	0.00
	Rock fragments	0.88	Shrink-swell	0.98
17A: Keomah-----	Fair		Poor	
	Depth to saturated zone	0.04	Low strength	0.00
	Too clayey	0.05	Depth to saturated zone	0.04
	Too acid	0.98	Shrink-swell	0.89
36B: Tama-----	Fair		Poor	
	Too clayey	0.64	Low strength	0.00
			Shrink-swell	0.87
36C2: Tama-----	Fair		Poor	
	Too clayey	0.64	Low strength	0.00
			Shrink-swell	0.87
43A: Ipava-----	Fair		Poor	
	Depth to saturated zone	0.14	Low strength	0.00
			Depth to saturated zone	0.14
			Shrink-swell	0.67
45A: Denny-----	Poor		Poor	
	Depth to saturated zone	0.00	Depth to saturated zone	0.00
	Too clayey	0.01	Low strength	0.00
			Shrink-swell	0.74

Table 25.--Construction Materials (Part 2)--Continued

Map symbol and soil name	Potential as source of topsoil		Potential as source of roadfill	
	Rating class and limiting features	Value	Rating class and limiting features	Value
46A: Herrick-----	Fair		Poor	
	Too clayey	0.05	Low strength	0.00
	Depth to saturated zone	0.14	Depth to saturated zone	0.14
	Too acid	0.98	Shrink-swell	0.47
50A: Virden-----	Poor		Poor	
	Depth to saturated zone	0.00	Depth to saturated zone	0.00
	Too clayey	0.02	Low strength	0.00
			Shrink-swell	0.35
68A: Sable-----	Poor		Poor	
	Depth to saturated zone	0.00	Depth to saturated zone	0.00
	Too clayey	0.98	Low strength	0.00
			Shrink-swell	0.87
86B: Osc-----	Fair		Poor	
	Too clayey	0.64	Low strength	0.00
			Shrink-swell	0.87
86C2: Osc-----	Fair		Poor	
	Too clayey	0.64	Low strength	0.00
			Shrink-swell	0.87
112A: Cowden-----	Poor		Poor	
	Depth to saturated zone	0.00	Depth to saturated zone	0.00
	Too clayey	0.05	Low strength	0.00
			Shrink-swell	0.55
119C2: Elco-----	Fair		Poor	
	Too clayey	0.57	Low strength	0.00
	Depth to saturated zone	0.98	Shrink-swell	0.43
			Depth to saturated zone	0.98
119D: Elco-----	Fair		Poor	
	Slope	0.04	Low strength	0.00
	Too clayey	0.57	Shrink-swell	0.38
	Depth to saturated zone	0.98	Depth to saturated zone	0.98
119D2: Elco-----	Fair		Poor	
	Slope	0.04	Low strength	0.00
	Too clayey	0.57	Shrink-swell	0.38
	Depth to saturated zone	0.98	Depth to saturated zone	0.98

Table 25.--Construction Materials (Part 2)--Continued

Map symbol and soil name	Potential as source of topsoil		Potential as source of roadfill	
	Rating class and limiting features	Value	Rating class and limiting features	Value
119D3: Elco-----	Fair		Poor	
	Slope	0.04	Low strength	0.00
	Too clayey	0.57	Shrink-swell	0.34
	Depth to saturated zone	0.98	Depth to saturated zone	0.98
127B: Harrison-----	Fair		Poor	
	Too clayey	0.67	Low strength	0.00
	Depth to saturated zone	0.98	Shrink-swell	0.87
			Depth to saturated zone	0.98
127C2: Harrison-----	Fair		Poor	
	Too clayey	0.67	Low strength	0.00
	Depth to saturated zone	0.98	Shrink-swell	0.87
			Depth to saturated zone	0.98
131C2: Alvin-----	Good		Good	
131D2: Alvin-----	Fair		Good	
	Slope	0.04		
134A: Camden-----	Fair		Fair	
	Too clayey	0.58	Shrink-swell	0.95
134B: Camden-----	Fair		Good	
	Too clayey	0.55		
134C2: Camden-----	Fair		Good	
	Too clayey	0.58		
138A: Shiloh-----	Poor		Poor	
	Depth to saturated zone	0.00	Depth to saturated zone	0.00
	Too clayey	0.18	Low strength	0.00
			Shrink-swell	0.12
198A: Elburn-----	Fair		Poor	
	Depth to saturated zone	0.14	Low strength	0.00
	Too clayey	0.81	Depth to saturated zone	0.14
			Shrink-swell	0.99
199A: Plano-----	Fair		Poor	
	Too clayey	0.67	Low strength	0.00
			Shrink-swell	0.98

Table 25.--Construction Materials (Part 2)--Continued

Map symbol and soil name	Potential as source of topsoil		Potential as source of roadfill	
	Rating class and limiting features	Value	Rating class and limiting features	Value
199B: Plano-----	Fair Too clayey	0.67	Poor Low strength Shrink-swell	0.00 0.99
212C2: Thebes-----	Fair Too clayey Too acid	0.57 0.98	Good	
242A: Kendall-----	Fair Depth to saturated zone Too clayey Too acid	0.04 0.57 0.99	Poor Low strength Depth to saturated zone Shrink-swell	0.00 0.04 0.95
244A: Hartsburg-----	Poor Depth to saturated zone Too clayey	0.00 0.98	Poor Depth to saturated zone Low strength	0.00 0.00
249A: Edinburg-----	Poor Depth to saturated zone Too clayey	0.00 0.00	Poor Depth to saturated zone Low strength Shrink-swell	0.00 0.00 0.17
257A: Clarksdale-----	Fair Too clayey Depth to saturated zone	0.01 0.04	Poor Low strength Depth to saturated zone Shrink-swell	0.00 0.04 0.50
259C2: Assumption-----	Fair Too clayey Depth to saturated zone	0.64 0.98	Poor Low strength Shrink-swell Depth to saturated zone	0.00 0.31 0.98
259D2: Assumption-----	Fair Slope Too clayey Depth to saturated zone	0.04 0.64 0.98	Poor Low strength Shrink-swell Depth to saturated zone	0.00 0.38 0.98
279B: Rozetta-----	Fair Too clayey	0.57	Poor Low strength Shrink-swell	0.00 0.92
280C2: Fayette-----	Fair Too clayey Too acid	0.57 0.98	Poor Low strength Shrink-swell	0.00 0.87

Table 25.--Construction Materials (Part 2)--Continued

Map symbol and soil name	Potential as source of topsoil		Potential as source of roadfill	
	Rating class and limiting features	Value	Rating class and limiting features	Value
280D2: Fayette-----	Fair		Poor	
	Slope	0.04	Low strength	0.00
	Too clayey	0.57	Shrink-swell	0.87
	Too acid	0.98		
280D3: Fayette-----	Fair		Poor	
	Slope	0.04	Low strength	0.00
	Too clayey	0.57	Shrink-swell	0.87
	Too acid	0.98		
533: Urban land-----	Not rated		Not rated	
536: Dumps-----	Not rated		Not rated	
549G: Marseilles-----	Poor		Poor	
	Slope	0.00	Depth to bedrock	0.00
	Too clayey	0.39	Slope	0.00
	Too acid	0.88	Low strength	0.00
	Depth to bedrock	0.90	Shrink-swell	0.87
567C2: Elkhart-----	Fair		Poor	
	Too clayey	0.64	Low strength	0.00
567D2: Elkhart-----	Fair		Poor	
	Slope	0.04	Low strength	0.00
	Too clayey	0.64		
630C2: Navlys-----	Fair		Poor	
	Too clayey	0.57	Low strength	0.00
630D2: Navlys-----	Fair		Poor	
	Slope	0.04	Low strength	0.00
	Too clayey	0.57		
630D3: Navlys-----	Fair		Poor	
	Slope	0.04	Low strength	0.00
	Too clayey	0.57		
679B: Blackberry-----	Fair		Poor	
	Too clayey	0.63	Low strength	0.00
	Depth to saturated zone	0.98	Shrink-swell	0.93
			Depth to saturated zone	0.98
684B: Broadwell-----	Fair		Poor	
	Too clayey	0.64	Low strength	0.00
			Shrink-swell	0.98

Table 25.--Construction Materials (Part 2)--Continued

Map symbol and soil name	Potential as source of topsoil		Potential as source of roadfill	
	Rating class and limiting features	Value	Rating class and limiting features	Value
684C2: Broadwell-----	Fair Too clayey	0.64	Poor Low strength Shrink-swell	0.00 0.97
685B: Middletown-----	Fair Too clayey	0.57	Fair Shrink-swell	0.99
685C2: Middletown-----	Fair Too clayey	0.57	Poor Low strength Shrink-swell	0.00 0.97
705A: Buckhart-----	Fair Depth to saturated zone	0.98	Poor Low strength Shrink-swell Depth to saturated zone	0.00 0.87 0.98
705B: Buckhart-----	Fair Too clayey Depth to saturated zone	0.67 0.98	Poor Low strength Shrink-swell Depth to saturated zone	0.00 0.87 0.98
712A: Spaulding-----	Poor Depth to saturated zone Carbonate content Too clayey	0.00 0.68 0.98	Poor Depth to saturated zone Low strength Shrink-swell	0.00 0.00 0.98
801C: Orthents-----	Fair Too clayey	0.67	Poor Low strength Shrink-swell	0.00 0.87
830: Landfills-----	Not rated		Not rated	
862: Pits, sand-----	Not rated		Not rated	
864: Pits, quarries-----	Not rated		Not rated	
3073A: Ross-----	Fair Hard to reclaim	0.98	Fair Low strength	0.22
3074A: Radford-----	Fair Depth to saturated zone	0.14	Poor Low strength Depth to saturated zone	0.00 0.14

Table 25.--Construction Materials (Part 2)--Continued

Map symbol and soil name	Potential as source of topsoil		Potential as source of roadfill	
	Rating class and limiting features	Value	Rating class and limiting features	Value
3077A: Huntsville-----	Good		Poor Low strength Shrink-swell	0.00 0.87
3107A: Sawmill-----	Poor Depth to saturated zone Too clayey	0.00 0.98	Poor Depth to saturated zone Low strength Shrink-swell	0.00 0.00 0.00 0.87
3284A: Tice-----	Fair Depth to saturated zone Too clayey	0.14 0.64	Poor Low strength Depth to saturated zone Shrink-swell	0.00 0.14 0.87
3405A: Zook-----	Poor Depth to saturated zone Too clayey	0.00 0.00	Poor Depth to saturated zone Low strength Shrink-swell	0.00 0.00 0.00 0.12
3451A: Lawson-----	Fair Depth to saturated zone	0.14	Poor Low strength Depth to saturated zone	0.00 0.14
7037A: Worthen-----	Good		Poor Low strength	0.00
7075B: Drury-----	Good		Good	
7148A: Proctor-----	Fair Too clayey	0.67	Fair Shrink-swell	0.97
7242A: Kendall-----	Fair Depth to saturated zone Too clayey	0.04 0.57	Poor Low strength Depth to saturated zone Shrink-swell	0.00 0.04 0.95
8396A: Vesser-----	Poor Depth to saturated zone	0.00	Poor Depth to saturated zone Low strength Shrink-swell	0.00 0.00 0.00 0.87

Table 26.--Water Management (Part 1)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
8D: Hickory-----	Somewhat limited Seepage Slope	0.72 0.02	Somewhat limited Piping	0.03	Very limited Deep to water	1.00
8D2: Hickory-----	Somewhat limited Seepage Slope	0.72 0.02	Somewhat limited Piping	0.01	Very limited Deep to water	1.00
8D3: Hickory-----	Somewhat limited Seepage Slope	0.72 0.02	Somewhat limited Piping	0.11	Very limited Deep to water	1.00
8F: Hickory-----	Somewhat limited Seepage Slope	0.72 0.34	Somewhat limited Piping	0.14	Very limited Deep to water	1.00
17A: Keomah-----	Somewhat limited Seepage	0.72	Very limited Depth to saturated zone Piping	1.00 0.23	Somewhat limited Slow refill Cutbanks cave	0.28 0.10
36B: Tama-----	Somewhat limited Seepage	0.72	Somewhat limited Piping	0.03	Very limited Deep to water	1.00
36C2: Tama-----	Somewhat limited Seepage	0.72	Not limited		Very limited Deep to water	1.00
43A: Ipava-----	Somewhat limited Seepage	0.72	Very limited Depth to saturated zone Piping	1.00 0.01	Somewhat limited Slow refill Cutbanks cave	0.28 0.10
45A: Denny-----	Somewhat limited Seepage	0.04	Very limited Ponding Depth to saturated zone Piping	1.00 1.00 0.08	Somewhat limited Slow refill Cutbanks cave	0.28 0.10
46A: Herrick-----	Somewhat limited Seepage	0.04	Very limited Depth to saturated zone	1.00	Somewhat limited Slow refill Cutbanks cave	0.96 0.10

Table 26.--Water Management (Part 1)--Continued

Map symbol and soil name	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
50A: Virden-----	Somewhat limited Seepage	0.04	Very limited Ponding Depth to saturated zone	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.28 0.10
68A: Sable-----	Somewhat limited Seepage	0.72	Very limited Ponding Depth to saturated zone	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.28 0.10
86B: Osco-----	Somewhat limited Seepage	0.72	Somewhat limited Piping	0.01	Very limited Deep to water	1.00
86C2: Osco-----	Somewhat limited Seepage	0.72	Not limited		Very limited Deep to water	1.00
112A: Cowden-----	Somewhat limited Seepage	0.04	Very limited Ponding Depth to saturated zone	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.28 0.10
119C2: Elco-----	Somewhat limited Seepage	0.72	Somewhat limited Depth to saturated zone	0.68	Very limited Deep to water	1.00
119D: Elco-----	Somewhat limited Seepage Slope	0.72 0.02	Somewhat limited Depth to saturated zone Piping	0.68 0.01	Very limited Deep to water	1.00
119D2: Elco-----	Somewhat limited Seepage Slope	0.72 0.02	Somewhat limited Depth to saturated zone	0.68	Very limited Deep to water	1.00
119D3: Elco-----	Somewhat limited Seepage Slope	0.72 0.02	Somewhat limited Depth to saturated zone	0.68	Very limited Deep to water	1.00
127B: Harrison-----	Somewhat limited Seepage	0.72	Somewhat limited Depth to saturated zone	0.68	Very limited Deep to water	1.00
127C2: Harrison-----	Somewhat limited Seepage	0.72	Somewhat limited Depth to saturated zone	0.68	Very limited Deep to water	1.00

Table 26.--Water Management (Part 1)--Continued

Map symbol and soil name	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
131C2: Alvin-----	Very limited Seepage	1.00	Somewhat limited Seepage	0.15	Very limited Deep to water	1.00
131D2: Alvin-----	Very limited Seepage Slope	1.00 0.02	Somewhat limited Seepage	0.22	Very limited Deep to water	1.00
134A: Camden-----	Very limited Seepage	1.00	Somewhat limited Piping Seepage	0.94 0.10	Very limited Deep to water	1.00
134B: Camden-----	Somewhat limited Seepage	0.72	Somewhat limited Piping Seepage	0.78 0.02	Very limited Deep to water	1.00
134C2: Camden-----	Very limited Seepage	1.00	Somewhat limited Piping Seepage	0.93 0.10	Very limited Deep to water	1.00
138A: Shiloh-----	Somewhat limited Seepage	0.04	Very limited Ponding Depth to saturated zone	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.96 0.10
198A: Elburn-----	Very limited Seepage	1.00	Very limited Depth to saturated zone Piping Seepage	1.00 0.37 0.10	Very limited Cutbanks cave	1.00
199A: Plano-----	Very limited Seepage	1.00	Somewhat limited Piping	0.67	Very limited Deep to water	1.00
199B: Plano-----	Very limited Seepage	1.00	Somewhat limited Piping Seepage	0.71 0.10	Very limited Deep to water	1.00
212C2: Thebes-----	Very limited Seepage	1.00	Somewhat limited Seepage	0.17	Very limited Deep to water	1.00
242A: Kendall-----	Somewhat limited Seepage	0.72	Very limited Depth to saturated zone Piping	1.00 0.95	Somewhat limited Slow refill Cutbanks cave	0.28 0.10

Table 26.--Water Management (Part 1)--Continued

Map symbol and soil name	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
244A: Hartsburg-----	Somewhat limited Seepage	0.72	Very limited Ponding Depth to saturated zone Piping	1.00 1.00 0.04	Somewhat limited Slow refill Cutbanks cave	0.28 0.10
249A: Edinburg-----	Somewhat limited Seepage	0.54	Very limited Ponding Depth to saturated zone Hard to pack	1.00 1.00 0.64	Somewhat limited Slow refill Cutbanks cave	0.28 0.10
257A: Clarksdale-----	Somewhat limited Seepage	0.04	Very limited Depth to saturated zone	1.00	Somewhat limited Slow refill Cutbanks cave	0.96 0.10
259C2: Assumption-----	Somewhat limited Seepage	0.72	Somewhat limited Depth to saturated zone	0.68	Very limited Deep to water	1.00
259D2: Assumption-----	Somewhat limited Seepage Slope	0.72 0.02	Somewhat limited Depth to saturated zone	0.68	Very limited Deep to water	1.00
279B: Rozetta-----	Somewhat limited Seepage	0.72	Not limited		Very limited Deep to water	1.00
280C2: Fayette-----	Somewhat limited Seepage	0.72	Somewhat limited Piping	0.03	Very limited Deep to water	1.00
280D2: Fayette-----	Somewhat limited Seepage Slope	0.72 0.02	Somewhat limited Piping	0.03	Very limited Deep to water	1.00
280D3: Fayette-----	Somewhat limited Seepage Slope	0.72 0.02	Somewhat limited Piping	0.12	Very limited Deep to water	1.00
533: Urban land-----	Not rated		Not rated		Not rated	
536: Dumps-----	Not rated		Not rated		Not rated	
549G: Marseilles-----	Somewhat limited Slope Depth to bedrock	0.99 0.04	Somewhat limited Thin layer Hard to pack	0.70 0.03	Very limited Deep to water	1.00

Table 26.--Water Management (Part 1)--Continued

Map symbol and soil name	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
567C2: Elkhart-----	Somewhat limited Seepage	0.72	Somewhat limited Piping	0.03	Very limited Deep to water	1.00
567D2: Elkhart-----	Somewhat limited Seepage Slope	0.72 0.02	Somewhat limited Piping	0.03	Very limited Deep to water	1.00
630C2: Navlys-----	Somewhat limited Seepage	0.72	Somewhat limited Piping	0.20	Very limited Deep to water	1.00
630D2: Navlys-----	Somewhat limited Seepage Slope	0.72 0.02	Somewhat limited Piping	0.20	Very limited Deep to water	1.00
630D3: Navlys-----	Somewhat limited Seepage Slope	0.72 0.02	Somewhat limited Piping	0.10	Very limited Deep to water	1.00
679B: Blackberry-----	Somewhat limited Seepage	0.72	Somewhat limited Piping Depth to saturated zone	0.86 0.68	Somewhat limited Slow refill Deep to water Cutbanks cave	0.28 0.14 0.10
684B: Broadwell-----	Very limited Seepage	1.00	Somewhat limited Piping Seepage	0.98 0.50	Very limited Deep to water	1.00
684C2: Broadwell-----	Very limited Seepage	1.00	Somewhat limited Piping Seepage	0.32 0.10	Very limited Deep to water	1.00
685B: Middletown-----	Very limited Seepage	1.00	Somewhat limited Piping Seepage	0.77 0.10	Very limited Deep to water	1.00
685C2: Middletown-----	Very limited Seepage	1.00	Somewhat limited Seepage Piping	0.10 0.05	Very limited Deep to water	1.00
705A: Buckhart-----	Somewhat limited Seepage	0.72	Somewhat limited Depth to saturated zone Piping	0.68 0.03	Somewhat limited Slow refill Deep to water Cutbanks cave	0.28 0.14 0.10

Table 26.--Water Management (Part 1)--Continued

Map symbol and soil name	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
705B: Buckhart-----	Somewhat limited Seepage	0.72	Somewhat limited Depth to saturated zone Piping	0.68 0.03	Somewhat limited Slow refill Deep to water Cutbanks cave	0.28 0.14 0.10
712A: Spaulding-----	Somewhat limited Seepage	0.72	Very limited Ponding Depth to saturated zone	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.28 0.10
801C: Orthents-----	Somewhat limited Seepage	0.54	Somewhat limited Piping	0.50	Very limited Deep to water	1.00
830: Landfills-----	Not rated		Not rated		Not rated	
862: Pits, sand-----	Not rated		Not rated		Not rated	
864: Pits, quarries-----	Not rated		Not rated		Not rated	
3073A: Ross-----	Very limited Seepage	1.00	Very limited Piping Seepage	1.00 0.03	Very limited Deep to water	1.00
3074A: Radford-----	Somewhat limited Seepage	0.72	Very limited Depth to saturated zone Piping	1.00 0.22	Somewhat limited Slow refill Cutbanks cave	0.28 0.10
3077A: Huntsville-----	Somewhat limited Seepage	0.72	Somewhat limited Piping	0.50	Very limited Deep to water	1.00
3107A: Sawmill-----	Somewhat limited Seepage	0.72	Very limited Ponding Depth to saturated zone	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.28 0.10
3284A: Tice-----	Somewhat limited Seepage	0.72	Very limited Depth to saturated zone Piping	1.00 0.02	Somewhat limited Slow refill Cutbanks cave	0.28 0.10
3405A: Zook-----	Somewhat limited Seepage	0.02	Very limited Depth to saturated zone Hard to pack	1.00 1.00	Somewhat limited Slow refill Cutbanks cave	0.96 0.10

Table 26.--Water Management (Part 1)--Continued

Map symbol and soil name	Pond reservoir areas		Embankments, dikes, and levees		Aquifer-fed excavated ponds	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
3451A: Lawson-----	Somewhat limited Seepage	0.72	Very limited Depth to saturated zone Piping	1.00 0.60	Somewhat limited Slow refill Cutbanks cave	0.28 0.10
7037A: Worthen-----	Somewhat limited Seepage	0.72	Somewhat limited Piping	0.68	Very limited Deep to water	1.00
7075B: Drury-----	Somewhat limited Seepage	0.72	Somewhat limited Piping	0.99	Very limited Deep to water	1.00
7148A: Proctor-----	Very limited Seepage	1.00	Somewhat limited Piping Seepage	0.59 0.10	Very limited Deep to water	1.00
7242A: Kendall-----	Somewhat limited Seepage	0.72	Very limited Depth to saturated zone Piping Seepage	1.00 0.74 0.10	Somewhat limited Slow refill Cutbanks cave	0.28 0.10
8396A: Vesser-----	Somewhat limited Seepage	0.72	Very limited Depth to saturated zone Piping	1.00 0.03	Somewhat limited Slow refill Cutbanks cave	0.28 0.10

Table 27.--Water Management (Part 2)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Grassed waterways and surface drains		Terraces and diversions		Drainage	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
8D: Hickory-----	Very limited Slope	1.00	Very limited Water erosion	1.00	Somewhat limited Slope	0.96
	Water erosion	1.00	Slope	1.00	Cutbanks cave	0.10
8D2: Hickory-----	Very limited Slope	1.00	Very limited Slope	1.00	Somewhat limited Slope	0.96
	Water erosion	0.89	Water erosion	0.89	Cutbanks cave	0.10
8D3: Hickory-----	Very limited Slope	1.00	Very limited Slope	1.00	Somewhat limited Slope	0.96
	Water erosion	0.56	Water erosion	0.56	Cutbanks cave	0.10
8F: Hickory-----	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Slope	1.00
	Water erosion	0.89	Water erosion	0.89	Cutbanks cave	0.10
17A: Keomah-----	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Very limited Depth to	1.00
	Depth to saturated zone	1.00	Depth to saturated zone	1.00	Frost action	1.00
	Restricted permeability	0.96	Restricted permeability	0.96	Cutbanks cave	0.10
36B: Tama-----	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Very limited Frost action	1.00
	Slope	0.25	Slope	0.25	Cutbanks cave	0.10
36C2: Tama-----	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Very limited Frost action	1.00
	Slope	0.99	Slope	0.99	Cutbanks cave	0.10
43A: Ipava-----	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Very limited Depth to	1.00
	Depth to saturated zone	1.00	Depth to saturated zone	1.00	Frost action	1.00
	Restricted permeability	0.21	Restricted permeability	0.21	Cutbanks cave	0.10
45A: Denny-----	Very limited Depth to	1.00	Very limited Water erosion	1.00	Very limited Ponding	1.00
	saturated zone	1.00	Ponding	1.00	Depth to	1.00
	Water erosion	1.00	Depth to	1.00	saturated zone	1.00
	Restricted permeability	0.96	saturated zone	0.96	Frost action	1.00
			Restricted permeability	0.96	Cutbanks cave	0.10

Table 27.--Water Management (Part 2)--Continued

Map symbol and soil name	Grassed waterways and surface drains		Terraces and diversions		Drainage	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
46A: Herrick-----	Very limited Water erosion Depth to saturated zone Restricted permeability	 1.00 1.00 0.21	Very limited Water erosion Depth to saturated zone Restricted permeability	 1.00 1.00 0.21	Very limited Depth to saturated zone Frost action Cutbanks cave	 1.00 1.00 0.10
50A: Viriden-----	Very limited Depth to saturated zone Water erosion Restricted permeability	 1.00 1.00 0.21	Very limited Water erosion Ponding Depth to saturated zone Restricted permeability	 1.00 1.00 1.00 0.21	Very limited Ponding Depth to saturated zone Frost action Cutbanks cave	 1.00 1.00 1.00 0.10
68A: Sable-----	Very limited Depth to saturated zone Water erosion	 1.00 1.00	Very limited Water erosion Ponding Depth to saturated zone	 1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Frost action Cutbanks cave	 1.00 1.00 1.00 0.10
86B: Osco-----	Very limited Water erosion Slope	 1.00 0.25	Very limited Water erosion Slope	 1.00 0.25	Very limited Frost action Depth to saturated zone Cutbanks cave	 1.00 0.15 0.10
86C2: Osco-----	Very limited Water erosion Slope	 1.00 0.99	Very limited Water erosion Slope	 1.00 0.99	Very limited Frost action Depth to saturated zone Cutbanks cave	 1.00 0.15 0.10
112A: Cowden-----	Very limited Depth to saturated zone Water erosion Restricted permeability	 1.00 1.00 0.96	Very limited Water erosion Ponding Depth to saturated zone Restricted permeability	 1.00 1.00 1.00 0.96	Very limited Ponding Depth to saturated zone Frost action Cutbanks cave	 1.00 1.00 1.00 0.10
119C2: Elco-----	Very limited Water erosion Slope Restricted permeability Depth to saturated zone	 1.00 0.99 0.43 0.25	Very limited Water erosion Depth to saturated zone Slope Restricted permeability	 1.00 1.00 0.99 0.43	Very limited Frost action Depth to saturated zone Cutbanks cave	 1.00 0.99 0.10

Table 27.--Water Management (Part 2)--Continued

Map symbol and soil name	Grassed waterways and surface drains		Terraces and diversions		Drainage	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
119D: Elco-----	Very limited Slope	1.00	Very limited Water erosion	1.00	Very limited Frost action	1.00
	Water erosion	1.00	Slope	1.00	Depth to	0.99
	Restricted	0.43	Depth to	1.00	saturated zone	
	permeability		saturated zone		Slope	0.96
	Depth to	0.25	Restricted	0.43	Cutbanks cave	0.10
	saturated zone		permeability			
119D2: Elco-----	Very limited Slope	1.00	Very limited Water erosion	1.00	Very limited Frost action	1.00
	Water erosion	1.00	Slope	1.00	Depth to	0.99
	Restricted	0.43	Depth to	1.00	saturated zone	
	permeability		saturated zone		Slope	0.96
	Depth to	0.25	Restricted	0.43	Cutbanks cave	0.10
	saturated zone		permeability	0.43		
119D3: Elco-----	Very limited Slope	1.00	Very limited Water erosion	1.00	Very limited Frost action	1.00
	Water erosion	1.00	Slope	1.00	Depth to	0.99
	Restricted	0.43	Depth to	1.00	saturated zone	
	permeability		saturated zone		Slope	0.96
	Depth to	0.25	Restricted	0.43	Cutbanks cave	0.10
	saturated zone		permeability			
127B: Harrison-----	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Very limited Frost action	1.00
	Slope	0.25	Depth to	1.00	Depth to	0.99
	Depth to	0.25	saturated zone		saturated zone	
	saturated zone		Slope	0.25	Cutbanks cave	0.10
127C2: Harrison-----	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Very limited Frost action	1.00
	Slope	0.99	Depth to	1.00	Depth to	0.99
	Depth to	0.25	saturated zone		saturated zone	
	saturated zone		Slope	0.99	Cutbanks cave	0.10
131C2: Alvin-----	Somewhat limited Slope	0.99	Somewhat limited Slope	0.99	Very limited Cutbanks cave	1.00
	Water erosion	0.17	Water erosion	0.17		
131D2: Alvin-----	Very limited Slope	1.00	Very limited Slope	1.00	Very limited Cutbanks cave	1.00
	Water erosion	0.17	Water erosion	0.17	Slope	0.96
134A: Camden-----	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Very limited Cutbanks cave	1.00
					Frost action	1.00
134B: Camden-----	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Very limited Frost action	1.00
	Slope	0.25	Slope	0.25	Cutbanks cave	0.10

Table 27.--Water Management (Part 2)--Continued

Map symbol and soil name	Grassed waterways and surface drains		Terraces and diversions		Drainage	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
134C2: Camden-----	Very limited Water erosion Slope	1.00 0.99	Very limited Water erosion Slope	1.00 0.99	Very limited Cutbanks cave Frost action	1.00 1.00
138A: Shiloh-----	Very limited Depth to saturated zone Restricted permeability Water erosion	1.00 0.21 0.17	Very limited Ponding Depth to saturated zone Restricted permeability Water erosion	1.00 1.00 0.21 0.17	Very limited Ponding Depth to saturated zone Frost action Cutbanks cave	1.00 1.00 1.00 0.10
198A: Elburn-----	Very limited Water erosion Depth to saturated zone	1.00 1.00	Very limited Water erosion Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Cutbanks cave Frost action	1.00 1.00 1.00
199A: Plano-----	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Very limited Cutbanks cave Frost action	1.00 1.00
199B: Plano-----	Very limited Water erosion Slope	1.00 0.25	Very limited Water erosion Slope	1.00 0.25	Very limited Cutbanks cave Frost action	1.00 1.00
212C2: Thebes-----	Very limited Water erosion Slope	1.00 0.99	Very limited Water erosion Slope	1.00 0.99	Very limited Cutbanks cave Frost action	1.00 1.00
242A: Kendall-----	Very limited Water erosion Depth to saturated zone	1.00 1.00	Very limited Water erosion Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Frost action Cutbanks cave	1.00 1.00 1.00 0.10
244A: Hartsburg-----	Very limited Depth to saturated zone Water erosion	1.00 1.00	Very limited Water erosion Ponding Depth to saturated zone	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Frost action Cutbanks cave	1.00 1.00 1.00 1.00 0.10
249A: Edinburg-----	Very limited Depth to saturated zone Water erosion Restricted permeability	1.00 1.00 0.96	Very limited Water erosion Ponding Depth to saturated zone Restricted permeability	1.00 1.00 1.00 0.96	Very limited Ponding Depth to saturated zone Frost action Cutbanks cave	1.00 1.00 1.00 1.00 0.10

Table 27.--Water Management (Part 2)--Continued

Map symbol and soil name	Grassed waterways and surface drains		Terraces and diversions		Drainage	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
257A:						
Clarksdale-----	Very limited		Very limited		Very limited	
	Water erosion	1.00	Water erosion	1.00	Depth to	1.00
	Depth to	1.00	Depth to	1.00	saturated zone	
	saturated zone		saturated zone	1.00	Frost action	1.00
	Restricted	0.21	Restricted	0.21	Cutbanks cave	0.10
	permeability		permeability			
259C2:						
Assumption-----	Very limited		Very limited		Very limited	
	Water erosion	1.00	Water erosion	1.00	Frost action	1.00
	Slope	0.99	Depth to	1.00	Depth to	0.99
	Restricted	0.43	saturated zone		saturated zone	
	permeability		Slope	0.99	Cutbanks cave	0.10
	Depth to	0.25	Restricted	0.43		
	saturated zone		permeability			
259D2:						
Assumption-----	Very limited		Very limited		Very limited	
	Slope	1.00	Water erosion	1.00	Frost action	1.00
	Water erosion	1.00	Slope	1.00	Depth to	0.99
	Restricted	0.43	Depth to	1.00	saturated zone	
	permeability		saturated zone		Slope	0.96
	Depth to	0.25	Restricted	0.43	Cutbanks cave	0.10
	saturated zone		permeability			
279B:						
Rozetta-----	Very limited		Very limited		Very limited	
	Water erosion	1.00	Water erosion	1.00	Frost action	1.00
	Slope	0.25	Slope	0.25	Depth to	0.15
					saturated zone	
					Cutbanks cave	0.10
280C2:						
Fayette-----	Very limited		Very limited		Very limited	
	Water erosion	1.00	Water erosion	1.00	Frost action	1.00
	Slope	0.99	Slope	0.99	Cutbanks cave	0.10
280D2:						
Fayette-----	Very limited		Very limited		Very limited	
	Slope	1.00	Water erosion	1.00	Frost action	1.00
	Water erosion	1.00	Slope	1.00	Slope	0.96
					Cutbanks cave	0.10
280D3:						
Fayette-----	Very limited		Very limited		Very limited	
	Slope	1.00	Water erosion	1.00	Frost action	1.00
	Water erosion	1.00	Slope	1.00	Slope	0.96
					Cutbanks cave	0.10
533:						
Urban land-----	Not rated		Not rated		Not rated	
536:						
Dumps-----	Not rated		Not rated		Not rated	

Table 27.--Water Management (Part 2)--Continued

Map symbol and soil name	Grassed waterways and surface drains		Terraces and diversions		Drainage	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
549G: Marseilles-----	Very limited Slope	1.00	Very limited Water erosion	1.00	Very limited Slope	1.00
	Water erosion	1.00	Slope	1.00	Frost action	1.00
	Restricted permeability	0.96	Restricted permeability	0.96	Cutbanks cave	0.10
	Depth to soft bedrock	0.10	Depth to soft bedrock	0.10	Depth to soft bedrock	0.10
567C2: Elkhart-----	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Very limited Frost action	1.00
	Slope	0.99	Slope	0.99	Depth to saturated zone	0.16
					Cutbanks cave	0.10
567D2: Elkhart-----	Very limited Slope	1.00	Very limited Water erosion	1.00	Very limited Frost action	1.00
	Water erosion	1.00	Slope	1.00	Slope	0.96
					Depth to saturated zone	0.16
					Cutbanks cave	0.10
630C2: Navlys-----	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Very limited Frost action	1.00
	Slope	0.99	Slope	0.99	Depth to saturated zone	0.15
					Cutbanks cave	0.10
630D2: Navlys-----	Very limited Slope	1.00	Very limited Water erosion	1.00	Very limited Frost action	1.00
	Water erosion	1.00	Slope	1.00	Slope	0.96
					Depth to saturated zone	0.15
					Cutbanks cave	0.10
630D3: Navlys-----	Very limited Slope	1.00	Very limited Water erosion	1.00	Very limited Frost action	1.00
	Water erosion	1.00	Slope	1.00	Slope	0.96
					Depth to saturated zone	0.15
					Cutbanks cave	0.10
679B: Blackberry-----	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Very limited Frost action	1.00
	Slope	0.25	Depth to saturated zone	1.00	Depth to saturated zone	0.99
	Depth to saturated zone	0.25	Slope	0.25	Cutbanks cave	0.10
684B: Broadwell-----	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Very limited Cutbanks cave	1.00
	Slope	0.25	Slope	0.25	Frost action	1.00

Table 27.--Water Management (Part 2)--Continued

Map symbol and soil name	Grassed waterways and surface drains		Terraces and diversions		Drainage	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
684C2: Broadwell-----	Very limited Water erosion Slope	1.00 0.99	Very limited Water erosion Slope	1.00 0.99	Very limited Cutbanks cave Frost action	1.00 1.00
685B: Middletown-----	Very limited Water erosion Slope	1.00 0.25	Very limited Water erosion Slope	1.00 0.25	Very limited Cutbanks cave Frost action	1.00 1.00
685C2: Middletown-----	Very limited Water erosion Slope	1.00 0.99	Very limited Water erosion Slope	1.00 0.99	Very limited Cutbanks cave Frost action	1.00 1.00
705A: Buckhart-----	Very limited Water erosion Depth to saturated zone	1.00 0.25	Very limited Water erosion Depth to saturated zone	1.00 1.00	Very limited Frost action Depth to saturated zone Cutbanks cave	1.00 0.99 0.10
705B: Buckhart-----	Very limited Water erosion Slope Depth to saturated zone	1.00 0.25 0.25	Very limited Water erosion Depth to saturated zone Slope	1.00 1.00 0.25	Very limited Frost action Depth to saturated zone Cutbanks cave	1.00 0.99 0.10
712A: Spaulding-----	Very limited Depth to saturated zone Water erosion	1.00 1.00	Very limited Water erosion Ponding Depth to saturated zone	1.00 1.00 1.00	Very limited Ponding Depth to saturated zone Frost action Cutbanks cave	1.00 1.00 1.00 0.10
801C: Orthents-----	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Very limited Frost action	1.00
830: Landfills-----	Not rated		Not rated		Not rated	
862: Pits, sand-----	Not rated		Not rated		Not rated	
864: Pits, quarries-----	Not rated		Not rated		Not rated	
3073A: Ross-----	Somewhat limited Water erosion	0.89	Somewhat limited Water erosion	0.89	Very limited Flooding Depth to saturated zone Cutbanks cave	1.00 0.15 0.10

Table 27.--Water Management (Part 2)--Continued

Map symbol and soil name	Grassed waterways and surface drains		Terraces and diversions		Drainage	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
3074A: Radford-----	Very limited Water erosion Depth to saturated zone	1.00 1.00	Very limited Water erosion Depth to saturated zone	1.00 1.00	Very limited Flooding Depth to saturated zone Frost action Cutbanks cave	1.00 1.00 1.00 0.10
3077A: Huntsville-----	Somewhat limited Water erosion	0.89	Somewhat limited Water erosion	0.89	Very limited Flooding Frost action Depth to saturated zone Cutbanks cave	1.00 1.00 0.15 0.10
3107A: Sawmill-----	Very limited Depth to saturated zone Water erosion	1.00 0.56	Very limited Ponding Depth to saturated zone Water erosion	1.00 1.00 0.56	Very limited Ponding Flooding Depth to saturated zone Frost action Cutbanks cave	1.00 1.00 1.00 1.00 0.10
3284A: Tice-----	Very limited Depth to saturated zone Water erosion	1.00 0.89	Very limited Depth to saturated zone Water erosion	1.00 0.89	Very limited Flooding Depth to saturated zone Frost action Cutbanks cave	1.00 1.00 1.00 0.10
3405A: Zook-----	Very limited Depth to saturated zone Restricted permeability Water erosion	1.00 0.96 0.56	Very limited Depth to saturated zone Restricted permeability Water erosion	1.00 0.96 0.56	Very limited Flooding Depth to saturated zone Frost action Cutbanks cave Too clayey	1.00 1.00 1.00 0.10 0.01
3451A: Lawson-----	Very limited Depth to saturated zone Water erosion	1.00 0.89	Very limited Depth to saturated zone Water erosion	1.00 0.89	Very limited Flooding Depth to saturated zone Frost action Cutbanks cave	1.00 1.00 1.00 0.10
7037A: Worthen-----	Somewhat limited Water erosion	0.89	Somewhat limited Water erosion	0.89	Very limited Frost action Cutbanks cave	1.00 0.10
7075B: Drury-----	Very limited Water erosion Slope	1.00 0.25	Very limited Water erosion Slope	1.00 0.25	Very limited Frost action Cutbanks cave	1.00 0.10

Table 27.--Water Management (Part 2)--Continued

Map symbol and soil name	Grassed waterways and surface drains		Terraces and diversions		Drainage	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
7148A: Proctor-----	Very limited Water erosion	1.00	Very limited Water erosion	1.00	Very limited Frost action Cutbanks cave	1.00 0.10
7242A: Kendall-----	Very limited Water erosion Depth to saturated zone	1.00 1.00	Very limited Water erosion Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Frost action Cutbanks cave	1.00 1.00 1.00 0.10
8396A: Vesser-----	Very limited Depth to saturated zone Water erosion	1.00 1.00	Very limited Water erosion Depth to saturated zone	1.00 1.00	Very limited Depth to saturated zone Frost action Flooding Cutbanks cave	1.00 1.00 1.00 0.60 0.10

Table 28.--Water Management (Part 3)

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value column range from 0.01 to 1.00. The larger the value, the greater the limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Sprinkler irrigation	
	Rating class and limiting features	Value
8D: Hickory-----	Somewhat limited Slope	0.98
8D2: Hickory-----	Somewhat limited Slope	0.98
8D3: Hickory-----	Somewhat limited Slope	0.98
8F: Hickory-----	Very limited Slope	1.00
17A: Keomah-----	Very limited Wetness	1.00
36B: Tama-----	Not limited	
36C2: Tama-----	Very limited Erosion hazard Slope	1.00 0.06
43A: Ipava-----	Very limited Wetness	1.00
45A: Denny-----	Very limited Ponding Wetness	1.00 1.00
46A: Herrick-----	Very limited Wetness	1.00
50A: Virden-----	Very limited Ponding Wetness	1.00 1.00
68A: Sable-----	Very limited Ponding Wetness	1.00 1.00

Table 28.--Water Management (Part 3)--Continued

Map symbol and soil name	Sprinkler irrigation	
	Rating class and limiting features	Value
86B: Osco-----	Not limited	
86C2: Osco-----	Very limited Erosion hazard Slope	1.00 0.06
112A: Cowden-----	Very limited Ponding Wetness	1.00 1.00
119C2: Elco-----	Very limited Erosion hazard Slope	1.00 0.06
119D: Elco-----	Very limited Erosion hazard Slope	1.00 0.98
119D2: Elco-----	Very limited Erosion hazard Slope	1.00 0.98
119D3: Elco-----	Very limited Erosion hazard Slope	1.00 0.98
127B: Harrison-----	Not limited	
127C2: Harrison-----	Very limited Erosion hazard Slope	1.00 0.06
131C2: Alvin-----	Somewhat limited Slope	0.06
131D2: Alvin-----	Somewhat limited Slope	0.98
134A: Camden-----	Not limited	
134B: Camden-----	Very limited Erosion hazard	1.00
134C2: Camden-----	Very limited Erosion hazard Slope	1.00 0.06

Table 28.--Water Management (Part 3)--Continued

Map symbol and soil name	Sprinkler irrigation	
	Rating class and limiting features	Value
138A:		
Shiloh-----	Very limited	
	Ponding	1.00
	Wetness	1.00
198A:		
Elburn-----	Very limited	
	Wetness	1.00
199A:		
Plano-----	Not limited	
199B:		
Plano-----	Not limited	
212C2:		
Thebes-----	Very limited	
	Erosion hazard	1.00
	Slope	0.06
242A:		
Kendall-----	Very limited	
	Wetness	1.00
244A:		
Hartsburg-----	Very limited	
	Ponding	1.00
	Wetness	1.00
249A:		
Edinburg-----	Very limited	
	Ponding	1.00
	Wetness	1.00
257A:		
Clarksdale-----	Very limited	
	Wetness	1.00
259C2:		
Assumption-----	Very limited	
	Erosion hazard	1.00
	Slope	0.06
259D2:		
Assumption-----	Very limited	
	Erosion hazard	1.00
	Slope	0.98
279B:		
Rozetta-----	Very limited	
	Erosion hazard	1.00
280C2:		
Fayette-----	Very limited	
	Erosion hazard	1.00
	Slope	0.06
280D2:		
Fayette-----	Very limited	
	Erosion hazard	1.00
	Slope	0.98

Table 28.--Water Management (Part 3)--Continued

Map symbol and soil name	Sprinkler irrigation	
	Rating class and limiting features	Value
280D3: Fayette-----	Very limited Erosion hazard Slope	1.00 0.98
533: Urban land-----	Not rated	
536: Dumps-----	Not rated	
549G: Marseilles-----	Very limited Soft bedrock within a depth of 40 inches Slope Limited available water capacity	1.00 1.00 0.11
567C2: Elkhart-----	Very limited Erosion hazard Slope	1.00 0.06
567D2: Elkhart-----	Very limited Erosion hazard Slope	1.00 0.98
630C2: Navlys-----	Very limited Erosion hazard Slope	1.00 0.06
630D2: Navlys-----	Very limited Erosion hazard Slope	1.00 0.98
630D3: Navlys-----	Very limited Erosion hazard Slope	1.00 0.98
679B: Blackberry-----	Not limited	
684B: Broadwell-----	Not limited	
684C2: Broadwell-----	Very limited Erosion hazard Slope	1.00 0.06
685B: Middletown-----	Very limited Erosion hazard	1.00

Table 28.--Water Management (Part 3)--Continued

Map symbol and soil name	Sprinkler irrigation	
	Rating class and limiting features	Value
685C2: Middletown-----	Very limited Erosion hazard Slope	1.00 0.06
705A: Buckhart-----	Not limited	
705B: Buckhart-----	Not limited	
712A: Spaulding-----	Very limited Ponding Wetness	1.00 1.00
801C: Orthents-----	Very limited Permeability Erosion hazard Slope	1.00 1.00 0.06
830: Landfills-----	Not rated	
862: Pits, sand-----	Not rated	
864: Pits, quarries-----	Not rated	
3073A: Ross-----	Very limited Flooding	1.00
3074A: Radford-----	Very limited Flooding Wetness	1.00 1.00
3077A: Huntsville-----	Very limited Flooding	1.00
3107A: Sawmill-----	Very limited Ponding Flooding Wetness	1.00 1.00 1.00
3284A: Tice-----	Very limited Flooding Wetness	1.00 1.00
3405A: Zook-----	Very limited Flooding Wetness Limited available water capacity	1.00 1.00 0.20

Table 28.--Water Management (Part 3)--Continued

Map symbol and soil name	Sprinkler irrigation	
	Rating class and limiting features	Value
3451A: Lawson-----	Very limited Flooding Wetness	1.00 1.00
7037A: Worthen-----	Not limited	
7075B: Drury-----	Very limited Erosion hazard	1.00
7148A: Proctor-----	Not limited	
7242A: Kendall-----	Very limited Wetness	1.00
8396A: Vesser-----	Very limited Wetness	1.00

Table 29.--Engineering Index Properties
(Absence of an entry indicates that the data were not estimated.)

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
8D:												
Hickory-----	0-4	Silt loam	CL	A-4, A-6	0	0-5	95-100	90-100	90-100	75-95	20-35	3-15
	4-12	Loam, clay loam	CL, ML, CL-ML	A-6, A-4	0	0-5	95-100	90-100	75-100	55-100	20-35	3-15
	12-46	Clay loam, silty clay loam, gravelly clay loam, loam	CL	A-6, A-7	0-1	0-5	95-100	75-100	70-95	65-80	30-50	15-30
	46-58	Gravelly clay loam, loam, clay loam	CL, CL-ML	A-4, A-6	0-1	0-5	85-100	75-95	70-95	60-80	20-40	5-20
	58-63	Loam, sandy loam, gravelly clay loam	CL, CL-ML, SC, SC-SM	A-2, A-4, A-6	0-1	0-5	85-100	70-95	45-95	25-75	20-40	5-20
8D2:												
Hickory-----	0-6	Loam	CL	A-4, A-6	0	0-5	95-100	90-100	90-100	75-95	20-35	8-15
	6-47	Clay loam, silty clay loam, gravelly clay loam	CL	A-6, A-7	0-1	0-5	95-100	75-100	70-95	65-80	30-50	15-30
	47-60	Sandy loam, loam, gravelly clay loam	CL, CL-ML	A-4, A-6	0-1	0-5	85-100	75-95	70-95	60-80	20-40	5-20
8D3:												
Hickory-----	0-8	Clay loam	CL	A-6, A-7	0	0-5	95-100	90-100	80-95	70-85	30-50	15-30
	8-46	Clay loam, loam, gravelly clay loam	CL	A-6, A-7	0-1	0-5	95-100	75-100	70-95	65-80	30-50	15-30
	46-58	Clay loam, loam, gravelly clay loam	CL, CL-ML, SC, SC-SM	A-2, A-4, A-6	0-1	0-5	85-100	70-95	45-95	25-75	20-40	5-20
	58-80	Loam, sandy loam, gravelly clay loam	CL, CL-ML, SC, SC-SM	A-2, A-4, A-6	0-1	0-5	85-100	70-95	45-95	25-75	20-40	5-20
8F:												
Hickory-----	0-12	Silt loam	CL	A-4, A-6	0	0-5	95-100	90-100	75-100	55-100	20-35	5-15
	12-46	Clay loam, silty clay loam, loam, gravelly clay loam	CL	A-6, A-7	0-1	0-5	85-100	70-100	65-95	50-85	30-50	15-30
	46-58	Clay loam, loam, gravelly clay loam	CL, CL-ML, SC, SC-SM	A-2, A-4, A-6	0-1	0-5	85-100	70-95	45-95	25-75	20-40	5-20
	58-80	Loam, sandy loam, gravelly clay loam	CL, CL-ML, SC, SC-SM	A-2, A-4, A-6	0-1	0-5	85-100	70-95	45-95	25-75	20-40	5-20

Table 29.--Engineering Index Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
17A:												
Keomah-----	0-11	Silt loam	CL, ML	A-4, A-6	0	0	100	100	100	95-100	25-35	10-15
	11-18	Silt loam	CL, ML	A-4, A-6	0	0	100	100	100	95-100	25-35	10-20
	18-33	Silty clay, silty clay loam	CH, CL	A-7-6	0	0	100	100	100	95-100	45-55	25-30
	33-51	Silty clay loam	CL, ML	A-6, A-7-6	0	0	100	100	100	95-100	35-45	15-25
	51-89	Silt loam	CL, CL-ML, ML	A-6, A-4	0	0	100	100	100	95-100	25-35	5-15
36B:												
Tama-----	0-19	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	100	95-100	25-40	5-15
	19-58	Silty clay loam	CL	A-7	0	0	100	100	100	95-100	40-50	15-25
	58-80	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-45	15-25
36C2:												
Tama-----	0-8	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	100	95-100	25-40	5-15
	8-30	Silty clay loam	CL	A-7	0	0	100	100	100	95-100	40-50	15-25
	30-60	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-45	15-25
43A:												
Ipava-----	0-20	Silt loam, silty clay loam	CL	A-6	0	0	100	100	95-100	90-100	25-40	10-20
	20-35	Silty clay loam, silty clay	CH, CL	A-7	0	0	100	100	95-100	90-100	45-70	25-40
	35-60	Silt loam, silty clay loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	90-100	25-40	5-20
45A:												
Denny-----	0-9	Silt loam	CL	A-4, A-6	0	0	100	100	95-100	95-100	30-40	8-15
	9-22	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	95-100	25-40	5-15
	22-45	Silty clay loam, silty clay	CH, CL	A-6, A-7	0	0	100	100	95-100	95-100	35-60	15-35
	45-70	Silt loam, silty clay loam	CL	A-6	0	0	100	100	95-100	95-100	25-40	11-20
46A:												
Herrick-----	0-13	Silt loam	CL, ML	A-4, A-6	0	0	100	100	95-100	90-100	30-40	5-15
	13-39	Silty clay loam, silty clay	CH, CL	A-7-6	0	0	100	100	95-100	90-100	45-60	25-40
	39-60	Silty clay loam, silt loam	CL	A-6, A-7-6	0	0	100	100	95-100	90-100	35-50	20-35
	60-80	Silt loam, loam, silty clay loam, clay loam	CL	A-6	0	0	100	100	90-100	80-100	30-40	10-20

Table 29.--Engineering Index Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
50A: Virden-----	0-16	Silty clay loam	CL	A-6, A-7	0	0	100	100	95-100	95-100	30-50	10-25
	16-49	Silty clay, silty clay loam	CH, CL	A-7-6	0	0	100	100	95-100	95-100	40-60	20-40
	49-60	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	95-100	90-100	30-50	10-25
68A: Sable-----	0-17	Silty clay loam	CH, CL, MH, ML	A-7	0	0	100	100	95-100	95-100	30-55	10-25
	17-23	Silty clay loam	CH, CL, MH, ML	A-7	0	0	100	100	95-100	95-100	41-65	15-35
	23-60	Silty clay loam, silt loam	CH, CL	A-7	0	0	100	100	95-100	95-100	40-55	20-35
86B: Osc-----	0-14	Silt loam	CL, ML	A-6, A-4	0	0	100	100	100	95-100	35-45	7-20
	14-55	Silty clay loam, silt loam	CL	A-6, A-7-6	0	0	100	100	100	95-100	40-50	15-25
	55-60	Silt loam, silty clay loam	CL, ML	A-6, A-4	0	0	100	100	100	95-100	35-45	7-25
86C2: Osc-----	0-9	Silt loam	CL, ML	A-6, A-4	0	0	100	100	95-100	95-100	35-45	10-20
	9-34	Silty clay loam, silt loam	CL	A-6, A-7-6	0	0	100	100	95-100	95-100	40-50	15-25
	34-60	Silt loam, silty clay loam	CL, ML	A-6, A-4	0	0	100	100	95-100	95-100	35-45	7-25
112A: Cowden-----	0-8	Silt loam	CL	A-6	0	0	100	100	95-100	90-100	30-40	15-25
	8-19	Silt loam	CL	A-6	0	0	100	100	95-100	90-100	30-40	15-25
	19-50	Silty clay loam, silty clay	CH	A-7-6	0	0	100	100	95-100	95-100	50-60	30-35
	50-80	Silt loam, loam, silty clay loam, clay loam	CL, ML	A-6, A-7-6	0	0	100	95-100	85-100	70-100	30-50	7-30
119C2: Elco-----	0-8	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	90-100	25-40	5-15
	8-31	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	95-100	85-100	25-45	10-30
	31-60	Silty clay loam, loam, clay	CL	A-6, A-7	0	0	100	90-100	80-100	60-95	25-50	10-30

Table 29.--Engineering Index Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
119D:												
Elco-----	0-4	Silt loam	CL	A-4, A-6	0	0	100	100	95-100	95-100	25-40	5-15
	4-12	Silt loam	CL-ML, CL	A-4, A-6	0	0	100	100	95-100	95-100	25-40	5-15
	12-26	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	95-100	85-100	25-45	10-30
	26-80	Silty clay loam, loam, clay	CL	A-6, A-7	0	0	100	90-100	80-100	60-95	25-50	10-30
119D2:												
Elco-----	0-6	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	95-100	25-40	5-15
	6-28	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	95-100	85-100	25-45	10-30
	28-60	Silty clay loam, loam, clay	CL	A-6, A-7	0	0	100	90-100	80-100	60-95	25-50	10-30
119D3:												
Elco-----	0-5	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	30-45	15-30
	5-26	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	95-100	85-100	25-45	10-30
	26-60	Clay loam, silty clay loam, loam, clay	CL	A-6, A-7	0	0	100	90-100	80-100	60-95	25-50	10-30
127B:												
Harrison----	0-10	Silt loam	CL	A-4, A-6	0	0	100	100	100	95-100	30-40	8-15
	10-45	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	30-45	10-25
	45-65	Silty clay loam, clay loam, silt loam	CL	A-6, A-7	0	0-5	95-100	85-100	80-85	70-80	30-50	10-25
	65-80	Clay loam, clay, silty clay loam	CH, CL	A-6, A-7	0-1	0-5	95-100	85-100	80-95	70-90	35-55	15-30
127C2:												
Harrison----	0-8	Silt loam	CL	A-4, A-6	0	0	100	100	100	95-100	30-40	8-15
	8-45	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	30-45	10-25
	45-65	Silty clay loam, clay loam, silt loam	CL	A-6, A-7	0	0-5	95-100	85-100	80-85	70-80	30-50	10-25
	65-80	Clay loam, clay, silty clay loam	CH, CL	A-6, A-7	0-1	0-5	95-100	85-100	80-95	70-90	35-55	15-30

Table 29.--Engineering Index Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
131C2:												
Alvin-----	0-7	Fine sandy loam	ML, SM	A-2, A-4	0	0	100	100	80-95	30-60	0-25	NP-4
	7-42	Very fine sandy loam, sandy loam, loam	CL, ML, SC, SM	A-2, A-4, A-6	0	0	100	95-100	70-100	30-55	15-40	NP-15
	42-80	Very fine sand, fine sandy loam, loamy fine sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0	95-100	95-100	45-95	4-35	15-20	NP-4
131D2:												
Alvin-----	0-7	Fine sandy loam	ML, SM	A-2, A-4	0	0	100	100	80-95	30-60	0-25	NP-4
	7-14	Fine sandy loam, sandy loam, loamy fine sand	ML, SM	A-2, A-4	0	0	100	100	80-95	30-60	0-25	NP-4
	14-47	Fine sandy loam, sandy loam, loam	CL, ML, SC, SM	A-2, A-4, A-6	0	0	100	95-100	70-100	30-55	15-40	NP-15
	47-60	Fine sand, fine sandy loam, loamy fine sand	SM, SP, SP-SM	A-1, A-2, A-3	0	0	95-100	90-100	45-95	4-35	0-20	NP-4
134A:												
Camden-----	0-7	Silt loam	CL-ML, CL	A-4, A-6	0	0	100	100	95-100	90-100	20-35	5-15
	7-12	Silt loam	CL-ML, CL	A-4, A-6	0	0	100	100	95-100	90-100	20-35	5-15
	12-26	Silty clay loam, silt loam	CL	A-6	0	0	100	100	95-100	90-100	25-40	15-25
	26-53	Clay loam, silt loam, sandy loam	ML, CL, SM, SC	A-2, A-4, A-6	0	0-5	90-100	85-100	60-100	30-85	20-40	3-15
	53-60	Stratified loamy sand to silt loam	SM, SC, ML, CL	A-2-4, A-4	0	0-5	90-100	80-100	50-85	15-85	15-25	3-10
134B:												
Camden-----	0-9	Silt loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	95-100	95-100	25-35	5-15
	9-14	Silt loam	CL, ML	A-6	0	0	100	100	95-100	95-100	25-35	10-20
	14-22	Silty clay loam, silt loam	CL, ML	A-6	0	0	100	100	97-100	95-100	25-35	10-20
	22-35	Silty clay loam	CL, ML	A-6, A-7	0	0	98-100	95-100	90-100	80-100	35-45	15-25
	35-52	Clay loam, loam, sandy loam	CL, ML, SC-SM, SC	A-6	0	0	95-100	85-100	70-90	45-80	25-35	10-15
	52-80	Stratified sandy loam to loam to sandy clay loam	SC, CL-ML, CL, SC-SM	A-6, A-4, A-2	0	0	95-100	80-100	65-95	20-60	25-30	5-15

Table 29.--Engineering Index Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
134C2:												
Camden-----	0-7	Silt loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	95-100	90-100	20-35	3-15
	7-34	Silt loam, silty clay loam	CL	A-6	0	0	100	100	95-100	90-100	25-40	15-25
	34-55	Clay loam, sandy loam, silt loam	CL, ML, SC, SM	A-2, A-4, A-6	0	0-5	90-100	85-100	60-100	30-70	20-40	3-15
	55-80	Stratified loamy sand to silt loam	CL, ML, SC, SM	A-2, A-4	0	0-5	90-100	70-95	60-90	15-70	0-25	NP-10
138A:												
Shiloh-----	0-27	Silty clay loam, silty clay	CL	A-7	0	0	100	100	95-100	90-100	40-50	15-25
	27-52	Silty clay, silty clay loam	CH, CL	A-7	0	0	100	100	95-100	90-100	40-65	15-40
	52-80	Silty clay loam, silty clay, silt loam	CL	A-6, A-7	0	0	100	100	95-100	90-100	30-50	15-30
198A:												
Elburn-----	0-16	Silt loam	CL, CL-ML, ML	A-6, A-4	0	0	100	100	100	95-100	25-35	5-15
	16-49	Silty clay loam, silt loam	CL, ML	A-6, A-7-6	0	0	100	100	100	95-100	35-45	15-25
	49-58	Stratified sandy loam to silt loam	CL-ML, CL, SC-SM, ML	A-4, A-6	0	0	95-100	95-100	85-95	55-75	20-30	5-15
	58-62	Stratified loamy sand to silt loam	SC-SM, SM	A-2-4, A-4	0	0	95-100	90-100	50-80	20-45	20-25	NP-5
199A:												
Plano-----	0-14	Silt loam	CL, CL-ML, ML	A-4, A-6	0	0	100	100	95-100	90-100	20-30	5-15
	14-49	Silty clay loam, silt loam	CL, ML	A-6	0	0	100	100	95-100	90-100	25-40	10-25
	49-60	Clay loam, loam, sandy loam	CL, ML, SC, SM	A-4, A-6, A-7	0	0-1	90-100	85-95	60-90	35-75	20-45	5-25
	60-72	Stratified loamy sand to silt loam	SC, SM, CL, ML	A-2, A-4	0	0-5	90-100	70-95	40-80	15-55	0-25	NP-10
199B:												
Plano-----	0-15	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	95-100	20-30	5-15
	15-45	Silty clay loam, silt loam	CL	A-6	0	0	100	100	95-100	90-100	25-40	10-25
	45-60	Clay loam, loam, sandy loam	SC-SM, CL, CL-ML, SC	A-4, A-6, A-7	0	0-1	90-100	85-95	60-90	35-75	20-45	5-25
	60-72	Stratified silt loam to loamy sand	CL, ML, SC, SM	A-2, A-4	0	0-5	90-100	70-95	40-80	15-55	0-25	NP-10

Table 29.--Engineering Index Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
212C2: Thebes-----	0-9	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	100	95-100	25-35	5-15
	9-31	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-45	15-25
	31-40	Loam, sandy loam, clay loam	CL, ML, SC, SM	A-4, A-6	0	0	100	100	90-100	40-70	15-30	NP-13
	40-80	Fine sand, loamy fine sand, loamy sand, sandy loam	SM, SP, SP-SM	A-2, A-3	0	0	100	90-100	75-95	4-35	0-20	NP-4
242A: Kendall-----	0-7	Silt loam	CL-ML, CL	A-4, A-6	0	0	100	100	95-100	90-100	20-35	5-15
	7-11	Silt loam	CL-ML, CL	A-4, A-6	0	0	100	100	95-100	90-100	20-35	5-15
	11-51	Silty clay loam	CL	A-6, A-7	0	0	100	100	95-100	90-100	30-45	10-20
	51-58	Loam, silt loam, clay loam	CL, SC-SM	A-4, A-6	0	0	95-100	90-100	70-90	50-70	25-35	8-15
	58-74	Stratified sandy loam to silt loam	CL, CL-ML, SC-SM, SC	A-4	0	0-3	95-100	90-100	70-90	40-70	20-30	NP-10
	74-80	Stratified gravelly sandy loam to silt loam	SC-SM, SC, CL-ML, CL	A-4	0	0-5	65-95	60-80	50-70	35-50	20-30	NP-10
244A: Hartsburg---	0-17	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	17-34	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	95-100	95-100	30-50	15-30
	34-60	Silt loam, loam	CL	A-6	0	0	95-100	90-100	90-100	65-100	25-40	11-20
249A: Edinburg----	0-16	Silty clay loam	CL	A-6, A-7	0	0	100	100	95-100	90-100	35-50	16-25
	16-55	Silty clay loam, silty clay	CH, CL	A-7	0	0	100	100	95-100	90-100	45-70	25-45
	55-60	Silt loam, silty clay loam	CL	A-6, A-7	0	0	100	100	95-100	90-100	35-45	15-20
257A: Clarksdale--	0-8	Silt loam	CL	A-6	0	0	100	100	95-100	90-100	25-40	10-20
	8-16	Silt loam	CL	A-4, A-6	0	0	100	100	95-100	90-100	20-35	8-18
	16-47	Silty clay loam, silty clay	CH, CL	A-7	0	0	100	100	95-100	90-100	40-65	25-40
	47-67	Silt loam, silty clay loam	CL	A-6, A-7-6	0	0	100	100	95-100	90-100	25-45	10-25
	67-80	Silt loam	CL	A-6	0	0	95-100	95-100	95-100	90-100	25-40	10-20

Table 29.--Engineering Index Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
567C2:												
Elkhart-----	0-8	Silt loam	CL	A-4, A-6	0	0	100	100	100	95-100	25-35	8-15
	8-34	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	18-30
	34-60	Silt loam	CL	A-4, A-6	0	0	100	100	95-100	95-100	20-37	8-20
567D2:												
Elkhart-----	0-8	Silt loam	CL	A-4, A-6	0	0	100	100	100	95-100	25-35	8-15
	8-34	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	18-30
	34-60	Silt loam	CL	A-4, A-6	0	0	100	100	95-100	95-100	20-37	8-20
630C2:												
Navlys-----	0-6	Silt loam	CL	A-6, A-4	0	0	100	100	100	95-100	25-35	5-15
	6-27	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	20-30
	27-60	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	95-100	20-40	5-20
630D2:												
Navlys-----	0-6	Silt loam	CL	A-6, A-4	0	0	100	100	100	95-100	25-35	5-15
	6-27	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	20-30
	27-60	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	95-100	20-40	5-20
630D3:												
Navlys-----	0-6	Silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	20-30
	6-31	Silty clay loam, silt loam	CL	A-7, A-6	0	0	100	100	100	95-100	35-50	20-30
	31-60	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	95-100	20-40	5-20
679B:												
Blackberry--	0-16	Silt loam	CL-ML, CL	A-4, A-6	0	0	100	100	95-100	90-100	20-30	5-15
	16-47	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	95-100	90-100	25-45	10-25
	47-62	Stratified loam to silt loam, sandy loam	CL, CL-ML, ML, SC-SM, SC	A-4	0	0	95-100	95-100	70-90	45-75	20-30	5-10
	62-70	Stratified silt loam to loam to sandy loam	CL-ML, CL, ML, SC-SM, SM, SC	A-4	0	0	95-100	90-100	70-90	40-70	15-25	NP-10
684B:												
Broadwell---	0-15	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	90-100	25-45	5-20
	15-50	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	95-100	90-100	30-45	10-25
	50-55	Loamy fine sand, clay loam, fine sandy loam, loam	SC, SC-SM	A-4	0	0	100	95-100	80-90	35-50	20-30	5-10
	55-80	Loamy sand, fine sand, sand	SC-SM, SM, SP, SP-SM	A-2, A-3	0	0	100	100	75-95	4-35	0-20	NP-5

Table 29.--Engineering Index Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
684C2:												
Broadwell---	0-8	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	100	95-100	25-40	5-15
	8-46	Silt loam, silty clay loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-25
	46-49	Fine sandy loam, loam, loamy fine sand, clay loam	SC, SC-SM	A-4	0	0	100	95-100	80-90	35-50	20-30	5-10
	49-60	Loamy fine sand, loamy sand, fine sand	SC-SM, SM, SP-SM	A-2, A-3	0	0	100	90-100	75-95	5-40	0-20	NP-10
685B:												
Middletown--	0-9	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	100	90-100	25-40	5-15
	9-12	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	100	90-100	25-40	5-15
	12-44	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-30
	44-47	Clay loam, loam, fine sandy loam, loamy fine sand	CL, CL-ML	A-4, A-6	0	0	100	100	90-100	90-100	20-35	4-15
	47-80	Loamy fine sand, fine sand, loamy sand	SC, SC-SM, SM, SP-SM	A-2, A-3, A-4	0	0	100	90-100	75-95	5-40	0-20	NP-10
685C2:												
Middletown--	0-7	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	100	90-100	25-40	5-15
	7-46	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	100	95-100	35-50	15-30
	46-55	Clay loam, loam, fine sandy loam, loamy fine sand	CL, CL-ML	A-4, A-6	0	0	100	100	90-100	90-100	20-35	4-15
	55-60	Loamy fine sand, fine sand, loamy sand	SC, SC-SM, SM, SP-SM	A-2, A-3, A-4	0	0	100	90-100	75-95	5-40	0-20	NP-10
705A:												
Buckhart----	0-20	Silt loam, silty clay loam	CL, ML	A-6, A-7	0	0	100	100	100	95-100	35-45	10-20
	20-58	Silty clay loam, silt loam	CL	A-7	0	0	100	100	100	95-100	40-50	15-25
	58-60	Silty clay loam, silt loam	CL	A-6	0	0	100	100	100	95-100	30-40	11-20

Table 29.--Engineering Index Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plasticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
705B:												
Buckhart----	0-15	Silt loam	CL, ML	A-6, A-7	0	0	100	100	100	95-100	35-45	10-20
	15-67	Silty clay loam, silt loam	CL	A-7	0	0	100	100	100	95-100	40-50	15-25
	67-80	Silt loam	CL	A-6	0	0	100	100	100	95-100	30-40	11-20
712A:												
Spaulding---	0-22	Silty clay loam	CH, CL	A-7	0	0	100	100	95-100	95-100	45-60	20-35
	22-38	Silty clay loam, silt loam	CH, CL	A-7	0	0	100	100	95-100	95-100	40-60	20-35
	38-44	Silty clay loam, silt loam	CH, CL	A-6, A-7	0	0	100	100	95-100	95-100	35-55	20-35
	44-80	Silt loam	CL	A-6	0	0	100	100	95-100	95-100	30-40	10-20
801C:												
Orthents----	0-60	Silty clay loam, silt loam	CL, CL-ML	A-4, A-6, A-7	0	0	100	100	90-100	80-95	25-45	5-25
830:												
Landfills.												
862:												
Pits, sand.												
864:												
Pits, quarries.												
3073A:												
Ross-----	0-13	Silt loam, loam	CL-ML, ML, CL	A-4, A-6	0	0	90-100	90-100	80-100	65-95	20-35	NP-12
	13-43	Loam, silt loam, silty clay loam	CL, CL-ML, ML	A-4, A-6, A-7	0	0	90-100	85-100	70-100	55-95	22-45	3-20
	43-60	Gravelly sandy loam, stratified sandy loam to silt loam	CL, GM, ML, SM	A-2, A-4, A-6	0	0-5	65-100	45-100	30-100	25-80	0-30	NP-12
3074A:												
Radford----	0-12	Silt loam	ML, CL	A-4, A-6	0	0	100	100	95-100	85-100	30-40	5-15
	12-33	Silt loam	CL-ML, CL	A-4, A-6	0	0	100	100	95-100	85-100	25-35	5-15
	33-80	Silt loam, silty clay loam, clay loam	CL	A-6, A-7	0	0	100	100	95-100	70-95	35-50	15-25
3077A:												
Huntsville--	0-43	Silt loam	CL	A-6	0	0	100	95-100	90-100	85-100	25-40	10-20
	43-60	Silt loam	CL	A-6	0	0	100	95-100	90-100	85-100	20-35	10-20

Table 29.--Engineering Index Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
3107A:												
Sawmill-----	0-17	Silty clay loam	CL, ML	A-6, A-7-6	0	0	100	100	95-100	85-100	30-50	15-30
	17-32	Silty clay loam	CL, ML	A-6, A-7-6	0	0	100	95-100	95-100	85-100	30-50	15-30
	32-58	Silty clay loam	CL, ML	A-6, A-7-6	0	0	100	95-100	90-100	80-95	25-50	8-30
	58-65	Silty clay loam, clay loam, sandy loam	CL, ML, SC	A-4, A-6, A-7-6	0	0	100	90-100	75-100	40-95	20-50	8-30
3284A:												
Tice-----	0-14	Silty clay loam	CL	A-6, A-7	0	0	100	100	90-100	80-95	30-45	10-25
	14-52	Silty clay loam, silt loam	CH, CL	A-7	0	0	100	100	95-100	85-95	40-55	15-30
	52-72	Stratified loam to silty clay loam	CL, CL-ML	A-4, A-6, A-7	0	0	100	100	60-95	55-80	25-45	5-20
3405A:												
Zook-----	0-8	Silty clay loam	CH, CL	A-7	0	0	100	100	95-100	95-100	45-65	20-35
	8-55	Silty clay, silty clay loam	CH	A-7	0	0	100	100	95-100	95-100	60-85	35-55
	55-60	Silty clay loam, silty clay, silt loam	CH, CL, MH, ML	A-6, A-7	0	0	100	100	95-100	95-100	35-80	10-50
3451A:												
Lawson-----	0-14	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	90-100	85-100	20-40	5-20
	14-33	Silt loam, silty clay loam	CL, CL-ML	A-4	0	0	100	100	90-100	85-100	20-30	5-10
	33-80	Silty clay loam, silt loam	CL	A-6, A-7	0	0	100	100	90-100	60-100	20-45	10-25
7037A:												
Worthen-----	0-26	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	80-100	25-40	7-21
	26-60	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	80-100	25-40	7-21
	60-80	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	80-100	25-40	7-21
7075B:												
Drury-----	0-7	Silt loam	CL, CL-ML, ML	A-4, A-6	0	0	100	95-100	95-100	90-100	20-35	NP-15
	7-43	Silt loam	CL	A-4, A-6	0	0	100	95-100	95-100	90-100	25-35	8-15
	43-80	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	95-100	95-100	55-95	20-30	5-15
7148A:												
Proctor-----	0-16	Silt loam	CL	A-6	0	0	100	100	95-100	90-100	25-40	10-20
	16-34	Silty clay loam	CL	A-6, A-7	0	0	100	95-100	85-100	85-100	25-50	10-25
	34-53	Clay loam, loam, sandy loam	CL, CL-ML, SC, SC-SM	A-2, A-4, A-6, A-7	0	0	90-100	85-100	75-100	30-80	20-45	5-25
	53-60	Loamy sand, sandy loam	SC, SC-SM, SM	A-2, A-4	0	0	85-100	80-100	50-100	25-50	0-25	NP-10

Table 29.--Engineering Index Properties--Continued

Map symbol and soil name	Depth	USDA texture	Classification		Fragments		Percentage passing sieve number--				Liquid limit	Plas- ticity index
			Unified	AASHTO	>10 inches	3-10 inches	4	10	40	200		
	In				Pct	Pct					Pct	
7242A:												
Kendall-----	0-9	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	85-95	20-35	5-15
	9-14	Silt loam	CL, CL-ML	A-4, A-6	0	0	100	100	95-100	85-95	15-30	5-15
	14-54	Silty clay loam	CL	A-6, A-7	0	0	100	100	95-100	85-95	30-45	10-20
	54-60	Stratified sandy loam to clay loam	CL, CL-ML, SC, SC-SM	A-2, A-4	0	0-5	98-100	80-90	60-90	30-70	15-25	4-15
8396A:												
Vesser-----	0-14	Silt loam	CL	A-6	0	0	100	100	98-100	95-100	30-40	10-20
	14-26	Silt loam	CL	A-6	0	0	100	100	98-100	95-100	30-40	10-20
	26-80	Silty clay loam	CL	A-7	0	0	100	100	98-100	95-100	40-50	15-25

Table 30.--Physical Properties of the Soils

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer. Absence of an entry indicates that data were not estimated.)

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										Kw	Kf	T		
	In	Pct	Pct	Pct	g/cc	In/hr	In/in	Pct	Pct					
8D: Hickory-----	0-4	15-45	30-66	19-25	1.30-1.50	0.6-2	0.20-0.22	0.0-2.9	1.0-2.0	.32	.32	5	6	48
	4-12	15-45	33-70	15-30	1.30-1.50	0.6-2	0.20-0.22	0.0-2.9	0.0-0.5	.37	.37			
	12-46	15-45	20-58	20-35	1.45-1.65	0.6-2	0.15-0.19	3.0-5.9	0.0-0.5	.28	.32			
	46-58	30-45	23-55	15-32	1.50-1.70	0.6-2	0.11-0.19	0.0-2.9	0.0-0.5	.28	.32			
	58-63	30-45	23-55	15-30	1.50-1.75	0.6-2	0.10-0.15	0.0-2.9	0.0-0.5	.28	.32			
8D2: Hickory-----	0-6	15-45	30-66	19-25	1.30-1.50	0.6-2	0.20-0.22	0.0-2.9	1.0-2.0	.32	.32	5	6	48
	6-47	15-45	20-58	27-35	1.45-1.65	0.6-2	0.15-0.19	3.0-5.9	0.0-0.5	.28	.32			
	47-60	30-45	23-55	15-32	1.50-1.70	0.6-2	0.11-0.19	0.0-2.9	0.0-0.5	.28	.32			
8D3: Hickory-----	0-8	20-40	30-50	27-35	1.40-1.65	0.6-2	0.17-0.19	3.0-5.9	0.5-1.0	.28	.32	4	6	48
	8-46	20-45	30-50	24-35	1.45-1.65	0.6-2	0.15-0.19	3.0-5.9	0.0-0.5	.28	.32			
	46-58	25-49	28-50	15-32	1.50-1.70	0.6-2	0.11-0.19	0.0-2.9	0.0-0.5	.28	.32			
	58-80	30-55	25-50	15-30	1.50-1.75	0.6-2	0.10-0.15	0.0-2.9	0.0-0.5	.28	.32			
8F: Hickory-----	0-12	10-30	45-70	18-25	1.30-1.50	0.6-2	0.20-0.22	0.0-2.9	1.0-2.0	.32	.32	5	6	48
	12-46	15-45	30-50	24-35	1.45-1.65	0.6-2	0.15-0.19	3.0-5.9	0.0-0.5	.28	.32			
	46-58	25-49	28-50	15-32	1.50-1.70	0.6-2	0.11-0.19	0.0-2.9	0.0-0.5	.28	.32			
	58-80	30-55	25-50	15-30	1.50-1.75	0.6-2	0.10-0.15	0.0-2.9	0.0-0.5	.28	.32			
17A: Keomah-----	0-11	0-7	67-84	16-26	1.35-1.45	0.6-2	0.19-0.24	0.0-2.9	1.0-3.0	.43	.43	5	6	48
	11-18	0-7	67-84	16-26	1.40-1.60	0.2-0.6	0.17-0.21	0.0-2.9	0.1-1.0	.49	.49			
	18-33	0-7	51-65	35-42	1.30-1.40	0.06-0.2	0.15-0.19	6.0-8.9	0.1-0.5	.37	.37			
	33-51	0-7	58-73	27-35	1.35-1.45	0.2-0.6	0.16-0.20	3.0-5.9	0.1-0.5	.37	.37			
	51-89	0-7	66-85	15-27	1.40-1.60	0.6-2	0.19-0.22	0.0-2.9	0.0-0.2	.49	.49			
36B: Tama-----	0-19	0-7	66-76	24-27	1.25-1.30	0.6-2	0.22-0.24	3.0-5.9	3.0-4.0	.28	.28	5	6	48
	19-58	0-7	58-73	27-35	1.30-1.35	0.6-2	0.18-0.20	3.0-5.9	0.0-1.0	.37	.37			
	58-80	0-7	65-78	22-28	1.35-1.40	0.6-2	0.18-0.20	3.0-5.9	0.0-0.5	.49	.49			
36C2: Tama-----	0-8	0-7	66-76	24-27	1.25-1.30	0.6-2	0.22-0.24	3.0-5.9	2.0-3.0	.37	.37	5	6	48
	8-30	0-7	58-73	27-35	1.30-1.35	0.6-2	0.18-0.20	3.0-5.9	0.0-1.0	.37	.37			
	30-60	0-7	65-78	22-28	1.35-1.40	0.6-2	0.18-0.20	3.0-5.9	0.0-0.5	.49	.49			

Table 30.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind	Wind
										Kw	Kf	T	erodi- bility group	erodi- bility index
43A:														
Ipava -----	0-20	0-7	63-80	20-30	1.15-1.35	0.6-2	0.22-0.24	3.0-5.9	4.0-5.0	.28	.28	5	6	48
	20-35	0-7	50-65	35-43	1.25-1.50	0.2-0.6	0.11-0.20	6.0-8.9	0.5-1.0	.37	.37			
	35-60	0-7	63-80	20-30	1.30-1.55	0.2-0.6	0.20-0.22	3.0-5.9	0.0-0.5	.49	.49			
45A:														
Denny -----	0-9	0-7	66-80	20-27	1.25-1.45	0.6-2	0.22-0.24	0.0-2.9	2.0-3.0	.37	.37	5	6	48
	9-22	0-7	71-85	15-22	1.25-1.45	0.2-0.6	0.18-0.20	0.0-2.9	0.0-0.5	.43	.43			
	22-45	0-7	48-65	35-45	1.20-1.40	0.06-0.2	0.11-0.22	6.0-8.9	0.0-1.0	.37	.37			
	45-70	0-7	58-75	25-35	1.40-1.60	0.2-0.6	0.20-0.22	3.0-5.9	0.0-0.5	.43	.43			
46A:														
Herrick -----	0-13	1-7	66-78	20-27	1.15-1.30	0.6-2	0.22-0.24	3.0-5.9	3.0-4.0	.28	.28	5	6	48
	13-39	1-7	51-63	35-42	1.20-1.40	0.2-0.6	0.12-0.17	6.0-8.9	0.0-1.0	.37	.37			
	39-60	1-7	55-73	25-40	1.20-1.40	0.2-0.6	0.16-0.20	3.0-5.9	0.0-0.5	.37	.37			
	60-80	5-30	45-70	20-30	1.30-1.50	0.2-0.6	0.16-0.21	3.0-5.9	0.0-0.5	.49	.49			
50A:														
Viriden -----	0-16	0-7	58-73	27-35	1.20-1.40	0.6-2	0.21-0.24	3.0-5.9	3.0-6.0	.24	.24	5	7	38
	16-49	0-7	49-65	35-42	1.20-1.45	0.2-0.6	0.11-0.20	6.0-8.9	0.0-2.0	.37	.37			
	49-60	0-7	60-75	25-33	1.25-1.55	0.2-0.6	0.18-0.22	3.0-5.9	0.0-0.5	.43	.43			
68A:														
Sable -----	0-17	0-7	58-73	27-35	1.15-1.35	0.6-2	0.21-0.23	3.0-5.9	5.0-6.0	.24	.24	5	7	38
	17-23	0-7	58-73	27-35	1.20-1.40	0.6-2	0.18-0.20	3.0-5.9	2.0-4.0	.24	.24			
	23-60	0-7	58-76	24-35	1.30-1.50	0.6-2	0.18-0.20	3.0-5.9	0.2-1.0	.37	.37			
86B:														
Oscos -----	0-14	0-7	67-80	20-26	1.25-1.30	0.6-2	0.22-0.24	3.0-5.9	3.0-4.0	.28	.28	5	6	48
	14-55	0-7	58-76	24-35	1.30-1.35	0.6-2	0.18-0.20	3.0-5.9	0.0-1.0	.37	.37			
	55-60	0-7	63-80	20-30	1.35-1.40	0.6-2	0.18-0.20	3.0-5.9	0.0-0.5	.49	.49			
86C2:														
Oscos -----	0-9	0-7	67-80	20-26	1.25-1.30	0.6-2	0.22-0.24	3.0-5.9	2.0-3.0	.37	.37	5	6	48
	9-34	0-7	58-76	24-35	1.30-1.35	0.6-2	0.18-0.20	3.0-5.9	0.0-1.0	.37	.37			
	34-60	0-7	63-80	20-30	1.35-1.40	0.6-2	0.18-0.20	3.0-5.9	0.0-0.5	.49	.49			
112A:														
Cowden -----	0-8	1-7	68-80	17-27	1.30-1.50	0.6-2	0.22-0.24	0.0-2.9	2.0-3.0	.37	.37	3	6	48
	8-19	1-7	68-80	17-27	1.25-1.45	0.06-0.2	0.18-0.20	0.0-2.9	0.0-0.5	.49	.49			
	19-50	1-7	51-63	35-42	1.35-1.60	0.06-0.2	0.12-0.20	6.0-8.9	0.0-1.0	.37	.37			
	50-80	5-30	45-70	20-30	1.50-1.70	0.2-0.6	0.17-0.22	3.0-5.9	0.0-0.5	.49	.49			

Table 30.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										Kw	Kf	T		
	In	Pct	Pct	Pct	g/cc	In/hr	In/in	Pct	Pct					
119C2: Elco-----	0-8	0-7	66-80	20-27	1.20-1.35	0.6-2	0.22-0.24	0.0-2.9	1.0-2.0	.43	.43	5	6	48
	8-31	0-7	58-77	23-35	1.25-1.45	0.6-2	0.18-0.21	3.0-5.9	0.0-0.5	.37	.37			
	31-60	15-35	20-60	25-45	1.45-1.70	0.06-0.6	0.14-0.20	6.0-8.9	0.0-0.2	.28	.28			
119D: Elco-----	0-4	0-7	66-80	20-27	1.15-1.30	0.6-2	0.22-0.24	0.0-2.9	1.0-3.0	.43	.43	5	6	48
	4-12	0-7	66-82	18-27	1.20-1.35	0.6-2	0.22-0.24	0.0-2.9	0.1-1.0	.49	.49			
	12-26	0-7	58-77	23-35	1.25-1.45	0.6-2	0.18-0.21	3.0-5.9	0.0-0.5	.37	.37			
	26-80	15-35	20-60	25-45	1.45-1.70	0.06-0.6	0.14-0.20	6.0-8.9	0.0-0.2	.28	.28			
119D2: Elco-----	0-6	0-7	66-80	20-27	1.20-1.35	0.6-2	0.22-0.24	0.0-2.9	1.0-2.0	.43	.43	5	6	48
	6-28	0-7	58-77	23-35	1.25-1.45	0.6-2	0.18-0.21	3.0-5.9	0.0-0.5	.37	.37			
	28-60	15-35	20-60	25-45	1.45-1.70	0.06-0.6	0.14-0.20	6.0-8.9	0.0-0.2	.28	.28			
119D3: Elco-----	0-5	0-7	58-73	27-35	1.20-1.35	0.6-2	0.18-0.21	3.0-5.9	0.5-1.0	.37	.37	4	7	38
	5-26	0-7	58-77	23-35	1.25-1.45	0.6-2	0.18-0.21	3.0-5.9	0.0-0.5	.37	.37			
	26-60	15-35	20-60	25-45	1.45-1.70	0.06-0.6	0.14-0.20	6.0-8.9	0.0-0.2	.28	.28			
127B: Harrison-----	0-10	0-5	68-80	20-27	1.15-1.30	0.6-2	0.22-0.24	0.0-2.9	3.0-4.0	.28	.28	5	6	48
	10-45	0-5	60-75	25-35	1.25-1.40	0.6-2	0.18-0.22	3.0-5.9	0.2-1.0	.37	.37			
	45-65	5-30	45-75	20-35	1.30-1.45	0.6-2	0.14-0.20	3.0-5.9	0.0-0.2	.37	.37			
	65-80	5-30	30-65	30-50	1.50-1.70	0.06-0.2	0.10-0.19	6.0-8.9	0.0-0.2	.37	.37			
127C2: Harrison-----	0-8	0-5	68-80	20-27	1.15-1.30	0.6-2	0.22-0.24	0.0-2.9	2.0-3.0	.37	.37	5	6	48
	8-45	0-5	60-75	25-35	1.25-1.40	0.6-2	0.18-0.22	3.0-5.9	0.2-1.0	.37	.37			
	45-65	5-30	45-75	20-35	1.30-1.45	0.6-2	0.14-0.20	3.0-5.9	0.0-0.2	.37	.37			
	65-80	5-30	30-65	30-50	1.50-1.70	0.06-0.2	0.10-0.19	6.0-8.9	0.0-0.2	.37	.37			
131C2: Alvin-----	0-7	55-70	15-35	10-15	1.45-1.65	2-6	0.14-0.17	0.0-2.9	0.5-1.0	.20	.20	5	3	86
	7-42	45-70	12-40	15-18	1.40-1.65	2-6	0.14-0.18	0.0-2.9	0.0-0.5	.24	.24			
	42-80	65-95	2-32	3-10	1.45-1.65	2-6	0.10-0.15	0.0-2.9	0.0-0.3	.15	.15			
131D2: Alvin-----	0-7	55-70	15-35	10-15	1.45-1.65	2-6	0.14-0.17	0.0-2.9	0.5-1.0	.20	.20	5	3	86
	7-14	55-70	15-35	10-15	1.45-1.65	2-6	0.10-0.17	0.0-2.9	0.0-0.5	.24	.24			
	14-47	45-70	12-40	15-18	1.40-1.65	2-6	0.14-0.18	0.0-2.9	0.0-0.5	.24	.24			
	47-60	65-95	2-32	3-10	1.45-1.65	2-6	0.10-0.15	0.0-2.9	0.0-0.3	.15	.15			

Table 30.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										Kw	Kf	T		
	In	Pct	Pct	Pct	g/cc	In/hr	In/in	Pct	Pct					
134A: Camden-----	0-7	0-10	63-86	14-27	1.35-1.55	0.6-2	0.22-0.24	0.0-2.9	1.0-3.0	.43	.43	5	6	48
	7-12	0-10	65-88	12-25	1.35-1.55	0.6-2	0.20-0.22	0.0-2.9	0.1-1.0	.49	.49			
	12-26	0-10	55-78	22-35	1.40-1.60	0.6-2	0.14-0.24	3.0-5.9	0.1-0.5	.37	.37			
	26-53	15-65	5-67	18-30	1.45-1.65	0.6-2	0.11-0.22	3.0-5.9	0.0-0.5	.32	.32			
	53-60	45-80	0-50	5-25	1.40-1.70	0.6-6	0.08-0.20	0.0-2.9	0.0-0.5	.28	.28			
134B: Camden-----	0-9	2-7	66-83	15-27	1.35-1.55	0.6-2	0.22-0.24	0.0-2.9	1.0-2.5	.43	.43	5	5	56
	9-14	2-7	66-83	15-27	1.40-1.60	0.6-2	0.17-0.20	0.0-2.9	0.1-1.0	.49	.49			
	14-22	2-7	66-76	22-27	1.30-1.50	0.6-2	0.18-0.23	0.0-2.9	0.1-0.5	.37	.37			
	22-35	3-15	50-70	27-35	1.35-1.55	0.6-2	0.16-0.20	3.0-5.9	0.1-0.5	.43	.43			
	35-52	25-54	28-50	18-27	1.45-1.65	0.6-2	0.11-0.18	0.0-2.9	0.0-0.5	.32	.32			
	52-80	45-65	25-45	10-22	1.50-1.70	0.6-6	0.08-0.15	0.0-2.9	0.0-0.5	.28	.28			
134C2: Camden-----	0-7	0-10	63-86	14-27	1.35-1.55	0.6-2	0.22-0.24	0.0-2.9	1.0-2.0	.43	.43	5	6	48
	7-34	0-10	55-78	22-35	1.40-1.60	0.6-2	0.16-0.20	3.0-5.9	0.1-0.5	.37	.37			
	34-55	15-65	5-67	18-30	1.45-1.65	0.6-2	0.11-0.17	0.0-2.9	0.0-0.5	.32	.32			
	55-80	65-80	0-80	5-20	1.40-1.70	0.6-6	0.05-0.15	0.0-2.9	0.0-0.5	.28	.28			
138A: Shiloh-----	0-27	0-7	53-65	35-42	1.30-1.50	0.2-0.6	0.18-0.21	6.0-8.9	4.0-6.0	.24	.24	5	7	38
	27-52	0-7	48-65	35-45	1.35-1.55	0.2-0.6	0.09-0.18	6.0-8.9	0.5-2.0	.37	.37			
	52-80	0-7	48-75	25-45	1.30-1.50	0.2-0.6	0.18-0.20	6.0-8.9	0.2-0.5	.43	.43			
198A: Elburn-----	0-16	0-10	63-84	15-27	1.25-1.35	0.6-2	0.22-0.24	0.0-2.9	3.5-5.0	.28	.28	5	6	48
	16-49	0-10	55-74	25-35	1.35-1.45	0.6-2	0.18-0.21	3.0-5.9	0.5-1.5	.37	.37			
	49-58	30-50	23-55	15-27	1.40-1.60	0.6-2	0.14-0.17	0.0-2.9	0.0-0.5	.32	.32			
	58-62	45-80	5-50	5-15	1.55-1.75	2-6	0.08-0.11	0.0-2.9	0.0-0.1	.28	.28			
199A: Plano-----	0-14	0-10	63-82	18-27	1.10-1.30	0.6-2	0.22-0.24	0.0-2.9	3.0-5.0	.28	.28	5	6	48
	14-49	0-10	55-80	20-35	1.20-1.40	0.6-2	0.18-0.20	3.0-5.9	0.2-1.0	.37	.37			
	49-60	15-70	0-70	15-32	1.30-1.55	0.6-6	0.09-0.16	0.0-2.9	0.1-0.5	.32	.32			
	60-72	15-80	0-80	5-20	1.50-1.70	2-6	0.11-0.22	0.0-2.9	0.1-0.5	.28	.28			
199B: Plano-----	0-15	0-10	63-82	18-27	1.10-1.30	0.6-2	0.22-0.24	0.0-2.9	3.0-5.0	.28	.28	5	6	48
	15-45	0-10	55-80	20-35	1.20-1.40	0.6-2	0.18-0.20	3.0-5.9	0.2-1.0	.37	.37			
	45-60	15-70	0-70	15-30	1.30-1.55	0.6-6	0.09-0.16	0.0-2.9	0.1-0.5	.32	.32			
	60-72	15-80	0-80	5-20	1.50-1.70	2-6	0.11-0.22	0.0-2.9	0.1-0.5	.28	.28			

Table 30.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind erodi- bility group	Wind erodi- bility index
										Kw	Kf	T		
	In	Pct	Pct	Pct	g/cc	In/hr	In/in	Pct	Pct					
212C2: Thebes-----	0-9	5-20	40-80	15-25	1.30-1.35	0.6-2	0.20-0.22	0.0-2.9	1.0-2.0	.43	.43	5	6	48
	9-31	0-15	50-75	25-35	1.30-1.45	0.6-2	0.18-0.20	3.0-5.9	0.0-0.5	.37	.37			
	31-40	30-60	0-35	15-27	1.35-1.55	0.6-2	0.12-0.19	0.0-2.9	0.0-0.5	.32	.32			
	40-80	70-90	0-27	3-10	1.45-1.65	6-20	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
242A: Kendall-----	0-7	0-10	63-80	20-27	1.15-1.30	0.6-2	0.22-0.24	0.0-2.9	1.0-3.0	.43	.43	5	6	48
	7-11	0-10	65-82	18-25	1.35-1.55	0.6-2	0.20-0.22	0.0-2.9	0.1-1.0	.49	.49			
	11-51	0-10	55-73	27-35	1.30-1.50	0.6-2	0.18-0.20	3.0-5.9	0.0-0.5	.37	.37			
	51-58	30-50	33-50	15-27	1.45-1.55	0.6-2	0.11-0.14	0.0-2.9	0.0-0.5	.32	.32			
	58-74	30-50	35-55	5-20	1.40-1.60	0.6-2	0.14-0.17	0.0-2.9	0.0-0.5	.32	.37			
	74-80	30-50	30-50	5-20	1.55-1.75	0.6-2	0.11-0.15	0.0-2.9	0.0-0.2	.24	.32			
244A: Hartsburg-----	0-17	0-5	62-73	27-33	1.15-1.35	0.6-2	0.21-0.24	3.0-5.9	3.0-5.0	.24	.24	5	7	38
	17-34	0-5	60-75	25-35	1.20-1.50	0.6-2	0.18-0.20	3.0-5.9	0.2-1.0	.37	.37			
	34-60	0-30	43-80	20-27	1.30-1.55	0.6-2	0.20-0.22	0.0-2.9	0.0-0.2	.32	.32			
249A: Edinburg-----	0-16	0-7	58-73	27-35	1.10-1.30	0.6-2	0.21-0.24	6.0-8.9	3.0-6.0	.24	.24	3	7	38
	16-55	0-7	47-65	35-46	1.20-1.40	0.06-0.2	0.13-0.20	6.0-8.9	0.2-1.0	.37	.37			
	55-60	0-7	63-78	22-30	1.30-1.50	0.2-2	0.18-0.22	3.0-5.9	0.0-0.2	.49	.49			
257A: Clarksdale-----	0-8	0-7	66-80	20-27	1.30-1.50	0.6-2	0.22-0.25	3.0-5.9	2.0-3.0	.37	.37	5	6	48
	8-16	0-7	66-85	15-27	1.25-1.50	0.2-0.6	0.20-0.22	0.0-2.9	0.0-1.0	.43	.43			
	16-47	0-7	48-65	35-45	1.30-1.50	0.2-0.6	0.11-0.20	6.0-8.9	0.0-0.5	.37	.37			
	47-67	0-7	63-80	20-30	1.40-1.60	0.2-0.6	0.20-0.22	3.0-5.9	0.0-0.5	.43	.43			
	67-80	0-7	66-82	18-27	1.40-1.60	0.2-0.6	0.20-0.22	0.0-2.9	0.0-0.5	.49	.49			
259C2: Assumption-----	0-8	0-7	66-80	20-27	1.25-1.45	0.6-2	0.23-0.25	0.0-2.9	2.0-3.0	.37	.37	5	6	48
	8-24	0-7	58-75	25-35	1.20-1.40	0.6-2	0.18-0.22	3.0-5.9	0.0-1.0	.37	.37			
	24-60	20-30	25-50	30-45	1.45-1.65	0.06-0.6	0.14-0.20	6.0-8.9	0.0-0.5	.28	.28			
259D2: Assumption-----	0-7	0-7	66-80	20-27	1.25-1.45	0.6-2	0.23-0.25	0.0-2.9	2.0-3.0	.37	.37	5	6	48
	7-28	0-7	58-75	25-35	1.20-1.40	0.6-2	0.18-0.22	3.0-5.9	0.0-1.0	.37	.37			
	28-60	20-30	25-50	30-45	1.45-1.65	0.06-0.6	0.14-0.20	6.0-8.9	0.0-0.5	.28	.28			
279B: Rozetta-----	0-7	0-7	66-85	15-27	1.20-1.40	0.6-2	0.22-0.24	0.0-2.9	1.0-3.0	.43	.43	5	6	48
	7-11	0-7	66-88	12-27	1.20-1.40	0.6-2	0.22-0.24	0.0-2.9	0.1-1.0	.49	.49			
	11-55	0-7	58-73	27-35	1.35-1.55	0.6-2	0.18-0.22	3.0-5.9	0.0-0.5	.37	.37			
	55-60	0-7	63-80	20-30	1.40-1.60	0.6-2	0.20-0.22	0.0-2.9	0.0-0.5	.49	.49			

Table 30.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind	Wind
										Kw	Kf	T	erodi- bility group	erodi- bility index
	In	Pct	Pct	Pct	g/cc	In/hr	In/in	Pct	Pct					
280C2: Fayette-----	0-8	0-7	66-75	25-27	1.35-1.45	0.6-2	0.18-0.20	3.0-5.9	1.0-2.0	.43	.43	5	6	48
	8-64	0-7	58-75	25-35	1.30-1.45	0.6-2	0.18-0.20	3.0-5.9	0.0-0.5	.37	.37			
	64-80	0-7	67-78	22-26	1.45-1.50	0.6-2	0.18-0.20	3.0-5.9	0.0-0.5	.49	.49			
280D2: Fayette-----	0-6	0-7	66-75	25-27	1.35-1.45	0.6-2	0.18-0.20	3.0-5.9	1.0-2.0	.43	.43	5	6	48
	6-48	0-7	58-75	25-35	1.30-1.45	0.6-2	0.18-0.20	3.0-5.9	0.0-0.5	.37	.37			
	48-60	0-7	67-78	22-26	1.45-1.50	0.6-2	0.18-0.20	3.0-5.9	0.0-0.5	.49	.49			
280D3: Fayette-----	0-8	0-7	61-73	27-32	1.35-1.45	0.6-2	0.18-0.20	3.0-5.9	0.5-1.0	.37	.37	4	7	38
	8-36	0-7	58-75	25-35	1.30-1.45	0.6-2	0.18-0.20	3.0-5.9	0.0-0.5	.37	.37			
	36-60	0-7	67-78	22-26	1.45-1.50	0.6-2	0.18-0.20	3.0-5.9	0.0-0.5	.49	.49			
533: Urban land.														
536: Dumps.														
549G: Marseilles-----	0-10	0-15	58-80	20-27	1.20-1.40	0.6-2	0.20-0.24	0.0-2.9	1.0-3.0	.32	.32	3	6	48
	10-35	0-25	43-73	27-42	1.35-1.60	0.06-0.2	0.09-0.20	3.0-6.0	0.0-0.5	.37	.37			
	35-60	---	---	---	---	0.0015-0.2	---	---	---	---	---			
567C2: Elkhart-----	0-8	0-7	66-80	20-27	1.15-1.35	0.6-2	0.22-0.24	0.0-2.9	2.0-3.0	.37	.37	5	6	38
	8-34	0-7	58-75	25-35	1.25-1.45	0.6-2	0.18-0.20	3.0-5.9	0.0-1.0	.37	.37			
	34-60	0-7	66-80	18-27	1.35-1.55	0.6-2	0.20-0.22	0.0-2.9	0.0-0.5	.49	.49			
567D2: Elkhart-----	0-8	0-7	66-80	20-27	1.15-1.35	0.6-2	0.22-0.24	0.0-2.9	2.0-3.0	.37	.37	5	6	48
	8-34	0-7	58-75	25-35	1.25-1.45	0.6-2	0.18-0.20	3.0-5.9	0.0-1.0	.37	.37			
	34-60	0-7	66-80	18-27	1.35-1.55	0.6-2	0.20-0.22	0.0-2.9	0.0-0.5	.49	.49			
630C2: Navlys-----	0-6	0-7	66-80	20-27	1.20-1.40	0.6-2	0.20-0.22	3.0-5.9	1.0-2.0	.43	.43	5	6	48
	6-27	0-7	58-75	25-35	1.30-1.50	0.6-2	0.18-0.20	3.0-5.9	0.0-0.5	.37	.37			
	27-60	0-7	66-82	18-27	1.30-1.50	0.6-2	0.20-0.22	0.0-2.9	0.0-0.5	.49	.49			
630D2: Navlys-----	0-6	0-7	66-80	20-27	1.20-1.40	0.6-2	0.20-0.22	3.0-5.9	1.0-2.0	.43	.43	5	6	48
	6-27	0-7	58-75	25-35	1.30-1.50	0.6-2	0.18-0.20	3.0-5.9	0.0-0.5	.37	.37			
	27-60	0-7	66-82	18-27	1.30-1.50	0.6-2	0.20-0.22	0.0-2.9	0.0-0.5	.49	.49			

Table 30.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind	Wind
										Kw	Kf	T	erodi- bility group	erodi- bility index
	In	Pct	Pct	Pct	g/cc	In/hr	In/in	Pct	Pct					
630D3:														
Navlys-----	0-6	0-7	61-73	27-32	1.25-1.45	0.6-2	0.20-0.22	3.0-5.9	0.5-1.0	.37	.37	4	7	38
	6-31	0-7	58-75	25-35	1.30-1.50	0.6-2	0.18-0.20	3.0-5.9	0.0-0.5	.37	.37			
	31-60	0-7	66-82	18-27	1.30-1.50	0.6-2	0.20-0.22	0.0-2.9	0.0-0.5	.49	.49			
679B:														
Blackberry-----	0-16	0-10	63-82	18-27	1.10-1.30	0.6-2	0.22-0.24	0.0-2.9	3.0-5.0	.28	.28	5	6	48
	16-47	0-10	55-75	25-35	1.20-1.40	0.6-2	0.18-0.20	3.0-5.9	0.2-1.0	.37	.37			
	47-62	30-50	35-55	15-25	1.30-1.55	0.6-2	0.11-0.22	3.0-5.9	0.1-0.5	.32	.32			
	62-70	30-50	35-55	5-20	1.50-1.70	0.6-2	0.05-0.19	0.0-2.9	0.0-0.1	.37	.37			
684B:														
Broadwell-----	0-15	5-20	55-75	20-27	1.25-1.45	0.6-2	0.23-0.26	0.0-2.9	3.0-4.0	.28	.28	5	6	48
	15-50	0-10	55-76	24-35	1.35-1.60	0.6-2	0.14-0.24	3.0-5.9	0.0-1.0	.37	.37			
	50-55	40-80	0-35	10-28	1.30-1.35	2-6	0.11-0.17	0.0-2.9	0.0-0.5	.32	.32			
	55-80	50-90	0-47	3-10	1.55-1.75	6-20	0.08-0.11	0.0-2.9	0.0-0.5	.15	.15			
684C2:														
Broadwell-----	0-8	0-10	63-76	24-27	1.25-1.30	0.6-2	0.22-0.24	0.0-2.9	2.0-3.0	.37	.37	5	6	48
	8-46	0-10	55-80	20-35	1.30-1.35	0.6-2	0.19-0.22	3.0-5.9	0.0-1.0	.37	.37			
	46-49	40-80	0-35	10-28	1.30-1.35	2-6	0.11-0.17	0.0-2.9	0.0-0.5	.32	.32			
	49-60	50-90	0-47	3-10	1.30-1.35	6-20	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
685B:														
Middletown-----	0-9	0-10	63-80	20-27	1.20-1.40	0.6-2	0.22-0.24	0.0-2.9	1.0-3.0	.43	.43	5	6	48
	9-12	0-10	63-85	15-27	1.20-1.40	0.6-2	0.22-0.24	0.0-2.9	0.1-1.0	.49	.49			
	12-44	0-10	55-75	25-35	1.25-1.45	0.6-2	0.18-0.21	3.0-5.9	0.0-0.5	.37	.37			
	44-47	40-80	0-35	10-30	1.35-1.60	0.6-2	0.15-0.19	0.0-2.9	0.0-0.5	.32	.32			
	47-80	50-90	0-47	3-10	1.45-1.65	2-20	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
685C2:														
Middletown-----	0-7	0-10	63-80	20-27	1.20-1.40	0.6-2	0.22-0.24	0.0-2.9	1.0-2.0	.43	.43	5	6	48
	7-46	0-10	55-75	25-35	1.25-1.45	0.6-2	0.18-0.21	3.0-5.9	0.0-0.5	.37	.37			
	46-55	40-80	0-35	10-35	1.35-1.60	0.6-2	0.15-0.19	0.0-2.9	0.0-0.5	.32	.32			
	55-60	50-90	0-47	3-10	1.45-1.65	2-20	0.05-0.10	0.0-2.9	0.0-0.5	.15	.15			
705A:														
Buckhart-----	0-20	0-7	63-80	20-30	1.25-1.30	0.6-2	0.22-0.24	3.0-5.9	3.0-4.0	.28	.28	5	6	48
	20-58	0-7	58-75	25-35	1.30-1.35	0.6-2	0.18-0.20	3.0-5.9	0.2-1.0	.37	.37			
	58-60	0-7	66-82	18-27	1.35-1.45	0.6-2	0.20-0.22	3.0-5.9	0.0-0.5	.49	.49			
705B:														
Buckhart-----	0-15	0-7	67-80	20-26	1.25-1.30	0.6-2	0.22-0.24	3.0-5.9	3.0-4.0	.28	.28	5	6	48
	15-67	0-7	58-75	25-35	1.30-1.35	0.6-2	0.18-0.20	3.0-5.9	0.2-1.0	.37	.37			
	67-80	0-7	66-82	18-27	1.35-1.45	0.6-2	0.20-0.22	3.0-5.9	0.0-0.5	.49	.49			

Table 30.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind	Wind
										Kw	Kf	T	erodi- bility group	erodi- bility index
	In	Pct	Pct	Pct	g/cc	In/hr	In/in	Pct	Pct					
712A: Spaulding-----	0-22	0-7	58-73	27-35	1.05-1.25	0.6-2	0.21-0.24	3.0-5.9	4.0-6.0	.24	.24	5	4L	86
	22-38	0-7	58-75	25-35	1.20-1.50	0.6-2	0.18-0.22	3.0-5.9	0.5-2.0	.37	.37			
	38-44	0-7	58-78	22-35	1.25-1.55	0.6-2	0.17-0.22	3.0-5.9	0.5-1.0	.37	.37			
	44-80	0-7	66-80	20-27	1.30-1.55	0.6-2	0.20-0.22	0.0-2.9	0.0-0.5	.49	.49			
801C: Orthents-----	0-60	0-15	50-80	20-35	1.35-1.55	0.2-2	0.18-0.22	3.0-5.9	0.2-1.0	.49	.49	5	6	48
830: Landfills.														
862: Pits, sand.														
864: Pits, quarries.														
3073A: Ross-----	0-13	20-45	28-65	15-27	1.20-1.45	0.6-2	0.19-0.24	0.0-2.9	2.0-4.0	.32	.32	5	5	56
	13-43	20-45	23-62	18-32	1.20-1.50	0.6-2	0.16-0.22	0.0-2.9	1.0-3.0	.32	.32			
	43-60	40-70	5-55	5-25	1.35-1.60	0.6-6	0.05-0.18	0.0-2.9	0.2-0.5	.32	.32			
3074A: Radford-----	0-12	0-15	58-82	18-27	1.40-1.60	0.6-2	0.22-0.24	0.0-2.9	2.0-4.0	.32	.32	5	6	48
	12-33	0-15	58-82	18-27	1.40-1.60	0.6-2	0.20-0.22	0.0-2.9	0.0-2.0	.49	.49			
	33-80	5-30	35-71	24-35	1.35-1.55	0.6-2	0.18-0.20	3.0-5.9	0.0-1.0	.32	.32			
3077A: Huntsville-----	0-43	0-15	58-82	18-27	1.15-1.35	0.6-2	0.22-0.24	3.0-5.9	2.0-4.0	.32	.32	5	6	48
	43-60	5-30	43-77	18-27	1.20-1.40	0.6-2	0.20-0.22	3.0-5.9	0.2-0.5	.49	.49			
3107A: Sawmill-----	0-17	0-15	50-73	27-35	1.20-1.40	0.6-2	0.21-0.23	3.0-5.9	4.0-5.0	.28	.28	5	7	38
	17-32	0-15	50-73	27-35	1.20-1.40	0.6-2	0.21-0.23	3.0-5.9	1.0-4.0	.28	.28			
	32-58	5-20	45-68	27-35	1.30-1.45	0.6-2	0.17-0.20	3.0-5.9	0.2-2.0	.32	.32			
	58-65	5-55	10-77	18-35	1.35-1.50	0.6-2	0.15-0.19	3.0-5.9	0.2-1.0	.32	.32			
3284A: Tice-----	0-14	0-15	50-73	27-35	1.25-1.45	0.6-2	0.21-0.24	3.0-5.9	2.0-4.0	.28	.28	5	7	38
	14-52	5-20	45-71	24-35	1.30-1.50	0.6-2	0.18-0.20	3.0-5.9	0.0-1.0	.32	.32			
	52-72	5-40	30-80	15-30	1.40-1.60	0.6-2	0.11-0.18	3.0-5.9	0.0-1.0	.32	.32			

Table 30.--Physical Properties of the Soils--Continued

Map symbol and soil name	Depth	Sand	Silt	Clay	Moist bulk density	Permea- bility (Ksat)	Available water capacity	Linear extensi- bility	Organic matter	Erosion factors			Wind	Wind
										Kw	Kf	T	erodi- bility group	erodi- bility index
	In	Pct	Pct	Pct	g/cc	In/hr	In/in	Pct	Pct					
3405A: Zook-----	0-8	0-15	45-65	35-40	1.30-1.35	0.2-0.6	0.21-0.23	6.0-8.9	4.0-5.0	.28	.28	5	7	38
	8-55	0-15	40-64	36-45	1.30-1.45	0.06-0.2	0.11-0.13	6.0-8.9	2.0-4.0	.28	.28			
	55-60	0-15	40-80	20-45	1.30-1.45	0.06-0.6	0.11-0.22	6.0-8.9	0.0-1.0	.32	.32			
3451A: Lawson-----	0-14	0-15	58-90	10-27	1.20-1.55	0.6-2	0.22-0.24	0.0-2.9	2.0-4.0	.32	.32	5	5	56
	14-33	0-15	55-90	10-30	1.20-1.55	0.6-2	0.18-0.22	0.0-2.9	2.0-4.0	.32	.32			
	33-80	5-40	30-77	18-30	1.55-1.65	0.6-2	0.18-0.20	3.0-5.9	0.0-1.0	.49	.49			
7037A: Worthen-----	0-26	0-15	63-88	12-22	1.20-1.40	0.6-2	0.22-0.24	0.0-2.9	2.0-4.0	.32	.32	5	6	48
	26-60	0-15	59-85	15-26	1.20-1.40	0.6-2	0.20-0.22	0.0-2.9	0.5-2.0	.49	.49			
	60-80	10-25	51-75	15-24	1.20-1.40	0.6-2	0.20-0.22	0.0-2.9	0.0-0.5	.49	.49			
7075B: Drury-----	0-7	1-15	70-80	15-25	1.20-1.40	0.6-2	0.20-0.24	0.0-2.9	1.0-3.0	.43	.43	5	5	56
	7-43	1-15	65-80	18-25	1.25-1.45	0.6-2	0.20-0.22	0.0-2.9	0.0-0.2	.49	.49			
	43-80	5-25	65-77	15-22	1.30-1.50	0.6-2	0.12-0.21	0.0-2.9	0.0-0.2	.49	.49			
7148A: Proctor-----	0-16	0-10	63-82	18-27	1.10-1.30	0.6-2	0.22-0.24	0.0-2.9	3.0-4.0	.28	.28	5	6	48
	16-34	0-10	55-73	27-35	1.20-1.45	0.6-2	0.18-0.20	3.0-5.9	0.2-1.0	.37	.37			
	34-53	15-70	0-70	15-30	1.30-1.55	0.6-2	0.13-0.16	3.0-5.9	0.1-0.5	.32	.32			
	53-60	65-80	5-30	5-15	1.40-1.70	2-6	0.08-0.10	0.0-2.9	0.0-0.5	.28	.28			
7242A: Kendall-----	0-9	0-10	63-80	20-27	1.15-1.30	0.6-2	0.22-0.24	0.0-2.9	1.0-3.0	.43	.43	5	6	48
	9-14	0-10	65-82	18-25	1.25-1.45	0.6-2	0.20-0.22	0.0-2.9	0.1-1.0	.49	.49			
	14-54	0-10	55-73	27-35	1.30-1.50	0.6-2	0.18-0.20	3.0-5.9	0.0-0.5	.37	.37			
	54-60	65-80	5-30	5-15	1.55-1.70	0.6-2	0.11-0.20	0.0-2.9	0.0-0.5	.28	.28			
8396A: Vesser-----	0-14	0-15	59-80	20-26	1.30-1.35	0.6-2	0.20-0.24	3.0-5.9	2.0-3.0	.32	.32	5	6	48
	14-26	0-15	63-82	18-22	1.35-1.40	0.6-2	0.18-0.22	3.0-5.9	0.1-1.0	.37	.37			
	26-80	5-20	45-65	30-35	1.40-1.45	0.6-2	0.17-0.21	3.0-5.9	0.0-1.0	.32	.32			

Table 31.--Chemical Properties of the Soils

(Absence of an entry indicates that data were not estimated.)

Map symbol and soil name	Depth	Soil reaction	Cation- exchange capacity	Calcium carbonate equivalent
	In	pH	meq/100 g	Pct
8D:				
Hickory-----	0-4	4.5-7.3	14-19	0
	4-12	4.5-7.3	9.0-14	0
	12-46	4.5-7.3	16-22	0
	46-58	5.1-8.4	9.0-19	0-15
	58-63	5.6-8.4	5.0-15	0-25
8D2:				
Hickory-----	0-6	4.5-7.3	14-19	0
	6-47	4.5-7.3	16-22	0
	47-60	5.1-8.4	9.0-19	0-15
8D3:				
Hickory-----	0-8	4.5-7.3	17-23	0
	8-46	4.5-7.3	16-22	0
	46-58	5.1-7.8	9.0-19	0-15
	58-80	5.6-7.8	5.0-15	0-25
8F:				
Hickory-----	0-12	4.5-7.3	14-19	0
	12-46	4.5-7.3	16-22	0
	46-58	5.1-7.8	9.0-19	0-15
	58-80	5.6-7.8	5.0-15	0-25
17A:				
Keomah-----	0-11	5.1-7.3	10-26	0
	11-18	5.1-7.3	9.0-24	0
	18-33	5.1-5.5	28-41	0
	33-51	5.6-7.3	16-29	0
	51-89	6.1-7.3	8.0-18	0
36B:				
Tama-----	0-19	5.1-7.3	25-30	0
	19-58	5.1-6.5	25-30	0
	58-80	5.6-7.3	25-30	0
36C2:				
Tama-----	0-8	5.1-7.3	25-30	0
	8-30	5.1-6.5	25-30	0
	30-60	5.6-7.3	25-30	0
43A:				
Ipava-----	0-20	5.6-7.3	20-27	0
	20-35	5.6-7.8	22-27	0
	35-60	6.1-8.4	12-19	0-15
45A:				
Denny-----	0-9	5.6-7.3	18-24	0
	9-22	5.6-6.5	9.0-15	0
	22-45	5.6-6.5	21-29	0
	45-70	5.6-7.8	15-21	0
46A:				
Herrick-----	0-13	5.1-7.3	18-24	0
	13-39	4.5-6.0	21-25	0
	39-60	5.6-7.3	15-25	0
	60-80	5.6-7.8	12-18	0-10

Table 31.--Chemical Properties of the Soils--Continued

Map symbol and soil name	Depth	Soil reaction	Cation- exchange capacity	Calcium carbonate equivalent
	In	pH	meq/100 g	Pct
50A:				
Virден-----	0-16	5.6-7.8	24-30	0
	16-49	5.6-7.8	21-27	0
	49-60	5.6-8.4	15-20	0-25
68A:				
Sable-----	0-17	5.6-7.3	26-33	0
	17-23	5.6-7.3	20-30	0
	23-60	5.6-7.8	15-23	0
86B:				
Oscо-----	0-14	5.1-7.3	18-25	0
	14-55	5.1-6.5	15-23	0
	55-60	5.6-7.3	12-18	0
86C2:				
Oscо-----	0-9	5.1-7.3	18-25	0
	9-34	5.1-6.5	15-23	0
	34-60	5.1-7.3	12-18	0
112A:				
Cowden-----	0-8	5.6-7.3	14-22	0
	8-19	4.5-6.0	10-17	0
	19-50	4.5-7.3	21-27	0
	50-80	5.6-7.8	12-17	0
119C2:				
Elco-----	0-8	5.6-7.3	14-22	0
	8-31	5.1-7.8	14-22	0
	31-60	5.1-7.8	15-27	0
119D:				
Elco-----	0-4	5.6-7.3	14-22	0
	4-12	5.6-7.3	14-22	0
	12-26	5.1-7.8	14-22	0
	26-80	5.1-7.8	15-27	0
119D2:				
Elco-----	0-6	5.6-7.3	14-22	0
	6-28	5.1-7.8	14-22	0
	28-60	5.1-7.8	15-27	0
119D3:				
Elco-----	0-5	5.6-7.3	16-22	0
	5-26	5.1-7.8	14-22	0
	26-60	5.1-7.8	15-27	0
127B:				
Harrison-----	0-10	6.1-7.3	16-24	0
	10-45	5.1-6.5	15-23	0
	45-65	5.6-7.3	12-21	0
	65-80	5.1-7.8	18-30	0-20
127C2:				
Harrison-----	0-8	6.1-7.3	16-24	0
	8-45	5.1-6.5	15-23	0
	45-65	5.6-7.3	12-21	0
	65-80	5.1-7.8	18-30	0-20

Table 31.--Chemical Properties of the Soils--Continued

Map symbol and soil name	Depth		Soil reaction	Cation- exchange capacity	Calcium carbonate equivalent
	In	pH	meq/100 g	Pct	
131C2:					
Alvin-----	0-7	4.5-7.3	7.0-11	0	
	7-42	4.5-7.3	9.0-12	0	
	42-80	5.1-8.4	2.0-7.0	0-25	
131D2:					
Alvin-----	0-7	4.5-7.3	7.0-11	0	
	7-14	4.5-7.3	6.0-11	0	
	14-47	4.5-7.3	9.0-12	0	
	47-60	5.1-8.4	2.0-7.0	0-25	
134A:					
Camden-----	0-7	5.1-7.3	10-22	0	
	7-12	5.1-7.3	7.0-17	0	
	12-26	5.1-7.3	13-22	0	
	26-53	5.1-7.3	11-19	0	
	53-60	5.1-8.4	3.0-16	0-5	
134B:					
Camden-----	0-9	6.1-7.3	11-29	0	
	9-14	6.1-7.3	9.0-24	0	
	14-22	6.1-7.3	10-25	0	
	22-35	5.6-6.5	16-29	0	
	35-52	5.1-6.5	9.0-19	0	
	52-80	5.6-7.8	8.0-15	0-25	
134C2:					
Camden-----	0-7	5.1-7.3	10-20	0	
	7-34	5.1-7.3	13-22	0	
	34-55	5.1-7.3	10-19	0	
	55-80	5.1-8.4	3.0-12	0-5	
138A:					
Shiloh-----	0-27	6.1-7.3	29-34	0	
	27-52	6.1-7.8	22-31	0	
	52-80	6.1-8.4	15-28	0-10	
198A:					
Elburn-----	0-16	6.1-7.3	16-32	0	
	16-49	5.6-7.8	17-31	0	
	49-58	6.6-7.8	6.0-17	0-5	
	58-62	6.6-7.8	1.0-6.0	0-5	
199A:					
Plano-----	0-14	6.1-7.3	17-26	0	
	14-49	5.1-7.3	15-23	0	
	49-60	5.6-7.8	9.0-20	0	
	60-72	5.6-8.4	6.0-13	0-20	
199B:					
Plano-----	0-15	6.1-7.3	17-26	0	
	15-45	5.1-7.3	15-23	0	
	45-60	5.6-7.8	9.0-20	0	
	60-72	5.6-8.4	6.0-13	0-20	
212C2:					
Thebes-----	0-9	5.1-7.3	15-20	0	
	9-31	4.5-6.5	15-22	0	
	31-40	4.5-6.5	10-20	0	
	40-80	5.1-6.5	1.0-7.0	0	

Table 31.--Chemical Properties of the Soils--Continued

Map symbol and soil name	Depth	Soil reaction	Cation- exchange capacity	Calcium carbonate equivalent
	In	pH	meq/100 g	Pct
242A:				
Kendall-----	0-7	5.1-7.3	14-22	0
	7-11	5.1-7.3	11-17	0
	11-51	4.5-7.3	16-22	0
	51-58	5.1-7.8	9.0-19	0-15
	58-74	7.4-8.4	2.0-13	0-15
	74-80	7.4-8.4	3.0-10	0-20
244A:				
Hartsburg-----	0-17	6.1-7.8	22-30	0-5
	17-34	6.6-8.4	15-23	0-30
	34-60	7.4-8.4	12-17	10-40
249A:				
Edinburg-----	0-16	5.6-7.8	22-29	0
	16-55	5.6-7.3	21-28	0
	55-60	6.6-7.8	13-18	0-5
257A:				
Clarksdale-----	0-8	5.1-7.3	10-22	0
	8-16	5.1-7.3	9.0-18	0
	16-47	5.1-7.3	21-28	0
	47-67	6.1-8.4	12-19	0-15
	67-80	6.1-8.4	12-18	0-15
259C2:				
Assumption-----	0-8	5.6-7.3	18-24	0
	8-24	5.1-7.3	15-23	0
	24-60	5.1-7.3	18-28	0
259D2:				
Assumption-----	0-7	5.6-7.3	18-24	0
	7-28	5.1-7.3	15-23	0
	28-60	5.1-7.3	18-28	0
279B:				
Rozetta-----	0-7	5.1-7.3	10-22	0
	7-11	4.5-7.3	7.0-17	0
	11-55	4.5-6.0	16-22	0
	55-60	5.6-7.8	12-17	0-15
280C2:				
Fayette-----	0-8	5.1-7.3	18-25	0
	8-64	4.5-6.0	15-20	0
	64-80	5.1-7.8	15-20	0-15
280D2:				
Fayette-----	0-6	5.1-7.3	18-25	0
	6-48	4.5-6.0	15-20	0
	48-60	5.1-7.8	15-20	0-15
280D3:				
Fayette-----	0-8	5.1-7.3	25-30	0
	8-36	4.5-6.0	15-20	0
	36-60	5.1-7.8	15-20	0-15
533:				
Urban land.				
536:				
Dumps.				

Table 31.--Chemical Properties of the Soils--Continued

Map symbol and soil name	Depth	Soil reaction	Cation- exchange capacity	Calcium carbonate equivalent
	In	pH	meq/100 g	Pct
549G:				
Marseilles-----	0-10	5.1-7.3	14-22	0
	10-35	4.5-6.5	16-27	0
	35-60	---	---	---
567C2:				
Elkhart-----	0-8	5.6-7.8	16-24	0
	8-34	5.6-8.4	15-22	0-20
	34-60	7.4-8.4	12-21	10-40
567D2:				
Elkhart-----	0-8	5.6-7.8	16-24	0
	8-34	5.6-8.4	15-22	0-20
	34-60	7.4-8.4	12-21	10-40
630C2:				
Navlys-----	0-6	5.6-7.3	14-20	0
	6-27	5.6-7.3	15-23	0
	27-60	6.6-8.4	11-17	0-35
630D2:				
Navlys-----	0-6	5.6-7.3	14-20	0
	6-27	5.6-7.3	15-23	0
	27-60	6.6-8.4	11-17	0-35
630D3:				
Navlys-----	0-6	5.6-7.3	16-20	0
	6-31	5.6-7.3	15-23	0
	31-60	6.6-8.4	11-17	0-35
679B:				
Blackberry-----	0-16	6.1-7.3	17-26	0
	16-47	5.1-7.3	15-23	0
	47-62	5.6-8.4	9.0-22	0-10
	62-70	5.6-8.4	3.0-19	0-20
684B:				
Broadwell-----	0-15	5.6-7.3	18-27	0
	15-50	5.6-7.3	15-23	0
	50-55	5.6-7.3	15-20	0
	55-80	5.6-7.3	2.0-7.0	0
684C2:				
Broadwell-----	0-8	5.6-7.3	25-30	0
	8-46	5.6-7.3	25-30	0
	46-49	5.6-7.3	15-20	0
	49-60	5.6-7.3	5.0-10	0
685B:				
Middletown-----	0-9	6.1-7.3	14-22	0
	9-12	5.1-7.3	9.0-19	0
	12-44	4.5-6.5	15-22	0
	44-47	4.5-7.3	9.0-19	0
	47-80	5.1-7.3	1.0-7.0	0
685C2:				
Middletown-----	0-7	6.1-7.3	14-21	0
	7-46	4.5-6.5	15-22	0
	46-55	4.5-7.3	9.0-28	0
	55-60	5.1-7.3	1.0-7.0	0

Table 31.--Chemical Properties of the Soils--Continued

Map symbol and soil name	Depth	Soil reaction	Cation- exchange capacity	Calcium carbonate equivalent
	In	pH	meq/100 g	Pct
705A:				
Buckhart-----	0-20	5.6-7.3	18-25	0
	20-58	5.6-7.8	15-23	0
	58-60	6.6-7.8	12-18	0-15
705B:				
Buckhart-----	0-15	5.6-7.3	18-25	0
	15-67	5.6-7.8	15-23	0
	67-80	6.6-7.8	12-18	0-15
712A:				
Spaulding-----	0-22	7.4-8.4	24-33	10-40
	22-38	7.4-8.4	17-25	5-40
	38-44	7.4-8.4	14-23	5-40
	44-80	7.4-8.4	12-17	10-40
801C:				
Orthents-----	0-60	5.1-7.8	10-25	0-10
830:				
Landfills.				
862:				
Pits, sand.				
864:				
Pits, quarries.				
3073A:				
Ross-----	0-13	6.1-7.3	13-23	0
	13-43	6.1-7.3	12-26	0
	43-60	6.1-7.8	3.0-16	0-5
3074A:				
Radford-----	0-12	5.6-7.8	15-24	0
	12-33	6.1-7.8	11-20	0
	33-80	6.1-7.8	14-23	0-20
3077A:				
Huntsville-----	0-43	5.6-7.8	17-24	0
	43-60	5.6-7.8	11-17	0
3107A:				
Sawmill-----	0-17	6.1-7.8	24-31	0
	17-32	6.1-7.8	18-29	0
	32-58	6.1-7.8	17-25	0-10
	58-65	6.1-8.4	11-23	0-30
3284A:				
Tice-----	0-14	6.1-7.8	20-27	0
	14-52	5.6-7.8	16-23	0
	52-72	5.6-7.8	9.0-20	0-20
3405A:				
Zook-----	0-8	5.6-7.3	36-41	0
	8-55	5.6-7.8	36-41	0
	55-60	5.6-7.8	30-36	0
3451A:				
Lawson-----	0-14	6.1-7.8	11-28	0
	14-33	6.1-7.8	11-29	0
	33-80	6.1-7.8	11-23	0

Table 31.--Chemical Properties of the Soils--Continued

Map symbol and soil name	Depth	Soil reaction	Cation- exchange capacity	Calcium carbonate equivalent
	<i>In</i>	<i>pH</i>	<i>meq/100 g</i>	<i>Pct</i>
7037A:				
Worthen-----	0-26	5.6-7.3	15-21	0
	26-60	5.6-7.8	11-14	0
	60-80	6.1-8.4	9.0-14	0-25
7075B:				
Drury-----	0-7	5.6-8.4	8.0-16	0
	7-43	5.6-7.3	11-15	0
	43-80	6.1-7.8	9.0-12	0-15
7148A:				
Proctor-----	0-16	5.6-7.3	16-25	0
	16-34	5.6-6.5	16-23	0
	34-53	5.6-7.3	9.0-22	0
	53-60	6.1-7.3	3.0-7.0	0
7242A:				
Kendall-----	0-9	5.1-7.3	14-20	0
	9-14	5.1-7.3	11-16	0
	14-54	4.5-7.3	16-22	0
	54-60	5.6-8.4	6.0-16	0-15
8396A:				
Vesser-----	0-14	5.6-7.3	25-30	0
	14-26	5.1-6.5	20-25	0
	26-80	5.1-6.5	25-30	0

Table 32.--Water Features

(See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Hydro- logic group	Month	Water table			Ponding			Flooding	
			Upper limit	Lower limit	Kind	Surface water depth	Duration	Frequency	Duration	Frequency
			<i>Ft</i>	<i>Ft</i>		<i>Ft</i>				
8D: Hickory-----	B	Jan-Dec	>6.0	>6.0	---	---	---	---	---	None
8D2: Hickory-----	B	Jan-Dec	>6.0	>6.0	---	---	---	---	---	None
8D3: Hickory-----	B	Jan-Dec	>6.0	>6.0	---	---	---	---	---	None
8F: Hickory-----	B	Jan-Dec	>6.0	>6.0	---	---	---	---	---	None
17A: Keomah-----	C	Jan-May Jun-Dec	0.5-2.0 >6.0	>6.0 >6.0	Apparent ---	---	---	---	---	None None
36B: Tama-----	B	Jan-Dec	>6.0	>6.0	---	---	---	---	---	None
36C2: Tama-----	B	Jan-Dec	>6.0	>6.0	---	---	---	---	---	None
43A: Ipava-----	B	Jan-May Jun-Dec	1.0-2.0 >6.0	>6.0 >6.0	Apparent ---	---	---	---	---	None None
45A: Denny-----	D	Jan-May Jun-Dec	0.0-1.0 >6.0	>6.0 >6.0	Apparent ---	0.0-1.0 ---	Brief ---	Frequent ---	---	None None
46A: Herrick-----	B	Jan-May Jun-Dec	1.0-2.0 >6.0	>6.0 >6.0	Apparent ---	---	---	---	---	None None
50A: Virden-----	E/D	Jan-May Jun-Dec	0.0-1.0 >6.0	>6.0 >6.0	Apparent ---	0.0-0.5 ---	Brief ---	Frequent ---	---	None None
68A: Sable-----	E/D	Jan-May Jun-Dec	0.0-1.0 >6.0	>6.0 >6.0	Apparent ---	0.0-0.5 ---	Brief ---	Frequent ---	---	None None
86B: Osco-----	B	Jan Feb-Apr May-Dec	>6.0 4.0-6.0 >6.0	>6.0 >6.0 >6.0	--- Apparent ---	---	---	---	---	None None None
86C2: Osco-----	B	Jan Feb-Apr May-Dec	>6.0 4.0-6.0 >6.0	>6.0 >6.0 >6.0	--- Apparent ---	---	---	---	---	None None None
112A: Cowden-----	D	Jan-May Jun-Dec	0.0-1.0 >6.0	>6.0 >6.0	Apparent ---	0.0-0.5 ---	Brief ---	Frequent ---	---	None None

Table 32.--Water Features--Continued

Map symbol and soil name	Hydro- logic group	Month	Water table			Ponding			Flooding	
			Upper limit	Lower limit	Kind	Surface water depth	Duration	Frequency	Duration	Frequency
			<i>Ft</i>	<i>Ft</i>		<i>Ft</i>				
630C2: Navlys-----	B	Jan	>6.0	>6.0	---	---	---	---	---	None
		Feb-Apr	4.0-6.0	>6.0	Apparent	---	---	---	---	None
		May-Dec	>6.0	>6.0	---	---	---	---	---	None
630D2: Navlys-----	B	Jan	>6.0	>6.0	---	---	---	---	---	None
		Feb-Apr	4.0-6.0	>6.0	Apparent	---	---	---	---	None
		May-Dec	>6.0	>6.0	---	---	---	---	---	None
630D3: Navlys-----	B	Jan	>6.0	>6.0	---	---	---	---	---	None
		Feb-Apr	4.0-6.0	>6.0	Apparent	---	---	---	---	None
		May-Dec	>6.0	>6.0	---	---	---	---	---	None
679B: Blackberry-----	B	Jan	>6.0	>6.0	---	---	---	---	---	None
		Feb-Apr	2.0-3.5	>6.0	Apparent	---	---	---	---	None
		May-Dec	>6.0	>6.0	---	---	---	---	---	None
684B: Broadwell-----	B	Jan-Dec	>6.0	>6.0	---	---	---	---	---	None
684C2: Broadwell-----	B	Jan-Dec	>6.0	>6.0	---	---	---	---	---	None
685B: Middletown-----	B	Jan-Dec	>6.0	>6.0	---	---	---	---	---	None
685C2: Middletown-----	B	Jan-Dec	>6.0	>6.0	---	---	---	---	---	None
705A: Buckhart-----	B	Jan	>6.0	>6.0	---	---	---	---	---	None
		Feb-Apr	2.0-3.5	>6.0	Apparent	---	---	---	---	None
		May-Dec	>6.0	>6.0	---	---	---	---	---	None
705B: Buckhart-----	B	Jan	>6.0	>6.0	---	---	---	---	---	None
		Feb-Apr	2.0-3.5	>6.0	Apparent	---	---	---	---	None
		May-Dec	>6.0	>6.0	---	---	---	---	---	None
712A: Spaulding-----	B/D	Jan-May	0.0-1.0	>6.0	Apparent	0.0-0.5	Brief	Frequent	---	None
		Jun-Dec	>6.0	>6.0	---	---	---	---	---	None
801C: Orthents-----	B	Jan-Dec	>6.0	>6.0	---	---	---	---	---	None
830: Landfills-----	---	Jan-Dec	>6.0	>6.0	---	---	---	---	---	None
862: Pits, sand-----	---	Jan-Dec	>6.0	>6.0	---	---	---	---	---	---
864: Pits, quarries----	---	Jan-Dec	>6.0	>6.0	---	---	---	---	---	---

Table 32.--Water Features--Continued

Map symbol and soil name	Hydro- logic group	Month	Water table			Ponding			Flooding	
			Upper limit	Lower limit	Kind	Surface water depth	Duration	Frequency	Duration	Frequency
			<i>Ft</i>	<i>Ft</i>		<i>Ft</i>				
7242A: Kendall-----	B	Jan-May	0.5-2.0	>6.0	Apparent	---	---	---	---	Rare
		Jun	>6.0	>6.0	---	---	---	---	---	Rare
		Jul-Oct	>6.0	>6.0	---	---	---	---	---	None
		Nov-Dec	>6.0	>6.0	---	---	---	---	---	Rare
8396A: Vesser-----	C	Jan-May	0.0-1.0	>6.0	Apparent	---	---	---	Brief	Occasional
		Jun	>6.0	>6.0	---	---	---	---	Brief	Occasional
		Jul-Oct	>6.0	>6.0	---	---	---	---	---	None
		Nov-Dec	>6.0	>6.0	---	---	---	---	Brief	Occasional

Table 33.--Soil Features

(See text for definitions of terms used in this table. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

Map symbol and soil name	Restrictive layer		Potential for frost action	Risk of corrosion	
	Kind	Depth to top		Uncoated steel	Concrete
		In			
8D: Hickory-----	---	>80	Moderate----	Moderate----	Moderate.
8D2: Hickory-----	---	>80	Moderate----	Moderate----	Moderate.
8D3: Hickory-----	---	>80	Moderate----	Moderate----	Moderate.
8F: Hickory-----	---	>80	Moderate----	Moderate----	Moderate.
17A: Keomah-----	---	>80	High-----	High-----	Moderate.
36B: Tama-----	---	>80	High-----	Moderate----	Moderate.
36C2: Tama-----	---	>80	High-----	Moderate----	Moderate.
43A: Ipava-----	---	>80	High-----	High-----	Moderate.
45A: Denny-----	---	>80	High-----	High-----	Moderate.
46A: Herrick-----	---	>80	High-----	High-----	High.
50A: Virden-----	---	>80	High-----	High-----	Moderate.
68A: Sable-----	---	>80	High-----	High-----	Low.
86B: Osco-----	---	>80	High-----	Moderate----	Moderate.
86C2: Osco-----	---	>80	High-----	Moderate----	Moderate.
112A: Cowden-----	---	>80	High-----	High-----	Moderate.
119C2: Elco-----	---	>80	High-----	High-----	Moderate.
119D: Elco-----	---	>80	High-----	High-----	Moderate.
119D2: Elco-----	---	>80	High-----	High-----	Moderate.
119D3: Elco-----	---	>80	High-----	High-----	Moderate.

Table 33.--Soil Features--Continued

Map symbol and soil name	Restrictive layer		Potential for frost action	Risk of corrosion	
	Kind	Depth to top		Uncoated steel	Concrete
		In			
127B: Harrison-----	---	>80	High-----	High-----	Moderate.
127C2: Harrison-----	---	>80	High-----	High-----	Moderate.
131C2: Alvin-----	---	>80	Moderate---	Low-----	High.
131D2: Alvin-----	---	>80	Moderate---	Low-----	High.
134A: Camden-----	---	>80	High-----	Moderate---	Moderate.
134B: Camden-----	---	>80	High-----	Moderate---	Moderate.
134C2: Camden-----	---	>80	High-----	Low-----	Moderate.
138A: Shiloh-----	---	>80	High-----	High-----	Low.
198A: Elburn-----	---	>80	High-----	High-----	Moderate.
199A: Plano-----	---	>80	High-----	Moderate---	Low.
199B: Plano-----	---	>80	High-----	Moderate---	Low.
212C2: Thebes-----	---	>80	High-----	Moderate---	Moderate.
242A: Kendall-----	---	>80	High-----	High-----	High.
244A: Hartsburg-----	---	>80	High-----	High-----	Low.
249A: Edinburg-----	---	>80	High-----	High-----	Moderate.
257A: Clarksdale-----	---	>80	High-----	High-----	Moderate.
259C2: Assumption-----	---	>80	High-----	High-----	Moderate.
259D2: Assumption-----	---	>80	High-----	High-----	Moderate.
279B: Rozetta-----	---	>80	High-----	Moderate---	Moderate.
280C2: Fayette-----	---	>80	High-----	Moderate---	Moderate.
280D2: Fayette-----	---	>80	High-----	Moderate---	Moderate.

Table 33.--Soil Features--Continued

Map symbol and soil name	Restrictive layer		Potential for frost action	Risk of corrosion	
	Kind	Depth to top		Uncoated steel	Concrete
		In			
280D3: Fayette-----	---	>80	High-----	Moderate----	Moderate.
533: Urban land-----	---	>80	---	---	---
536: Dumps-----	---	>80	---	---	---
549G: Marseilles-----	Bedrock (paralithic)	20-40	High-----	High-----	Moderate.
567C2: Elkhart-----	---	>80	High-----	Moderate----	Moderate.
567D2: Elkhart-----	---	>80	High-----	Moderate----	Moderate.
630C2: Navlys-----	---	>80	High-----	Moderate----	Moderate.
630D2: Navlys-----	---	>80	High-----	Moderate----	Moderate.
630D3: Navlys-----	---	>80	High-----	Moderate----	Moderate.
679B: Blackberry-----	---	>80	High-----	High-----	Moderate.
684B: Broadwell-----	---	>80	High-----	Moderate----	Moderate.
684C2: Broadwell-----	---	>80	High-----	Moderate----	Moderate.
685B: Middletown-----	---	>80	High-----	High-----	High.
685C2: Middletown-----	---	>80	High-----	High-----	High.
705A: Buckhart-----	---	>80	High-----	Moderate----	Moderate.
705B: Buckhart-----	---	>80	High-----	Moderate----	Moderate.
712A: Spaulding-----	---	>80	High-----	High-----	Low.
801C: Orthents-----	---	>80	High-----	High-----	Moderate.
830: Landfills-----	---	>80	---	---	---
862: Pits, sand-----	---	>80	---	---	---

Table 33.--Soil Features--Continued

Map symbol and soil name	Restrictive layer		Potential for frost action	Risk of corrosion	
	Kind	Depth to top		Uncoated steel	Concrete
		In			
864: Pits, quarries-----	---	>80	---	---	---
3073A: Ross-----	---	>80	Moderate----	Low-----	Low.
3074A: Radford-----	---	>80	High-----	High-----	Moderate.
3077A: Huntsville-----	---	>80	High-----	Low-----	Low.
3107A: Sawmill-----	---	>80	High-----	High-----	Low.
3284A: Tice-----	---	>80	High-----	High-----	Low.
3405A: Zook-----	---	>80	High-----	High-----	Moderate.
3451A: Lawson-----	---	>80	High-----	Moderate----	Low.
7037A: Worthen-----	---	>80	High-----	Low-----	Low.
7075B: Drury-----	---	>80	High-----	Moderate----	Moderate.
7148A: Proctor-----	---	>80	High-----	Moderate----	Moderate.
7242A: Kendall-----	---	>80	High-----	High-----	Moderate.
8396A: Vesser-----	---	>80	High-----	High-----	Moderate.

NRCS Accessibility Statement

The Natural Resources Conservation Service (NRCS) is committed to making its information accessible to all of its customers and employees. If you are experiencing accessibility issues and need assistance, please contact our Helpdesk at 1-800-457-3642 or email at helpdesk@helpdesk.itc.nrcs.usda.gov. For assistance with publications that include maps, graphs, or similar forms of information, you may also wish to contact the State or local office. You can locate the correct office and phone number at http://offices.usda.gov/scripts/ndISAPI.dll/oip_public/USA_map.


USDA Nondiscrimination Statement

The United States Department of Agriculture (USDA) prohibits discrimination in all of its programs on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice or TDD).


To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue SW, Washington, DC 20250-9410, or call 202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Printing Soil Survey Maps


The soil survey maps were made at a scale of 1:12000 and were designed to be used at that scale. To print the maps at 1:12000 scale, set the view to Actual Size from the View pull down menu.




Using the pan tool, go to the area you would like to print. Select the Graphic Selection Tool by holding down the Text Selection Tool button and clicking on the Graphic Selection Tool button.








Then using the Graphic Selection Tool drag a box around the area you would like to print. Note dashed lines forming a box around area to print.



Select File Print. The Print Range will be set to Selected graphic. Click OK and the map will be sent to the printer.



CONVENTIONAL AND SPECIAL SYMBOLS LEGEND

DESCRIPTION	SYMBOL	DESCRIPTION	SYMBOL	DESCRIPTION	SYMBOL
CULTURAL FEATURES		CULTURAL FEATURES (cont.)		SPECIAL SYMBOLS FOR SOIL SURVEY AND SSURGO	
BOUNDARIES		MISCELLANEOUS CULTURAL FEATURES		 <p>SOIL DELINEATIONS AND SYMBOLS</p> <p>LANDFORM FEATURES</p> <p>ESCARPMENTS</p> <p>Bedrock </p> <p>Other than bedrock </p> <p>SHORT STEEP SLOPE </p> <p>GULLY </p> <p>DEPRESSION, closed </p> <p>SINKHOLE </p>	
• National, state, or province		Farmland, house (omit in urban areas)			
• County or parish		Church			
Minor civil division		School			
Reservation, (national forest or park, state forest or park)		Other Religion (label)			
Land grant		Located object (label)			
Limit of soil survey (label) and/or denied access areas		Tank (label)			
• Field sheet matchline & neatline		Lookout Tower			
Previously published survey		Oil and / or Natural Gas Wells			
OTHER BOUNDARY (label)		Windmill			
Airport, airfield		Lighthouse			
• Cemetery		HYDROGRAPHIC FEATURES			
City / county Park		STREAMS			
STATE COORDINATE TICK		Perennial, double line			
• LAND DIVISION CORNERS (section and land grants)		Perennial, single line			
• GEOGRAPHIC COORDINATE TICK		Intermittent			
TRANSPORTATION		Drainage end			
Divided roads		DRAINAGE AND IRRIGATION			
Other roads		Double line canal (label)			
# Trails		Perennial drainage and/or irrigation ditch			
ROAD EMBLEMS & DESIGNATIONS		Intermittent drainage and/or irrigation ditch			
• Interstate		SMALL LAKES, PONDS, AND RESERVOIRS			
• Federal		Perennial water			
• State		Miscellaneous water			
County, farm, or ranch		Flood pool line			
RAILROAD		MISCELLANEOUS WATER FEATURES			
POWER TRANSMISSION LINE (normally not shown)		Spring			
PIPELINE (normally not shown)		Well, artesian			
FENCE (normally not shown)		Well, irrigation			
LEVEES		RECOMMENDED AD HOC SOIL SYMBOLS			
Without road					
With road					
With railroad					
••••• Single side slope (showing actual feature location)					
DAMS					
Medium or small					
LANDFORM FEATURES					
Prominent Hill or Peak					
Soil Sample Site					

* Cultural features for use in Illinois

Descriptions of Special Features

Name	Description	Label
Blowout	A small saucer, cup, or trough-shaped hollow or depression formed by wind erosion, on a preexisting sand deposit. Typically 0.2 acre to 2.0 acres.	BLO
Borrow pit	An open excavation from which soil and underlying material have been removed, usually for construction purposes. Typically 0.2 acre to 2.0 acres.	BPI
Calcareous spot	An area in which the soil contains carbonates in the surface layer. The surface layer of the named soils in the surrounding map unit is non-calcareous. Typically 0.5 to 2.0 acres in size.	CSP
Clay spot	A spot where the surface layer is silty clay or clay in areas where the surface layer of the soils in the surrounding map unit is sandy loam, loam, silt loam, or coarser. Typically 0.2 acre to 2.0 acres.	CLA
Depression, closed	A shallow, saucer-shaped area that is slightly lower on the landscape than the surrounding area and is without a natural outlet for surface drainage. Typically 0.2 acre to 2.0 acres.	DEP
Disturbed soil spot	An area in which the soil has been removed and materials redeposited due to human activity. Typically 0.25 to 2.0 acres.	DSS
Dumps	Non-soil material that supports little or no vegetation. Typically 0.5 to 2.0 acres.	DMP
Escarpment, bedrock	A relatively continuous and steep slope or cliff, produced by erosion or faulting, that breaks the general continuity of more gently sloping land surfaces. Exposed material is hard or soft bedrock.	ESB
Escarpment, nonbedrock	A relatively continuous and steep slope or cliff, generally produced by erosion but in some places produced by faulting, that breaks the continuity of more gently sloping land surfaces. Exposed earthy material is nonsoil or very shallow soil.	ESO
Glacial till	An exposure of glacial till at the surface of the earth. Typically 0.25 to 2.0 acres.	GLA
Gravel pit	An open excavation from which soil and underlying material have been removed and used, without crushing, as a source of sand or gravel. Typically 0.2 acre to 2.0 acres.	GPI
Gravelly spot	A spot where the surface layer has more than 35 percent, by volume, rock fragments that are mostly less than 3 inches in diameter in an area with less than 15 percent rock fragments. Typically 0.2 acre to 2.0 acres.	GRA
Gully	A small channel with steep sides cut by running water through which water ordinarily runs only after a rain or after melting of ice or snow. It generally is an obstacle to wheeled vehicles and is too deep to be obliterated by ordinary tillage.	GUL
Iron bog	An accumulation of iron in the form of nodules, concretions, or soft masses on the surface or near the surface of soils. Typically 0.2 acre to 2.0 acres.	IBA
Landfill	An area of accumulated waste products of human habitation, either above or below natural ground level. Typically 0.2 acre to 2.0 acres.	LDL
Levee	An embankment that confines or controls water, especially one built along the LVS banks of a river to prevent overflow onto lowlands.	LVS

Name	Description	Label
Marsh or swamp	A water-saturated, very poorly drained area, intermittently or permanently covered by water. Sedges, cattails, and rushes are the dominant vegetation in marsh areas, and trees or shrubs are the dominant vegetation in swamps. Typically 0.2 acre to 2.0 acres.	MAR
Mine or quarry	An open excavation from which soil and underlying material have been removed and in which bedrock is exposed. Also denotes surface openings to underground mines. Typically 0.2 acre to 2.0 acres.	MPI
Mine subsided area	Areas that are lower than the soils in the surrounding map units due to subsurface coal mining. Typically 0.25 to 3 acres.	MSA
Miscellaneous water	A small, constructed water area that is used for industrial, sanitary, or mining applications and contains water most of the year. Typically 0.2 acre to 2.0 acres.	MIS
Muck spot	An area that occurs within an area of poorly drained or very poorly drained soil and that has a histic epipedon or an organic surface layer. The symbol is used only in map units consisting of mineral soil. Typically 0.2 acre to 2.0 acres.	MUC
Oil brine spot	An area of soil that has been severely damaged by the accumulation of oil brine, with or without liquid oily wastes. The area is typically barren but may have a vegetative cover of salt-tolerant plants. Typically 0.2 acre to 2.0 acres.	OBS
Perennial water	A small, natural or constructed lake, pond, or pit that contains water most of the year. Typically 0.2 acre to 2.0 acres.	WAT
Rock outcrop	An exposure of bedrock at the surface of the earth. Not used where the named soils of the surrounding map unit are shallow over bedrock or where "Rock outcrop" is a named component of the map unit. Typically 0.2 acre to 2.0 acres.	ROC
Saline spot	An area where the surface layer has an electrical conductivity of 8 mmhos/cm-1 more than the surface layer of the named soils in the surrounding map unit, in which electrical conductivity is 2 mmhos/cm-1 or less. Typically 0.2 acre to 2.0 acres.	SAL
Sandy spot	A spot where the surface layer is loamy fine sand or coarser in areas where the surface layer of the named soils in the surrounding map unit is very fine sandy loam or finer. Typically 0.2 acre to 2.0 acres.	SAN
Severely eroded spot	An area where, on the average, 75 percent or more of the original surface layer has been lost because of accelerated erosion. Not used in map units in which "severely eroded," "very severely eroded," or "gullied" is part of the map unit name. Typically 0.2 acre to 2.0 acres	ERO
Short, steep slope	A narrow area of soil having slopes that are at least two slope classes steeper than the slope class of the surrounding map unit.	SLP
Sinkhole	A closed depression formed either by solution of the surficial rock or by collapse of underlying caves. Typically 0.2 acre to 2.0 acres.	SNK

Name	Description	Label
Slide or slip	A prominent landform scar or ridge caused by fairly recent mass movement or descent of earthy material resulting from failure of earth or rock under shear stress along one or several surfaces. Typically 0.2 acre to 2.0 acres	SLI
Sodic spot	An area where the surface layer has a sodium adsorption ratio that is at least 10 more than that of the surface layer of the named soils in the surrounding map unit, which have a sodium adsorption ratio of 5 or less. Typically 0.2 acre to 2.0 acres.	SOD
Spoil area	A pile of earthy materials, either smoothed or uneven, resulting from human activity. Typically 0.2 acre to 2.0 acres.	SPO
Stony spot	A spot where 0.01 to 0.1 percent of the surface cover is rock fragments that are more than 10 inches in diameter in areas where the surrounding soil has no surface stones. Typically 0.2 acre to 2.0 acres.	STN
Unclassified water	A small, natural or manmade lake, pond, or pit that contains water, of an unspecified nature, most of the year. Typically 0.2 acre to 2.0 acres.	UWT
Very stony spot	A spot where 0.1 to 3.0 percent of the surface cover is rock fragments that are more than 10 inches in diameter in areas where the surface cover of the surrounding soil is less than 0.01 percent stones. Typically 0.2 acre to 2.0 acres.	STV
Wet depression	A shallow, concave area within an area of poorly drained or very poorly drained soils in which water is ponded for intermittent periods. The concave area is saturated for appreciably longer periods of time than the surrounding soil. Typically 0.2 acre to 2.0 acres.	WDP
Wet spot	A somewhat poorly drained to very poorly drained area that is at least two drainage classes wetter than the named soils in the surrounding map unit. Typically 0.2 acre to 2.0 acres.	WET

Sangamon County, Illinois Index to atlas sheets.

Click on a blue number to view soil map of area

